

# Problem set 7

Saturday, November 12, 2016 12:27 AM

1. Suppose  $X$  be a non-empty set. For any subset  $A$  of  $X$ , let  $\mathbb{1}_A$  be the characteristic function of  $A$ . That means,

$$\mathbb{1}_A: X \rightarrow \{0, 1\}, \quad \mathbb{1}_A(x) = \begin{cases} 1 & \text{if } x \in A, \\ 0 & \text{if } x \notin A. \end{cases}$$

(a) Suppose  $X$  is finite. What is  $\sum_{x \in X} \mathbb{1}_A(x)$ ?

(b) Use part (a) and  $\mathbb{1}_{A \cup B} = \mathbb{1}_A + \mathbb{1}_B - \mathbb{1}_{A \cap B}$  to conclude

$$|A \cup B| = |A| + |B| - |A \cap B|.$$

(c) Suppose  $A_1, \dots, A_n$  are subsets of  $X$ . By induction on  $n$ ,

show that  $\mathbb{1}_{(A_1 \cup A_2 \cup \dots \cup A_n)^c} = (1 - \mathbb{1}_{A_1})(1 - \mathbb{1}_{A_2}) \dots (1 - \mathbb{1}_{A_n})$ .

(Hint. Use  $(A \cup B)^c = A^c \cap B^c$ ,  $\mathbb{1}_{A^c} = 1 - \mathbb{1}_A$ ,  $\mathbb{1}_{A \cap B} = \mathbb{1}_A \cdot \mathbb{1}_B$ .)

(d) Use part (c) to conclude

$$\begin{aligned} \mathbb{1}_{(A_1 \cup \dots \cup A_n)^c} &= 1 - (\mathbb{1}_{A_1} + \dots + \mathbb{1}_{A_n}) + \underbrace{(\mathbb{1}_{A_1 \cap A_2} + \dots + \mathbb{1}_{A_{n-1} \cap A_n})}_{\text{all } \mathbb{1}_{A_i \cap A_j}} \\ &\quad - \underbrace{(\mathbb{1}_{A_1 \cap A_2 \cap A_3} + \dots + \mathbb{1}_{A_{n-2} \cap A_{n-1} \cap A_n})}_{\text{all } \mathbb{1}_{A_i \cap A_j \cap A_k}} + \dots \end{aligned}$$

(signs alternate)  $+ (-1)^n \mathbb{1}_{A_1 \cap A_2 \cap \dots \cap A_n}$ .

Then, using (a), deduce

$$|(A_1 \cup A_2 \cup \dots \cup A_n)^c| = |X| - \sum_i |A_i| + \sum_{i_1 < i_2} |A_{i_1} \cap A_{i_2}| - \dots + (-1)^n |A_1 \cap \dots \cap A_n|.$$

(This is called inclusion-exclusion formula).

# Problem set 7

Saturday, November 12, 2016 12:59 AM

2.(a) Suppose  $f: X \rightarrow Y$  is a function. Prove that if

$g: Y \rightarrow X$  is a left inverse of  $f$  and  $h: Y \rightarrow X$  is a right inverse of  $f$ , then  $g = h$ .

Hint ① You are allowed to use the fact that for three functions  $X_1 \xrightarrow{f_1} X_2 \xrightarrow{f_2} X_3 \xrightarrow{f_3} X_4$  we have  $f_3 \circ (f_2 \circ f_1) = (f_3 \circ f_2) \circ f_1$ .

② Consider  $g \circ f \circ h$ .)

(b) Use part (a) and a theorem proved in class to show:

If a function  $X \xrightarrow{f} Y$  is a bijection, then there is a unique function  $Y \xrightarrow{g} X$  such that  $g \circ f = I_X$  and  $f \circ g = I_Y$ .

(This function is called the inverse of  $f$ , and it is denoted by  $f^{(-1)}$ .)

3. Suppose  $X \xrightarrow{f} Y$  and  $Y \xrightarrow{g} Z$  are two bijections.

(a) Prove that  $g \circ f$  is a bijection.

(b) Prove that  $f^{(-1)}$  is a bijection. ( $f^{(-1)}$  is as above.)

4. Suppose  $X \xrightarrow{f} Y$  and  $Y \xrightarrow{g} Z$  are two functions, and  $g \circ f$  is a bijection. Prove that  $g$  is injective  $\iff f$  is surjective.

Hint.  $g$  bijective  $\implies f = \underbrace{g^{(-1)}}_{\text{bijective}} \circ \underbrace{(g \circ f)}_{\text{bijective}}$   
 $f$  bijective  $\implies g = \underbrace{(g \circ f)}_{\text{bijective}} \circ \underbrace{f^{(-1)}}_{\text{bijective}}$  } Use problem 3 and a theorem from class.)

## Problem set 7

Saturday, November 12, 2016 1:44 AM

This is NOT part of your homework assignment. It is for your practice.

5. Suppose  $X \xrightarrow{f} Y$  is a function. Suppose  $|X| \geq 2$ .

Prove that

(a)  $f$  has a unique left inverse  $\iff f$  is bijective.

(b)  $f$  has a unique right inverse  $\iff f$  is bijective.

(Hint. Look at the way we proved

$f$  has a left inverse  $\iff f$  is injective

and

$f$  has a right inverse  $\iff f$  is surjective,

assume is not bijective and find more than 1 left or right inverse in each case.)