

FOURIER EXPANSIONS OF COMPLEX-VALUED EISENSTEIN SERIES ON FINITE UPPER HALF PLANES

ANTHONY SHAHEEN AND AUDREY TERRAS

ABSTRACT. We consider complex-valued modular forms on finite upper half planes H_q and obtain Fourier expansions of Eisenstein series invariant under the groups $\Gamma = SL(2, \mathbb{F}_p)$ and $GL(2, \mathbb{F}_p)$. The expansions are analogous to those of Maass wave forms on the ordinary Poincaré upper half plane - the K-Bessel functions being replaced by Kloosterman sums.

1. INTRODUCTION

Before outlining our results, let us give a brief summary of the classical results for which we have found finite analogs. This work is part of a continuing project to seek out finite analogs of Terras [16], Chapter 3.

The usual Poincaré upper half plane H consists of complex numbers $z = x + iy$ with $y > 0$. The Poincaré arc length is defined by $ds^2 = y^{-2} (dx^2 + dy^2)$ and the corresponding Laplacian is $\Delta = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$. More information about H , its non-Euclidean geometry, and modular forms can be found in Terras [16].

Maass wave forms (of weight 0) for $\Gamma = SL(2, \mathbb{Z})$ are functions $f : H \rightarrow \mathbb{C}$ which are eigenfunctions of the Poincaré Laplacian $\Delta = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$ such that $f(\gamma z) = f\left(\frac{az+b}{cz+d}\right) = f(z)$ for all $z \in H$ and $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ and with $f(z)$ growing at most as a polynomial in y as $y \rightarrow \infty$.

An example of a Maass wave form for $SL(2, \mathbb{Z})$ is provided by the **Maass Eisenstein series** on H defined, for $z \in H$ and $s \in \mathbb{C}$ with $\text{Re } s > 1$, by

$$(1.1) \quad E_s(z) = \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} \text{Im}(\gamma z)^s.$$

Here Γ_∞ is the subgroup of Γ consisting of matrices which fix ∞ (i.e., the lower left element is 0).

Define the **gamma function** for $\text{Re } s > 0$ by

$$(1.2) \quad \Gamma(s) = \int_0^\infty y^s e^{-y} \frac{dy}{y}.$$

Next define the **K-Bessel function** for $\text{Re } a > 0$ by

$$(1.3) \quad K_s(a) = \frac{1}{2} \int_0^\infty y^s e^{-\frac{a}{2}\left(y+\frac{1}{y}\right)} \frac{dy}{y}.$$

Set

$$\Lambda(s) = \pi^{-s} \Gamma(s) \zeta(2s),$$

where $\zeta(s)$ is Riemann's zeta function. Then the **Fourier expansion** of $E_s(z)$ is

$$(1.4) \quad y^s + \frac{\Lambda(1-s)}{\Lambda(s)} y^{1-s} + \frac{2}{\Lambda(s)} \sum_{n \neq 0} |n|^{s-\frac{1}{2}} \sigma_{1-2s}(n) y^{\frac{1}{2}} K_{s-\frac{1}{2}}(2\pi |n| y) e^{2\pi i n x}.$$

Here the **divisor function** is

$$\sigma_s(n) = \sum_{0 < d|n} d^s.$$

See Terras [16] pp. 208-9 for a sketch of three proofs.

Date: July 20, 2006.

1991 Mathematics Subject Classification. 11T60, 11T23, 11L05, 11F99.

Key words and phrases. finite upper half planes, Eisenstein series, Kloosterman sums.

The upper half plane H has a finite analog called the **finite “upper” half plane** H_q attached to a finite field \mathbb{F}_q with q elements. We assume q is an odd number. It is likely that most of our results go through in characteristic 2, but we leave this for a future paper. Then we take a fixed non-square $\delta \in \mathbb{F}_q$, and define

$$H_q = \left\{ z = x + y\sqrt{\delta} \mid x, y \in \mathbb{F}_q, y \neq 0 \right\}.$$

The finite upper half plane is considered in detail in Terras [17], Chapter 19. See also Shaheen [13]. An application to coding theory is to be found in Tiu and Wallace [19]. References which cover finite upper half planes in both even and odd characteristic include: Angel [1], Angel and Evans [3], Evans [8], as well as Soto-Andrade and Vargas [15].

Recall that an element of the **general linear group**

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL(2, \mathbb{F}_q)$$

has entries in \mathbb{F}_q and non-zero determinant. Then for $z \in H_q$

$$gz = \frac{az + b}{cz + d} \in H_q.$$

Define the **“distance”** between $z, w \in H_q$ by

$$(1.5) \quad d(z, w) = \frac{N(z - w)}{\text{Im } z \text{ Im } w}.$$

It is easily checked that for $g \in GL(2, \mathbb{F}_q)$, $d(gw, gw) = d(z, w)$. For $a \in \mathbb{F}_q$, define the **finite upper half plane graphs** $X_q(\delta, a)$ to have vertices the elements of H_q and an edge between z and w in H_q iff $d(z, w) = a$. For $a \neq 0, 4\delta$, the graph $X_q(\delta, a)$ is $(q + 1)$ -regular, connected, and Ramanujan. See Terras [17] for more information. Ramanujan graphs are of interest to computer scientists because they provide efficient communications networks. Chai and Li [7] have proved some interesting connections of the spectra of the finite upper half plane graphs and modular forms of function fields.

The $GL(2, \mathbb{F}_q)$ -invariant operators on H_q analogous to the non-Euclidean Laplacian on the Poincaré upper half plane H are the **adjacency operators** on the finite upper half plane graphs defined for $a \in \mathbb{F}_q$ by

$$(1.6) \quad A_a f(z) = \sum_{\substack{w \in H_q \\ d(z, w) = a}} f(w).$$

These operators generate a commutative algebra of operators.

One can view H_q as a finite symmetric space G/K as in Terras [17], Chapter 19. From that point of view, the adjacency operators A_a correspond to G -invariant differential operators on a symmetric space. This subject may also be studied from the point of view of association schemes (see Bannai and Ito [4]) or Hecke operators (see Krieg [11]).

A (complex-valued) **modular form** on the finite upper half plane H_q is a function $f : H_q \rightarrow \mathbb{C}$ having an invariance property for a subgroup Γ of $GL(2, \mathbb{F}_q)$. See Definition 3 below. Here we consider $\Gamma = SL(2, \mathbb{F}_p)$ and $GL(2, \mathbb{F}_p)$. Such modular forms which are eigenfunctions of the adjacency operators for the finite upper half plane graphs attached to H_q are analogs of Maass wave forms on the Poincaré upper half plane.

Now we consider one of our finite upper half plane analogs of the Maass Eisenstein series in formula (1.1). Let Γ be a subgroup of $GL(2, \mathbb{F}_q)$ and let χ be a multiplicative character on \mathbb{F}_q^* . Define the **Eisenstein “series”** for $z \in H_q$ as

$$(1.7) \quad E_{\chi, \Gamma}(z) = \sum_{\gamma \in \Gamma} \chi(\text{Im}(\gamma z)).$$

We will see that $E_{\chi, \Gamma}$ is an eigenfunction of the adjacency operators of finite upper half plane graphs if it does not vanish identically on H_q . To see when this happens, recall that \mathbb{F}_q^* is a cyclic group with generator g . (See Terras [17] and the references given there for more information about finite fields.) Thus a **multiplicative character** χ of \mathbb{F}_q^* has the following form for integers a, b :

$$(1.8) \quad \chi(g^b) = e^{\frac{2\pi i ab}{q-1}}, \quad 0 \leq a, b \leq q-2.$$

We will find that for $q = p^n$, with $q > 2$ and $n > 2$, $E_{\chi, GL(2, \mathbb{F}_q)}$ is non-zero if and only if $\chi = \chi_a$ when a is a multiple of $(p-1)$. This last condition is equivalent to saying that $\chi_a|_{\mathbb{F}_p^*} = 1$. When $n = 2$, we will find that $E_{\lambda, GL(2, \mathbb{F}_q)}$ is non-zero, for $\chi = 1$ or $\chi = \lambda$ as in formula (1.15) below. We will also find experimentally that $E_{\chi, GL(2, \mathbb{F}_q)}$ is non-zero for $p = 3, 5$ when a is a multiple of $(p-1)$ (see Tables 1 and 2 at the end of this paper).

As we saw in formula (1.4), Fourier expansions of Maass Eisenstein series on H involve K-Bessel functions. In the finite case under consideration the Bessel functions will be replaced with Kloosterman sums. Thus we need to define these sums as well as Gauss sums in order to state the Fourier expansion of the Eisenstein series $E_{\chi, GL(2, \mathbb{F}_p)}(z)$.

Let Ψ be an **additive character** of \mathbb{F}_q . Then $\Psi = \Psi_b$ where

$$(1.9) \quad \Psi_b(u) = e^{\frac{2\pi i \text{Tr}(bu)}{p}}, \quad \text{for } b, u \in \mathbb{F}_q.$$

Here the **trace** in the exponent is

$$(1.10) \quad \text{Tr}(u) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(u) = u + u^p + u^{p^2} + \cdots + u^{p^{n-1}}.$$

Let Ψ be an additive character and χ a multiplicative character. Define the **Gauss sum** by

$$(1.11) \quad \Gamma(\chi, \Psi) = \Gamma_q(\chi, \Psi) = \sum_{t \in \mathbb{F}_q^*} \chi(t) \Psi(t).$$

The Gauss sum is analogous to the gamma function of formula (1.2).

See Ireland and Rosen [10] for more information on Gauss sums. In particular, they show that, if χ and Ψ_b are not trivial, then

$$(1.12) \quad \Gamma(\chi, \Psi_b) = \chi(b^{-1}) \Gamma(\chi, \Psi_1).$$

Another basic result says that if χ and Ψ are not trivial, then

$$(1.13) \quad |\Gamma_q(\chi, \Psi)| = \sqrt{q}.$$

The **Kloosterman sum** for $a, b \in \mathbb{F}_q$ is defined by

$$(1.14) \quad K_\Psi(\chi|a, b) = \sum_{t \in \mathbb{F}_q^*} \chi(t) \Psi(at + bt^{-1}).$$

The Kloosterman sum is analogous to the K-Bessel function of formula (1.3). These sums have great importance in number theory. See Sarnak [12] who discusses the connection with modular forms and ‘‘Kloostermania’’.

Define the **quadratic residue character** λ on \mathbb{F}_q^* by

$$(1.15) \quad \lambda(y) = \begin{cases} 1, & \text{if } y \text{ is a square in } \mathbb{F}_q^*; \\ -1, & \text{if } y \text{ is not a square in } \mathbb{F}_q^*; \\ 0, & y = 0. \end{cases}$$

One of our main results (see formula (2.8)) is a finite analog of formula (1.4) and says that the **Fourier expansion of the Eisenstein series** $E_{\chi, GL(2, \mathbb{F}_q)}(z)$ is given by

$$(1.16) \quad \frac{1}{p(p-1)^2} E_{\chi, GL(2, \mathbb{F}_q)}(z) = \chi(y) + \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1) \Gamma(\lambda \chi, \Psi_1)}{\Gamma(\chi, \Psi_1)} \chi^{-1}(-\delta y) \lambda(-\delta) \\ + \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1)}{\Gamma(\chi, \Psi_1)} \chi(y) \sum_{\substack{b \in \mathbb{F}_q^* \\ \text{Tr}(b)=0}} \frac{\chi(b)}{\lambda(b)} K_{\Psi_b} \left(\lambda \chi \left| -\delta y^2, -\frac{1}{4} \right. \right) \Psi_b(-x).$$

We will also prove the analogous result for $\Gamma = SL(2, \mathbb{F}_p)$ in formula (2.7). Both of these Fourier expansions result from that of Theorem 1 which gives the Fourier expansion of what we call a G-type Eisenstein series (see Definition 2).

The main idea of the proof of (1.16) is to imitate the second proof of formula (1.4) given in Terras [16]. This proof goes back to Bateman and Grosswald [5] and is extended in Terras [18]. The main fact needed is the Poisson summation formula for \mathbb{Z} as a subgroup of \mathbb{R} . For our finite upper half plane analog we will use Poisson summation for the additive subgroup \mathbb{F}_p of \mathbb{F}_q . See Proposition 2 below or p. 199 of Terras [17].

One can also consider finite field valued modular forms on finite upper half planes. These are perhaps more analogous to holomorphic modular forms. See Shaheen [13] where an analog of the Selberg trace formula is developed which is more like that of Selberg (see Hejhal [9]) than the version in Terras [17].

2. LATTICES, MODULAR FORMS ON H_q , AND FOURIER EXPANSIONS

Classically Eisenstein series are sums over lattices such as $\mathbb{Z}w_1 \oplus \mathbb{Z}w_2$ for elements w_i in \mathbb{C} which are linearly independent over \mathbb{R} . Our finite analog of a lattice is defined as follows. Here $q = p^n$, with $n > 1$.

Definition 1. An \mathbb{F}_p -lattice in $\mathbb{C}_q = \mathbb{F}_q(\sqrt{\delta})$ is a set of the form $L = \mathbb{F}_p w_1 \oplus \mathbb{F}_p w_2$, where $w_1, w_2 \in \mathbb{C}_q$ are linearly independent over \mathbb{F}_q .

Note that if $z \in H_q$, we have an example of an \mathbb{F}_p -lattice in $\mathbb{C}_q = \mathbb{F}_q(\sqrt{\delta})$ by setting $L_z = \mathbb{F}_p z \oplus \mathbb{F}_p$. Next we define a G-type Eisenstein series.




FIGURE 1. A Tessellation of H_{49} from $GL(2, \mathbb{F}_7)$. Any 7 points of different colors form a fundamental domain for $GL(2, \mathbb{F}_7) \backslash H_{49}$.

Definition 2. Let $\chi : \mathbb{F}_q(\sqrt{\delta})^* \rightarrow \mathbb{C}$ be a multiplicative character. Define the **G-Eisenstein series** $G_\chi(z)$, for $z \in H_q$, by

$$(2.1) \quad G_\chi(z) = \sum_{\substack{0 \neq \omega \\ \omega \in L_z = \mathbb{F}_p z \oplus \mathbb{F}_p}} \chi\left(\frac{1}{\omega}\right).$$

Proposition 1. If $q = p^n$, $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL(2, \mathbb{F}_p)$ and $z \in H_q$, then

$$G_\chi(\gamma z) = \chi(cz + d)G_\chi(z).$$

Proof. Imitate the classical proof found in [14]. □

We now give our most general definition of a complex-valued modular form.

Definition 3. Let $\chi : \mathbb{F}_q(\sqrt{\delta})^* \rightarrow \mathbb{C}$ be a multiplicative character. We say that $f : H_q \rightarrow \mathbb{C}$ is a **complex-valued modular form for χ and a subgroup Γ of $GL(2, \mathbb{F}_q)$** if

$$f(\gamma z) = \chi(cz + d)f(z) \text{ for all } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma \text{ and } z \in H_q.$$

We denote the complex vector space of such forms by $M_\chi(\Gamma)$.

Proposition 1 can then be restated to say that, as a function of z , $G_\chi(z) \in M_\chi(\Gamma)$. If $f \in M_\chi(\Gamma)$ and $g \in M_\psi(\Gamma)$, then $fg \in M_{\chi\psi}(\Gamma)$.

Note that $M_\chi(GL(2, \mathbb{F}_p)) \neq \{0\}$ implies that $\chi(a) = 1$ for all $a \in \mathbb{F}_p^*$. Similarly $M_\chi(SL(2, \mathbb{F}_p)) \neq \{0\}$ implies that $\chi(-1) = 1$. This is analogous to the fact that classical holomorphic modular forms for $SL(2, \mathbb{Z})$ must have even weight.

Fundamental domains for $\Gamma \backslash H_q$ are discussed in Shaheen [13]. It turns out that the classical perpendicular bisector method for constructing such fundamental domains (discussed in Terras [16], for example) has a finite analog.

Tessellations of H_q given by letting Γ act on a fundamental domain are as beautiful as those for the modular group acting on the Poincaré upper half plane. One such tessellation is given in Figure 1.

Let χ be a multiplicative character of \mathbb{F}_q^* and let $N = N_{\mathbb{F}_q(\sqrt{\delta})/\mathbb{F}_q}$ be the norm map. The character $\chi \circ N^{-1}$ seems like a reasonable character to investigate. In fact, it comes up when we consider the Eisenstein series $E_{\chi, \Gamma}$ from formula (1.7) of the introduction when $\Gamma = GL(2, \mathbb{F}_p)$ as we shall soon see.

Here we will need the Poisson summation formula for a subgroup B of a finite abelian group A proved in Terras [17], p. 199. The formula relates Fourier analysis on A/B and that on A itself.

Proposition 2. Poisson Summation Formula for Finite Abelian Groups.

Suppose B is a subgroup of the finite additive abelian group A and $f : A \rightarrow \mathbb{C}$. Then

$$(2.2) \quad \frac{1}{|B|} \sum_{b \in B} f(a+b) = \frac{1}{|A|} \sum_{\Psi \in B^\#} \widehat{f}(\Psi) \Psi(a).$$

We write \widehat{A} to denote the group of additive characters of A . Here the **dual group** to B in A is

$$B^\# = \{ \Psi \in \widehat{A} \mid \Psi(b) = 1, \forall b \in B \} \cong (\widehat{A/B}),$$

and the **Fourier transform of f** at $\Psi \in \widehat{A}$ is

$$\widehat{f}(\Psi) = \sum_{u \in A} f(u) \overline{\Psi}(u).$$

Note that the left hand side of formula (2.2) is a function of the coset $a+B$ in A/B while the right hand side involves Fourier transforms on A .

We want to apply the Poisson summation formula to $B = \mathbb{F}_p$ as an additive subgroup of $A = \mathbb{F}_q$. Therefore, we will need to know the characters in $\mathbb{F}_p^\#$. As in formula (1.9), $\Psi_a(u) = e^{\frac{2\pi i Tr(au)}{p}}$ is an arbitrary element of $\widehat{\mathbb{F}_q}$. Here the trace is $Tr = Tr_{\mathbb{F}_q/\mathbb{F}_p}$ as in formula (1.10). Then, $\Psi_a(u) = e^{\frac{2\pi i Tr(au)}{p}} = 1$ for all $u \in \mathbb{F}_p$, if and only if, p divides $Tr(au) = uTr(a)$ for all $u \in \mathbb{F}_p$. Therefore, $\Psi_a \in \mathbb{F}_p^\#$, if and only if, $Tr(a) = 0$ in \mathbb{F}_p . We have proved the following lemma.

Lemma 1. Using the notation of the preceding proposition, if we consider \mathbb{F}_p as a subgroup of the additive group \mathbb{F}_q , the dual group to \mathbb{F}_p in \mathbb{F}_q is

$$\mathbb{F}_p^\# = \left\{ \Psi_a \in \widehat{\mathbb{F}_q} \mid a \in \mathbb{F}_q, Tr(a) = Tr_{\mathbb{F}_q/\mathbb{F}_p}(a) = 0 \right\}.$$

We will also find that the Kloosterman sum appears in a different guise in our computations. So let us define the finite k -Bessel function in analogy to that considered on the Poincaré upper half plane in Terras [16], p. 136.

Definition 4. For a multiplicative character χ of \mathbb{F}_q^* and an additive character Ψ of \mathbb{F}_q , if $z \in H_q$, define the **k -Bessel function** by

$$k(z|\chi, \Psi) = \sum_{u \in \mathbb{F}_q} \chi \left(\operatorname{Im} \left(\frac{-1}{z+u} \right) \right) \Psi(u).$$

It follows from Lemma 2 below that $k(z) = k(z|\chi, \Psi)$ is an eigenfunction of all the adjacency operators A_a for the finite upper half plane graphs from formula (1.6).

The following proposition is proved in Celniker et al [6] (see also Angel et al [2]). It is a finite analog of a classical integral formula for K -Bessel functions.

Proposition 3. Relation Between k -Bessel function and Kloosterman sum.

For a non-trivial multiplicative character χ of \mathbb{F}_q^* and a non-trivial additive character Ψ_a of \mathbb{F}_q and $z \in H_q$,

$$\Gamma(\chi, \Psi_a) k(z|\chi, \Psi_a) = g_a \chi(y) \Psi_a(-x) K_{\Psi_a}(\chi \lambda | -\delta y^2, -\frac{1}{4}),$$

where $g_a = \sum_{u \in \mathbb{F}_q} \Psi_a(u^2)$ is a Gauss sum, the Kloosterman sum K is defined by formula (1.14), and the multiplicative character λ is given in formula (1.15).

Why do we say that g_a is a Gauss sum? It is easily seen that for $a \neq 0$

$$(2.3) \quad g_a = \sum_{u \in \mathbb{F}_q} \Psi_a(u^2) = \sum_{u \in \mathbb{F}_q} (1 + \lambda(u)) \Psi_a(u) = \lambda \left(a^{-1} \right) \Gamma(\lambda, \Psi_1).$$

Theorem 1. Let $N : \mathbb{F}_q(\sqrt{\delta})^* \rightarrow \mathbb{F}_q^*$ be the norm map, $\chi = \chi_a$ be the multiplicative character of \mathbb{F}_q^* given by (1.8), and $z \in H_q$. If $a = 0$, then $G_{\chi_0}(z) = p^2 - 1$. If $a \in \{ \frac{p-1}{2}, \frac{2(p-1)}{2}, \frac{3(p-1)}{2}, \frac{4(p-1)}{2}, \dots, \frac{m(p-1)}{2} \}$ where $\frac{m}{2} = \frac{q-1}{p-1} - 1$, then for $z = x + y\sqrt{\delta} \in H_q$, we have the Fourier expansion

$$\begin{aligned} \frac{1}{p-1} G_{\chi \circ N^{-1}}(z) &= 1 + \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1) \Gamma(\lambda \chi, \Psi_1)}{\Gamma(\chi, \Psi_1)} \chi^{-1}(-\delta y^2) \lambda^{-1}(-\delta) \\ &\quad + \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1)}{\Gamma(\chi, \Psi_1)} \sum_{\substack{0 \neq b \in \mathbb{F}_q \\ Tr(b)=0}} (\chi \lambda)(b) K_{\Psi_b} \left(\lambda \chi \left| -\delta y^2, -\frac{1}{4} \right. \right) \psi_b(-x). \end{aligned}$$

Here ψ_b is the additive character of \mathbb{F}_q given by (1.9), $\Gamma(\chi, \Psi)$ is the Gauss sum from formula (1.11), K_ψ is the Kloosterman sum from formula (1.14), λ is the multiplicative character from formula (1.15). If

$$a \notin \left\{0, \frac{p-1}{2}, \frac{2(p-1)}{2}, \frac{3(p-1)}{2}, \frac{4(p-1)}{2}, \dots, \frac{m(p-1)}{2}\right\},$$

then $G_{\chi \circ N^{-1}} = 0$.

Proof. First note that $a \in \left\{0, \frac{p-1}{2}, \frac{2(p-1)}{2}, \frac{3(p-1)}{2}, \frac{4(p-1)}{2}, \dots, \frac{m(p-1)}{2}\right\}$ is equivalent to $\chi_a^2 = \text{identity}$ on \mathbb{F}_p^* . To see this, suppose that $\mathbb{F}_q^* = \langle g \rangle$ and $\chi = \chi_a$ as in formula (1.8). It follows that $\mathbb{F}_p^* = \left\langle g^{\frac{q-1}{p-1}} \right\rangle$. Therefore, $\chi^2 = \text{identity}$ on \mathbb{F}_p^* if and only if $(p-1)$ divides $2a$.

Henceforth, we will assume that $\chi^2 = \text{identity}$ on \mathbb{F}_p^* . The case $\chi = \chi_0$ is trivial since G_χ is constant. Thus for the remainder of this proof we also assume that $\chi \neq \chi_0$.

If $z \in H_q$, then

$$\begin{aligned} G_{\chi \circ N^{-1}}(z) &= \sum_{\substack{(c,d) \in \mathbb{F}_p^2 \\ (c,d) \neq (0,0)}} \chi \left(\frac{1}{N(cz+d)} \right) = \sum_{d \in \mathbb{F}_p^*} \chi \left(\frac{1}{d^2} \right) + \sum_{c \in \mathbb{F}_p^*} \chi \left(\frac{1}{c^2} \right) \sum_{d \in \mathbb{F}_p} \chi^{-1} \left(N(z + \frac{d}{c}) \right) \\ &= \sum_{v \in \mathbb{F}_p^*} \chi^2(v) + \sum_{v \in \mathbb{F}_p^*} \chi^2(v) \sum_{u \in \mathbb{F}_p} \chi^{-1}(N(z+u)) \\ &= (p-1) \begin{cases} 0, & \text{if } \chi^2 \neq \text{identity on } \mathbb{F}_p^*; \\ 1 + \sum_{u \in \mathbb{F}_p} \chi^{-1}(N(z+u)), & \text{if } \chi^2 = \text{identity on } \mathbb{F}_p^*. \end{cases} \end{aligned}$$

We can now apply Poisson summation (from Proposition 2.2) and Lemma 1 to see that

$$\sum_{u \in \mathbb{F}_p} \chi^{-1}(N(z+u)) = \frac{p}{q} \sum_{\substack{0 \neq b \in \mathbb{F}_q \\ \text{Tr}(b)=0}} \sum_{u \in \mathbb{F}_q} \chi^{-1}(N(z+u)) \overline{\Psi}_b(u).$$

Expanding the inner sum, we have, using Definition 4, for $b \neq 0$

$$\begin{aligned} \sum_{u \in \mathbb{F}_q} \chi^{-1}(N(z+u)) \Psi_{-b}(u) &= \sum_{u \in \mathbb{F}_q} \chi \left(\frac{y \frac{1}{y}}{N(z+u)} \right) \Psi_{-b}(u) = \chi^{-1}(y) k(z|\chi, \Psi_{-b}) \\ &= \frac{g^{-b}}{\Gamma(\chi, \Psi_{-b})} \Psi_{-b}(-x) K_{\Psi_{-b}}(\chi \lambda | -\delta y^2, -\frac{1}{4}) \end{aligned}$$

Here we make use of Proposition 3.

It follows from formula (2.3) that

$$\begin{aligned} \frac{1}{p-1} G_{\chi \circ N^{-1}}(z) &= 1 + \frac{p}{q \chi(y)} k(z|\chi, \Psi_0) \\ &+ \frac{p}{q} \sum_{\substack{0 \neq b \in \mathbb{F}_q \\ \text{Tr}(b)=0}} \frac{\Gamma(\lambda, \Psi_1)}{\Gamma(\chi, \Psi_b)} \lambda(b^{-1}) K_{\Psi_b}(\chi \lambda | -\delta y^2, -\frac{1}{4}) \psi_b(-x). \end{aligned}$$

Apply formula (1.12) to see that

$$\begin{aligned} \frac{1}{p-1} G_{\chi \circ N^{-1}}(z) &= 1 + \frac{p}{q \chi(y)} k(z|\chi, \Psi_0) + \\ &+ \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1)}{\Gamma(\chi, \Psi_1)} \sum_{\substack{0 \neq b \in \mathbb{F}_q \\ \text{Tr}(b)=0}} (\chi \lambda)(b) K_{\Psi_b}(\chi \lambda | -\delta y^2, -\frac{1}{4}) \psi_b(-x). \end{aligned}$$

But, what is $k(z|\chi, \Psi_0)$? To answer this, note that

$$k(z|\chi, \Psi_0) = \sum_{u \in \mathbb{F}_q} \chi \left(\text{Im} \left(\frac{-1}{z+u} \right) \right) = \frac{1}{\chi(y)} \sum_{v \in \mathbb{F}_q} \chi^{-1}(v^2 - \delta).$$

Plug in formula (1.12)

$$\chi^{-1}(v^2 - \delta) = \frac{\Gamma(\chi, \Psi_{v^2 - \delta})}{\Gamma(\chi, \Psi_1)}$$

to obtain

$$\begin{aligned} k(z|\chi, \Psi_0) &= \frac{1}{\chi(y)} \sum_{v \in \mathbb{F}_q} \frac{\Gamma(\chi, \Psi_{v^2-\delta})}{\Gamma(\chi, \Psi_1)} = \frac{1}{\chi(y)\Gamma(\chi, \Psi_1)} \sum_{v \in \mathbb{F}_q} \sum_{w \in \mathbb{F}_q^*} \chi(w) e^{\frac{2\pi i \text{Tr}(w(v^2-\delta))}{p}} \\ &= \frac{1}{\chi(y)\Gamma(\chi, \Psi_1)} \sum_{w \in \mathbb{F}_q^*} \chi(w) e^{\frac{2\pi i \text{Tr}(-\delta w)}{p}} \sum_{v \in \mathbb{F}_q} e^{\frac{2\pi i \text{Tr}(wv^2)}{p}}. \end{aligned}$$

When $w \neq 0$, the inner sum is the Gauss sum $g_w = \lambda(w^{-1})\Gamma(\lambda, \Psi_1)$. Therefore,

$$k(z|\chi, \Psi_0) = \frac{\Gamma(\lambda, \Psi_1)}{\chi(y)\Gamma(\chi, \Psi_1)} \sum_{w \in \mathbb{F}_q^*} (\lambda\chi)(w) e^{\frac{2\pi i \text{Tr}(-\delta w)}{p}} = \frac{\Gamma(\lambda, \Psi_1)\Gamma(\lambda\chi, \Psi_1)}{\chi(-\delta y)\lambda(-\delta)\Gamma(\chi, \Psi_1)}.$$

The theorem follows. \square

Furthermore, if $q \neq p^2$, then one can show that $G_{\chi \circ N^{-1}}(z)$ is not the zero function by showing that the constant term of its Fourier expansion is non-zero. When $q = p^2$, our later computations show that certain of these Eisenstein series must be non-zero also. But we leave the general $q = p^2$ question open for the moment.

Corollary 1. *If $q \neq p^2$ and $a \in \{0, \frac{p-1}{2}, \frac{2(p-1)}{2}, \frac{3(p-1)}{2}, \frac{4(p-1)}{2}, \dots, \frac{m(p-1)}{2}\}$ where $\frac{m}{2} = \frac{q-1}{p-1} - 1$, then $G_{\chi_a \circ N^{-1}}(z)$ is not the zero function. If $q = p^2$, $G_{\lambda \circ N^{-1}}(z)$ is not the zero function.*

Proof. If $a = 0$, then the result is clear. If $a \neq 0$, we show that the constant term of the Fourier expansion of $G_{\chi_a \circ N^{-1}}(z)$ is non-zero. We have two cases to consider.

If $\lambda\chi_a$ is the identity character, then the constant term of $\frac{1}{p-1}G_{\chi_a \circ N^{-1}}(z)$ is $1 + \frac{p}{q}\Gamma(1, \Psi_1) = 1 - \frac{p}{q} \neq 0$.

Suppose that $\lambda\chi_a$ is not the identity character. We take the absolute value of the second term in the constant term and note that it is not 1 using formula (1.13). Therefore, the constant term of $G_{\chi_a \circ N^{-1}}(z)$ is nonzero. \square

Next we consider the Fourier expansion of some Eisenstein series which are analogous to the Maass Eisenstein series (1.1). First we define the finite analog of the power function y^s on the Poincaré upper half plane.

Let $\chi : \mathbb{F}_q^* \rightarrow \mathbb{C}$ be a multiplicative character. Define the “**power function**” $p_\chi : H_q \rightarrow \mathbb{C}$, to be

$$(2.4) \quad p_\chi(z) = \chi(\text{Im}(z)).$$

Lemma 2. *(The power function is an eigenfunction of all adjacency operators) If A_a denotes the adjacency operator defined by formula (1.6), then*

$$A_a p_\chi = \mu(a, \chi) p_\chi, \quad \text{where} \quad \mu(a, \chi) = \sum_{d(w, \sqrt{\delta})=a} \chi(\text{Im}(w)).$$

Proof. See Terras [17], page 324. \square

Now we want to relate the Eisenstein series of E-type from formula (1.7) - an Eisenstein series analogous to the Maass Eisenstein series of formula (1.1) - and the G-type Eisenstein series from definition 2. If Γ is a subgroup of $GL(2, \mathbb{F}_q)$, χ is a multiplicative character on \mathbb{F}_q^* and $p_\chi(z)$ is the power function from formula (2.4), define the finite upper half plane E-type Eisenstein series $E_{\chi, \Gamma}(z)$ by formula (1.7). Then we see that

$$(2.5) \quad E_{\chi, \Gamma}(z) = \sum_{\gamma \in \Gamma} p_\chi(\gamma z) = \sum_{\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma} \chi \left(\frac{\det(\gamma) \text{Im}(z)}{N(cz + d)} \right).$$

Note that $E_{\chi, \Gamma} \in M_{\chi_1}(\Gamma)$, where χ_1 denotes the trivial character.

Lemma 3. *Given $c, d \in \mathbb{F}_p$ not both zero and $u \in \mathbb{F}_p^*$, there exist p pairs $(a, b) \in \mathbb{F}_p^2$ such that $\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = u$.*

Proof. We need to solve the equation $xd - yc = u \neq 0$ for $x, y \in \mathbb{F}_p$. If $c \neq 0$, then we can pick any $x \in \mathbb{F}_p$ and solve for $y = c^{-1}(xd - u)$. This gives us p different choices for x . If $c = 0$, then y can be any element of \mathbb{F}_p and $x = ud^{-1}$. Again, we have p choices for y . \square

We see from the preceding results that for our two favorite subgroups of $GL(2, \mathbb{F}_q)$ we have the following **relations between the E and G-type Eisenstein series**:

$$(2.6) \quad \begin{aligned} E_{\chi, SL(2, \mathbb{F}_p)}(z) &= p\chi(\text{Im}(z))G_{\chi \circ N^{-1}}(z), \\ E_{\chi, GL(2, \mathbb{F}_p)}(z) &= p(p-1)\chi(\text{Im}(z))G_{\chi \circ N^{-1}}(z). \end{aligned}$$

Here the G-type Eisenstein series G_χ is defined by formula (2.1). In both cases of formula (2.6), when $\chi^2|_{\mathbb{F}_p^*} \neq 1$, the Eisenstein series is 0.

It follows from Lemma 2, that if $E_{\chi, \Gamma} \neq 0$ on H_q , then $E_{\chi, \Gamma}(z)$ is an eigenfunction of the adjacency operator A_a of formula (1.6) with eigenvalue

$$\mu(a, \chi) = \sum_{\substack{w \in H_q \\ d(w, \sqrt{\delta})=a}} \chi(\text{Im}(w)).$$

Thus the Eisenstein series, $E_{\chi, \Gamma}$, are the analogues of the classical Maass Eisenstein series E_s from formula (1.1) at least when they are not identically zero. The case of $\Gamma = K$, the subgroup of $GL(2, \mathbb{F}_q)$ fixing $\sqrt{\delta}$ was studied in Angel et al [2], where they were called spherical functions. The general case was introduced in [13] in an attempt to find elements of $L^2(\Gamma \backslash H_q)$ that simultaneously diagonalize the matrices $A_a^\Gamma = A_a|_{L^2(\Gamma \backslash H_q)}$.

We can now easily find the Fourier expansions of $E_{\chi, SL(2, \mathbb{F}_p)}$ and $E_{\chi, GL(2, \mathbb{F}_p)}$ using that of $G_{\chi \circ N^{-1}}(z)$ in Theorem 1.

Let $\chi = \chi_a$ be the multiplicative character of \mathbb{F}_q^* given by (1.8) and $z \in H_q$. If $a = 0$, then $E_{\chi, SL(2, \mathbb{F}_p)}(z) = p(p^2 - 1)$. If $a \in \{\frac{p-1}{2}, \frac{2(p-1)}{2}, \frac{3(p-1)}{2}, \frac{4(p-1)}{2}, \dots, \frac{m(p-1)}{2}\}$, where $\frac{m}{2} = \frac{q-1}{p-1} - 1$, then we have the Fourier expansion

$$(2.7) \quad \begin{aligned} \frac{1}{p(p-1)} E_{\chi, SL(2, \mathbb{F}_p)}(z) &= \chi(y) + \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1) \Gamma(\lambda\chi, \Psi_1)}{\Gamma(\chi, \Psi_1)} \chi^{-1}(-\delta y) \lambda(-\delta) \\ &\quad + \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1)}{\Gamma(\chi, \Psi_1)} \chi(y) \sum_{\substack{0 \neq b \in \mathbb{F}_q \\ \text{Tr}(b)=0}} (\chi\lambda)(b) K_{\Psi_b} \left(\lambda\chi \left| -\delta y^2, -\frac{1}{4} \right. \right) \psi_b(-x). \end{aligned}$$

If $a \notin \{0, \frac{p-1}{2}, \frac{2(p-1)}{2}, \frac{3(p-1)}{2}, \frac{4(p-1)}{2}, \dots, \frac{m(p-1)}{2}\}$, then $E_{\chi, SL(2, \mathbb{F}_p)}(z) = 0$.

If $q \neq p^2$ and $a \in \{0, \frac{p-1}{2}, \frac{2(p-1)}{2}, \frac{3(p-1)}{2}, \frac{4(p-1)}{2}, \dots, \frac{m(p-1)}{2}\}$ where $\frac{m}{2} = \frac{q-1}{p-1} - 1$, then $E_{\chi, SL(2, \mathbb{F}_p)}(z)$ is not the zero function.

Similar results hold for $E_{\chi, GL(2, \mathbb{F}_p)}(z)$. Let $\chi = \chi_a$ be the multiplicative character of \mathbb{F}_q^* given by (1.8) and $z \in H_q$. If $a = 0$, then $E_{\chi, GL(2, \mathbb{F}_p)}(z) = p(p-1)(p^2 - 1)$. If $a \in \{p-1, 2(p-1), \dots, m(p-1)\}$ where $m = \frac{q-1}{p-1} - 1$, then

$$(2.8) \quad \begin{aligned} \frac{1}{p(p-1)^2} E_{\chi, GL(2, \mathbb{F}_p)}(z) &= \chi(y) + \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1) \Gamma(\lambda\chi, \Psi_1)}{\Gamma(\chi, \Psi_1)} \chi^{-1}(-\delta y) \lambda(-\delta) \\ &\quad + \frac{p}{q} \frac{\Gamma(\lambda, \Psi_1)}{\Gamma(\chi, \Psi_1)} \chi(y) \sum_{\substack{0 \neq b \in \mathbb{F}_q \\ \text{Tr}(b)=0}} (\chi\lambda)(b) K_{\Psi_b} \left(\lambda\chi \left| -\delta y^2, -\frac{1}{4} \right. \right) \psi_b(-x). \end{aligned}$$

If $a \notin \{0, p-1, 2(p-1), \dots, m(p-1)\}$, then $E_{\chi, GL(2, \mathbb{F}_p)}(z) = 0$.

If $q \neq p^2$ and $a \in \{0, p-1, 2(p-1), \dots, m(p-1)\}$ where $m = \frac{q-1}{p-1} - 1$, then $E_{\chi, GL(2, \mathbb{F}_p)}(z)$ is not the zero function.

In order to compute some values of these Eisenstein series we need to think about fundamental domains. Here we will only consider $q = p^2$ for $p = 3, 5$. The fundamental domains in Examples 1 and 2 below and the Eisenstein series in Tables 1 and 2 below were calculated using Mathematica.

Example 1. $SL(2, \mathbb{F}_3) \backslash H_9$.

Write x for a root of $x^2 + 1 = 0$ and $\mathbb{F}_9 = \{a + bx \mid a, b \in \mathbb{F}\}$. Then let $\delta = 1 + x$, and suppose y is a root of the equation $y^2 = \delta$. Then, $H_9 = \{a + by \mid a, b \in \mathbb{F}_9, b \neq 0\}$. The elements of \mathbb{F}_9 are given the ordering $\{1 + 2x, 2 + 2x, 2x, 2, 0, 1, x, 1 + x, 2 + x\}$. Each $SL(2, \mathbb{F}_3)$ -orbit in figure 2 has a different shape and level of shading. The elements of the fundamental domain have been circled (or boxed). Reading from left to right increases $\text{Re}(z)$, and reading from bottom to top increases $\text{Im}(z)$. More explicitly a fundamental domain for $SL(2, \mathbb{F}_3) \backslash H_9$ is the set:

$$\{1 + 2x + (2 + x)y, 2 + (2 + x)y, 1 + 2x + (1 + x)y, 2 + (1 + x)y, x + (1 + x)y, 1 + 2x + xy\}.$$

Example 2. $SL(2, \mathbb{F}_5) \backslash H_{25}$.




FIGURE 2. The fundamental domain for $SL(2, \mathbb{F}_3 \setminus H_9)$ is given by the 6 points which are either circled or boxed.

Let x be a root of $x^2 + 2 = 0$ and $\mathbb{F}_{25} = \{a + bx \mid a, b \in \mathbb{F}_5, \}$. Suppose $\delta = 1 + x$, and y satisfies $y^2 = \delta$. Then we have $H_{25} = \{a + by \mid a, b \in \mathbb{F}_{25}, b \neq 0\}$. It can be shown that a fundamental domain for $SL(2, \mathbb{F}_5) \setminus H_{25}$ is given by the 10 points

$$\{1 + 3x + (4 + 2x)y, 1 + 4x + (4 + 2x)y, 3 + (4 + 2x)y, 2x + (4 + 2x)y, 3 + (3 + 2x)y, \\ 1 + 4x + (2 + 2x)y, x + (2 + 2x)y, 1 + 4x + (1 + 2x)y, 3 + (1 + 2x)y, 1 + 4x + (4 + x)y\}.$$

We shall write A,B,C,D,E,F,G,H,I,J for these 10 points to save space in our table 2.

More general results on fundamental domains can be found in [13].

Finally we compute values of $E_{\chi, SL(2, \mathbb{F}_p)}(z)$ for the preceding examples. One can compute $E_{\chi, GL(2, \mathbb{F}_p)}(z)$ for the same values of p and q using formula (2.6).

Table 1 below gives the values of $E_{\chi, SL(2, \mathbb{F}_5)}$ on $SL(2, \mathbb{F}_3) \setminus H_9$ for $\chi = \chi_a, 0 \leq a \leq 7$. Table 2 below gives the values of $E_{\chi, SL(2, \mathbb{F}_5)}$ on $SL(2, \mathbb{F}_5) \setminus H_{25}$ for $\chi = \chi_a, 0 \leq a \leq 23$. It follows from our computations that at least for these special cases the Eisenstein series on H_{p^2} for $SL(2, \mathbb{F}_p)$ do not vanish.

REFERENCES

- [1] J. Angel, Finite upper half planes over finite fields, *Finite Fields Appl.*, 2 (1996), 62-86.
- [2] J. Angel, N. Celniker, S. Poulos, A. Terras, C. Trimble, and E. Velasquez, Special functions on finite upper half planes, *Contemp. Math.*, 128 (1992), 1-26.
- [3] J. Angel, and R. Evans, Diameters of finite upper half plane graphs, *J. Graph Theory*, 23 (1996), 129-137.
- [4] E. Bannai and T. Ito, *Association Schemes, Algebraic Combinatorics I*, Benjamin/Cummings, Menlo Park, CA, 1984.
- [5] P. T. Bateman and E. Grosswald, On Epstein's zeta function, *Acta Arithmetica*, 9 (1964), 365-373.
- [6] N. Celniker, S. Poulos, A. Terras, C. Trimble, and E. Velasquez, Is there life on finite upper half planes?, *Contemp. Math.*, Vol. 143 (1993), 65-88.
- [7] Chai, C.-L. and Li, W.-C. W., Character sums, automorphic forms, equidistribution, and Ramanujan graphs, II, Eigenvalues of Terras graphs, *Forum Math.*, 16 (2004), 631-661.
- [8] R. Evans, Spherical functions for finite upper half planes with characteristic 2, *Finite Fields Appl.* 1 (1995), 37-394.
- [9] D. Hejhal, *The Selberg Trace Formula for $PSL(2, \mathbb{R})$, I*, *Lecture Notes in Math.*, Vol. 548, Springer-Verlag, N.Y., 1967.
- [10] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, N.Y., 1982.
- [11] A. Krieg, *Hecke Algebras*, *Memoirs of Amer. Math. Soc.*, 435, A.M.S., Providence, 1990.
- [12] P. Sarnak, Kloosterman, quadratic forms and modular forms, *Kloosterman Centennial Celebration, Nieuw. Arch. Wiskd.*, 5/1, nr.4 (Dec., 2000), 385-389.
- [13] A. Shaheen, *Finite Planes and Finite Upper Half Planes: Their Geometry, a Trace Formula, Modular Forms and Eisenstein Series*, Ph.D. Thesis, U.C.S.D., 2005.
- [14] J. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, Springer-Verlag, N.Y., 1994.
- [15] J. Soto-Andrade, and J. Vargas, Twisted spherical functions on the finite Poincaré upper half-plane, *J. Algebra*, 248 (2002), 724-46.
- [16] A. Terras, *Harmonic Analysis on Symmetric Spaces and Applications, Vol. I*, Springer-Verlag, New York, 1985.
- [17] A. Terras, *Fourier Analysis on Finite Groups and Applications*, Cambridge University Press, Cambridge, 1999.
- [18] A. Terras, Bessel series expansions of the Epstein zeta function and the functional equation, *Trans. Amer. Math. Soc.*, 183 (1973), 477-486.
- [19] P. D. Tiu and D. I. Wallace, Norm quadratic residue codes, *IEEE Trans. Information Theory*, 40 (1994), 946-949.

Table 1. Values of $E_{\chi, SL(2, \mathbb{F}_3)}$ on $SL(2, \mathbb{F}_3) \setminus H_9$ for $\chi = \chi_a, 0 \leq a \leq 7$.

z	$E_{\chi_0, SL(2, \mathbb{F}_3)}$	$E_{\chi_1, SL(2, \mathbb{F}_3)}$
$1 + 2x + (2 + x)y$	24	$-6i + (3 - 9i)\sqrt{2}$
$2 + (2 + x)y$	24	$(-6 + 12i) + (3 - 3i)\sqrt{2}$
$1 + 2x + (1 + x)y$	24	0
$2 + (1 + x)y$	24	$6i - (3 - 9i)\sqrt{2}$
$x + (1 + x)y$	24	0
$1 + 2x + xy$	24	$(6 - 12i) - (3 - 3i)\sqrt{2}$
z	$E_{\chi_2, SL(2, \mathbb{F}_3)}$	$E_{\chi_3, SL(2, \mathbb{F}_3)}$
$1 + 2x + (2 + x)y$	$-6 - 6i$	$6i - (3 + 9i)\sqrt{2}$
$2 + (2 + x)y$	$-6 - 6i$	$-6 - 12i - (3 + 3i)\sqrt{2}$
$1 + 2x + (1 + x)y$	$12 + 12i$	0
$2 + (1 + x)y$	$-6 - 6i$	$-6i + (3 + 9i)\sqrt{2}$
$x + (1 + x)y$	$12 + 12i$	0
$1 + 2x + xy$	$-6 - 6i$	$6 + 12i + (3 + 3i)\sqrt{2}$
z	$E_{\chi_4, SL(2, \mathbb{F}_3)}$	$E_{\chi_5, SL(2, \mathbb{F}_3)}$
$1 + 2x + (2 + x)y$	-12	$-6i - (3 - 9i)\sqrt{2}$
$2 + (2 + x)y$	12	$-6 + 12i - (3 - 3i)\sqrt{2}$
$1 + 2x + (1 + x)y$	0	0
$2 + (1 + x)y$	-12	$6i + (3 - 9i)\sqrt{2}$
$x + (1 + x)y$	0	0
$1 + 2x + xy$	12	$6 - 12i + (3 - 3i)\sqrt{2}$
z	$E_{\chi_6, SL(2, \mathbb{F}_3)}$	$E_{\chi_7, SL(2, \mathbb{F}_3)}$
$1 + 2x + (2 + x)y$	$-6 + 6i$	$6i + (3 + 9i)\sqrt{2}$
$2 + (2 + x)y$	$-6 + 6i$	$-6 - 12i + (3 + 3i)\sqrt{2}$
$1 + 2x + (1 + x)y$	$12 - 12i$	0
$2 + (1 + x)y$	$-6 + 6i$	$-6i - (3 + 9i)\sqrt{2}$
$x + (1 + x)y$	$12 - 12i$	0
$1 + 2x + xy$	$-6 + 6i$	$6 + 12i - (3 + 3i)\sqrt{2}$

Table 2. Values of $E_{\chi, \Gamma}$ on $\Gamma \backslash H_{25}$ for $\chi = \chi_a$, $0 \leq a \leq 23$, $\Gamma = SL(2, \mathbb{F}_5)$.

z	$E_{\chi_0, SL(2, \mathbb{F}_5)}(z)$	$E_{\chi_2, SL(2, \mathbb{F}_5)}(z)$	$E_{\chi_4, SL(2, \mathbb{F}_5)}(z)$
A	120	0	$20 + 20i\sqrt{3}$
B	120	$-30 - 10i - (10 - 30i)\sqrt{3}$	$20 + 20i\sqrt{3}$
C	120	$30 - 40i + (20 - 10i)\sqrt{3}$	$-30 - 30i\sqrt{3}$
D	120	0	$20 + 20i\sqrt{3}$
E	120	$-50i + (10 + 20i)\sqrt{3}$	$-30 - 30i\sqrt{3}$
F	120	0	$20 + 20i\sqrt{3}$
G	120	0	$20 + 20i\sqrt{3}$
H	120	$50i - (10 + 20i)\sqrt{3}$	$-30 - 30i\sqrt{3}$
I	120	$30 + 10i + (10 - 30i)\sqrt{3}$	$20 + 20i\sqrt{3}$
J	120	$-30 + 40i - (20 - 10i)\sqrt{3}$	$-30 - 30i\sqrt{3}$
z	$E_{\chi_6, SL(2, \mathbb{F}_5)}(z)$	$E_{\chi_8, SL(2, \mathbb{F}_5)}(z)$	$E_{\chi_{10}, SL(2, \mathbb{F}_5)}(z)$
A	0	$-60 - 20i\sqrt{3}$	0
B	$-60 - 20i$	0	$-30 - 10i + (10 - 30i)\sqrt{3}$
C	$-60 - 20i$	$-30 - 10i\sqrt{3}$	$30 - 40i - (20 - 10i)\sqrt{3}$
D	0	$-60 - 20i\sqrt{3}$	0
E	$60 + 20i$	$30 + 10i\sqrt{3}$	$-50i - (10 + 20i)\sqrt{3}$
F	0	$60 + 20i\sqrt{3}$	0
G	0	$60 + 20i\sqrt{3}$	0
H	$-60 - 20i$	$30 + 10i\sqrt{3}$	$50i + (10 + 20i)\sqrt{3}$
I	$60 + 20i$	0	$30 + 10i - (10 - 30i)\sqrt{3}$
J	$60 + 20i$	$-30 - 10i\sqrt{3}$	$-30 + 40i + (20 - 10i)\sqrt{3}$
z	$E_{\chi_{12}, SL(2, \mathbb{F}_5)}(z)$	$E_{\chi_{14}, SL(2, \mathbb{F}_5)}(z)$	$E_{\chi_{16}, SL(2, \mathbb{F}_5)}(z)$
A	-40	0	$-60 + 20i\sqrt{3}$
B	80	$-30 + 10i + (10 + 30i)\sqrt{3}$	0
C	0	$30 + 40i - (20 + 10i)\sqrt{3}$	$-30 + 10i\sqrt{3}$
D	-40	0	$-60 + 20i\sqrt{3}$
E	0	$50i - (10 - 20i)\sqrt{3}$	$30 - 10i\sqrt{3}$
F	-40	0	$60 - 20i\sqrt{3}$
G	-40	0	$60 - 20i\sqrt{3}$
H	0	$-50i + (10 - 20i)\sqrt{3}$	$30 - 10i\sqrt{3}$
I	80	$30 - 10i - (10 + 30i)\sqrt{3}$	0
J	0	$-30 - 40i + (20 + 10i)\sqrt{3}$	$-30 + 10i\sqrt{3}$
z	$E_{\chi_{18}, SL(2, \mathbb{F}_5)}(z)$	$E_{\chi_{20}, SL(2, \mathbb{F}_5)}(z)$	$E_{\chi_{22}, SL(2, \mathbb{F}_5)}(z)$
A	0	$20 - 20i\sqrt{3}$	0
B	$-60 + 20i$	$20 - 20i\sqrt{3}$	$-30 + 10i - (10 + 30i)\sqrt{3}$
C	$-60 + 20i$	$-30 + 30i\sqrt{3}$	$30 + 40i + (20 + 10i)\sqrt{3}$
D	0	$20 - 20i\sqrt{3}$	0
E	$60 - 20i$	$-30 + 30i\sqrt{3}$	$50i + (10 - 20i)\sqrt{3}$
F	0	$20 - 20i\sqrt{3}$	0
G	0	$20 - 20i\sqrt{3}$	0
H	$-60 + 20i$	$-30 + 30i\sqrt{3}$	$-50i - (10 - 20i)\sqrt{3}$
I	$60 - 20i$	$20 - 20i\sqrt{3}$	$30 - 10i - (10 + 30i)\sqrt{3}$
J	$60 - 20i$	$-30 + 30i\sqrt{3}$	$-30 - 40i - (20 + 10i)\sqrt{3}$

ANTHONY SHAHEEN, MATH. DEPT., CALIFORNIA STATE UNIVERSITY, LOS ANGELES

AUDREY TERRAS, MATH. DEPT., UNIVERSITY OF CALIFORNIA AT SAN DIEGO, LA JOLLA, CA 92093-0112

E-mail address: aterras@ucsd.edu

URL: <http://www.math.ucsd.edu/~aterras>