

Conversely, suppose X is reduced and irreducible.

Let $V \subset X$ be open, $V \neq \emptyset$.

Suppose $f, g \in \mathcal{O}_X(V)$ with $fg = 0$.

Recall: For a locally ringed space (X, \mathcal{O}_X) ,

$\forall f \in \mathcal{O}_X(X)$, the set $X_f := \{x \in X \mid f(x) \notin \mathfrak{m}_x \subset \mathcal{O}_{X,x}\}$
is open.

endow V with the locally ringed space structure induced

from X : $V \subset U$ $\mathcal{O}_V(V) := \mathcal{O}_X(V)$.

Then $Z(f) := \{x \in V \mid f(x) \in \mathfrak{m}_x \subset \mathcal{O}_{V,x}\}$
is closed in $V = V \setminus V_f$

Similarly $Z(g)$ is closed in V .

$fg = 0 \Rightarrow Z(fg) = V \Rightarrow Z(f) \cup Z(g) = V$

Recall from homework that V is irreducible.

$$\Rightarrow Z(f) = V \quad \text{or} \quad Z(g) = V$$

Suppose $Z(f) = V$. Then for any affine open $\text{Spec} A \subset V$,

$$\forall p \in \text{Spec} A \quad \mathcal{O}_{V,p} = A_p$$

$$\text{and } \mathfrak{m}_p = \mathfrak{p} A_p$$

$$Z(f) = V \Rightarrow Z(f) \supset \text{Spec} A \Rightarrow \nexists (p) \in \mathfrak{p} A_p$$

$$\forall p \in \text{Spec} A$$

$f(p)$ = image of $f|_{\text{Spec} A}$ in the localization A_p

$$= \frac{f|_{\text{Spec} A}}{1}$$

$$\Rightarrow \forall p \in \text{Spec} A$$

$$\frac{f|_{\text{Spec} A}}{1} \in \mathfrak{p} \subset A$$

$$\begin{array}{c} \mathcal{O}_X(U) \\ \downarrow \\ A \end{array}$$

$\Rightarrow \bigcap_{p \in \text{Spec } A} p = \text{nilradical}$
 $p \in \text{Spec } A := \text{set of all nilpotent elements}$

$\Rightarrow \bigcap_{p \in \text{Spec } A} p \text{ is nilpotent } \in A$

we assumed X is reduced $\Rightarrow \bigcap_{p \in \text{Spec } A} p = 0$

This holds for all open affine in $U \Rightarrow f = 0$

because U is covered by open affine subsets.

$\Rightarrow \mathcal{O}_X(U)$ is an integral domain.

□

Proposition: A scheme is locally noetherian if and only if, for any open affine $\text{Spec } A \cong U \subset X$, A is noetherian. In particular, an affine scheme is locally

noetherian if and only if A is noetherian.

Proof: The "if" part is clear.

Assume X is locally noetherian, let $\text{Spec} A = U \subset X$ be open. We will show that A is a noetherian ring.

X has a cover by open affine sets. $X = \bigcup_{i \in I} \text{Spec} B_i$
Choose $i \in I$
 $\forall f \in B = B_i \quad B[f^{-1}] = B[x] / (fx - 1)$ is noetherian

by the Hilbert basis theorem.

\Rightarrow open affine sets with noetherian rings of sections form a basis of the topology of X .

$\Rightarrow U = \text{Spec} A$ has a covering by open sets with noetherian rings of sections.

$$U = \bigcup_{j \in J} V_j \quad V_j = \text{Spec } C_j \quad C_j \text{ noetherian}$$

$$\exists f \in A \text{ s.t. } U_f = \text{Spec } A[f^{-1}] \subset V_j = \text{Spec } C_j$$

$$\text{write } V_j = V, \quad C_j = C$$

$$U \supset V \quad A = G(U) \longrightarrow C = G(V)$$

$$f \longmapsto f|_V =: \bar{f}$$

$$\text{Claim: } U_f = V_{\bar{f}} \quad \left(\Rightarrow \begin{array}{l} A[f^{-1}] = C[\bar{f}^{-1}] \\ \text{ring of } U_f \quad \text{ring of } V_{\bar{f}} \end{array} \right)$$

$$U_f = \{ \mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p} \} = \{ \mathfrak{p} \in \text{Spec } A \mid f(\mathfrak{p}) \notin \mathfrak{p}A_{\mathfrak{p}} \}$$

$$= \{ \sigma \in \text{Spec } C \mid \bar{f}(\sigma) \notin \sigma C_{\sigma} \}$$

$$= \{ \sigma \in \text{Spec } C \mid \bar{f} \notin \sigma \} = V_{\bar{f}} \subset \text{Spec } C$$

U is affine, hence quasi-compact, so we can cover U with a finite number of open sets U_{f_1}, \dots, U_{f_n} whose rings are $A[f_1^{-1}] = C_{j_1}[f_1^{-1}], \dots, A[f_n^{-1}] = C_{j_n}[f_n^{-1}]$ which are noetherian.

Since $U = U_{f_1} \cup \dots \cup U_{f_n}$, the elements f_1, \dots, f_n generate the unit ideal in A .

Now we show A is noetherian.

Let $\mathcal{O}_1 \subset \mathcal{O}_2 \subset \dots \subset \mathcal{O}_n \subset \dots$

be an ascending chain of ideals. $\forall i$

$$\mathcal{O}_1[f_i^{-1}] \subset \mathcal{O}_2[f_i^{-1}] \subset \dots \subset A[f_i^{-1}]$$

is stationary because $A[f_i^{-1}]$ is noetherian.

The original chain $\sigma_1 \subset \sigma_2 \subset \dots$
 is stationary by the following lemma:

Lemma: Let A be a ring, $f_1, \dots, f_n \in A$ which
 generate the unit ideal. Then for any ideal $\sigma \subset A$,

$$\sigma = \bigcap_{i=1}^n \varphi_i^{-1}(\varphi_i(\sigma) A[f_i^{-1}])$$

where $\varphi_i: A \rightarrow A[f_i^{-1}]$ is the localization morphism.

Proof: Clearly $\sigma \subset \bigcap_{i=1}^n \varphi_i^{-1}(\varphi_i(\sigma) A[f_i^{-1}])$

For the reverse inclusion, choose $b \in \bigcap_{i=1}^n \varphi_i^{-1}(\varphi_i(\sigma) A[f_i^{-1}])$

$$\forall i \exists n_i \in \mathbb{Z}, \exists a_i \in \sigma \text{ s.t. } \varphi_i(b) = \frac{a_i}{f_i^{n_i}} = \frac{f_i^{n_i} a_i}{f_i^{n_i + n}} \quad \forall n$$

\Rightarrow can assume $\exists n, a_i \in \mathcal{O}$ s.t. $\forall i \varphi_i(b) = \frac{a_i}{f_i^n}$

$$\frac{1}{1} = \varphi_i(b) = \frac{a_i}{f_i^n}$$

$$\exists m_i \in \mathbb{Z}_+ \text{ s.t. } f_i^{m_i} (f_i^n b - a_i) = 0$$

increase m_i if necessary:

$$\exists m, n \text{ s.t. } f_i^m (f_i^n b - a_i) = 0$$

$$\Rightarrow \forall i \quad f_i^{m+n} b (= f_i^m a_i) \in \mathcal{O}$$

$$\langle f_1, \dots, f_r \rangle = A \Rightarrow \langle f_1^{m+n}, \dots, f_r^{m+n} \rangle = A$$

$$\Rightarrow \exists c_1, \dots, c_r \in A \text{ s.t. } 1 = c_1 f_1^{m+n} + \dots + c_r f_r^{m+n}$$

$$b = b \cdot 1 = c_1 f_1^{m+n} b + \dots + c_r f_r^{m+n} b \in \mathcal{O}.$$

□

First properties of morphisms of schemes:

Definition: (1) A morphism of schemes $f: X \rightarrow Y$ is locally of finite type if \exists a covering of Y by open affine subsets $V_i = \text{Spec } B_i$ s.t. $\forall i$ $f^{-1}(V_i)$ has a covering by open affine subsets $U_{ij} := \text{Spec } A_{ij}$ where $\forall i, j$

A_{ij} is a finitely generated B_i -algebra.

(a small explanation: $f: X \rightarrow Y$ $f^\# : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_X$
 $\forall i, f^{-1}(V_i) = \bigcup_j U_{ij} \rightarrow V_i$ or $\forall i, j, f|_{U_{ij}} : U_{ij} \rightarrow V_i$
 $f|_{U_{ij}} : \text{Spec } A_{ij} \rightarrow \text{Spec } B_i$
gives A_{ij} a structure of B_i -algebra.