

Consider the subset

$$B := \left\{ (x, H) \mid \begin{array}{l} x \in H, \text{ either } x \text{ is a singular} \\ \text{point of } X \cap H \text{ or } H \supset X \end{array} \right\}$$

$$\subset X \times_k \mathbb{P}^n / \mathcal{O}_{\mathbb{P}^n}(1)$$

Claim:  $B$  has a natural structure of a closed subscheme of  $X \times_k \mathbb{P}^n / \mathcal{O}_{\mathbb{P}^n}(1)$ .

Proof: If  $(x, H) \in B$ , let  $f \in \Gamma(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$  be an equation for  $H$ , i.e.,  $Z(f) = H$  and let  $\{P_1, \dots, P_r\}$  be a set of generators for  $I_X$ . Then,  $I_{X \cap H}$  is generated by  $(f, P_1, \dots, P_r)$ .

We have  $(x, H) \in B \iff$  the Zariski tangent space  $T_x X \cap H$  has dimension  $> d-1$  where  $d = \dim X$

By the description of the embedded tangent space to

$X \cap H$ ,  $T_x X \cap H$  has  $\dim > d-1$

$\iff$   $\left( \frac{\partial f}{\partial X_i}(x), \frac{\partial P_j}{\partial X_i}(x) \right)_{\substack{1 \leq j \leq r \\ 0 \leq i \leq n}}$  has rank  $< n-d+1$

So the minors of size  $(n-d+1) \times (n-d+1)$  of the above matrix give polynomial equations for  $B$  as a closed subscheme of  $X \times (\mathbb{C}^{pp^2}(1))$  in the coordinates

$(X_0, \dots, X_n)$  on  $X \subset \mathbb{P}^n$  and coordinates  
 $(\lambda_0, \dots, \lambda_n)$  on  $|\mathcal{O}_{\mathbb{P}^n}(1)|$  where  $f = \sum_{i=0}^n \lambda_i X_i$   
 $= (\mathbb{P}^n)^* \cong \mathbb{P}^n$   
 (abstract) □

We continue the proof of Bertini's theorem:

Choose  $(x, H) \in X \times |\mathcal{O}_{\mathbb{P}^n}(1)|$ ,  $H = Z(f)$

Let  $i$  be such that  $x \in U_i = D_+(X_i)$

On  $U_i$ , the ideal  $I_{H \cap U_i}$  is generated by  $\frac{f}{X_i}$

consider the linear map

$$\begin{array}{ccc}
 \varepsilon_x: \Gamma(\mathcal{O}_{\mathbb{P}^n}(1)) & \longrightarrow & \overbrace{\mathcal{O}_{X, x} / \mathfrak{m}_x^2}^{\text{dim. } d+1/\mathbb{R}} \\
 f & \longmapsto & \left( \frac{f}{X_i} \right)_x \text{ modulo } \mathfrak{m}_x^2 \supset \mathfrak{m}_x / \mathfrak{m}_x^2
 \end{array}$$

We have  $H \supset X$  and  $x$  is a singular point of  $X \cap H$

$$\Leftrightarrow f \in \ker \varepsilon_x.$$

Because  $(f/x_i)_x \in \mathcal{O}_{X,x}$  generates the ideal of  $H \cap X$  in  $\mathcal{O}_{X,x}$

$$H \supset X \Leftrightarrow (f/x_i)_x = 0 \text{ in } \mathcal{O}_{X,x} = \text{localization of ring of } X \cap U_i$$

(localization map is injective because  $X$  is integral.)

( $H \not\subset X$  and  $x$  is a non-singular point of  $X \cap H$ )

$$\Leftrightarrow T_x(X \cap H) \text{ has dim. } d-1.$$

$$\Leftrightarrow m_{X \cap H, x} / m_{X \cap H, x}^2 \text{ has dim. } d-1$$

Note that:  $\mathcal{O}_{X \cap H, x} = \mathcal{O}_{X, x} / (f/x_i)_x$

and  $\mathfrak{m}_{X \cap H, x} = \mathfrak{m}_{X, x} / (f/x_i)_x$   
( $x \in X \cap H \Leftrightarrow (f/x_i)_x \in \mathfrak{m}_{X, x}$ )

and  $\mathfrak{m}_{X \cap H, x} / \mathfrak{m}_{X \cap H, x}^2 = \mathfrak{m}_{X, x} / \mathfrak{m}_{X, x}^2 / \left( (f/x_i)_x \bmod \mathfrak{m}_{X, x}^2 \right)$

So  $\dim \mathfrak{m}_{X \cap H, x} / \mathfrak{m}_{X \cap H, x}^2 = d-1$

$\Leftrightarrow (f/x_i)_x \neq 0 \bmod \mathfrak{m}_{X, x}^2$

So, if  $x \in X \cap H$  is a singular point, then

$$\left(\frac{f}{X_i}\right)_x = 0 \text{ modulo } \mathfrak{m}_{X,x}^2.$$

This proves that  $(x, H) \in B \iff f \in \ker \varepsilon_x$ .

Note: that if we change the index  $i$  sit.

$x \in U_i$ ,  $\varepsilon_x$  changes by multiplication by  $\left(\frac{X_i}{X_j}\right)(x)$  which is a nonzero scalar. So  $\ker \varepsilon_x$  is independent of  $i$ .

Note:  $\varepsilon_x$  is surjective because  $\mathcal{M}_x$  is generated by images (in  $\mathcal{O}_{X,x}$ ) of some degree  $\leq 1$  polynomials, i.e., elements of  $T(\mathcal{O}_{\mathbb{P}^n}(1))$  (we are over an alg. closed field).


$$\mathcal{M}_{\mathbb{P}^n, x} \rightarrow \mathfrak{m}_{X,x}$$

$$\text{If } x = (a_0, \dots, a_n) \quad M_{\mathbb{P}^n, x} = \begin{pmatrix} a_i X_j - a_j X_i \\ 0 \leq i, j \leq n \end{pmatrix}$$

So  $\text{Ker } \varepsilon_x$  has dim.  $n+1 - (d+1) = n - d$  over  $k$ .

Now:

We show  $q_2(B)$  is a proper closed subset of  $|\mathcal{O}_{\mathbb{P}^n}(1)|$



By Ex. II.3.22 about the fibers of a morphism,

the underlying set of  $q_1^{-1}(x)$  is the set of preimages of  $x$  by  $q_1$ .

$$q_1^{-1}(x) = \{ (x, H) \mid (x, H) \in B \}$$

$$= \{ (x, H) \mid H \in \text{Ker } \varepsilon_x \} = \{x\} \times \mathbb{P} \text{Ker } \varepsilon_x$$

$$\begin{aligned} \text{So } \dim q_1^{-1}(x) &= \dim(\{x\} \times \mathbb{P} \text{Ker } \varepsilon_x) = \dim(\mathbb{P} \text{Ker } \varepsilon_x) \\ &= n - d - 1 \end{aligned}$$

$$\begin{aligned} \text{Ex. II.3.22 : } \dim B &= \dim q_1(B) + \dim q_1^{-1}(x) \\ &\leq \dim X + n - d - 1 = n - 1 \end{aligned}$$

Ex. II.4.4 the image of a proper scheme is proper.

$B$  is a closed subscheme of  $X \times_{\mathbb{k}} |\mathcal{O}_{\mathbb{P}^n}(1)|$ , hence it is proper /  $\mathbb{k}$ . So  $q_2(B)$  is also proper, hence it is closed in  $|\mathcal{O}_{\mathbb{P}^n}(1)|$ , and  $\dim q_2(B) \leq \dim B \leq n-1$ .

Since  $\dim \mathbb{P}^n = n$ ,  $q_2(B) \not\subset |\mathcal{O}_{\mathbb{P}^n}(1)|$ .

□