

A Note on Constructive Methods for Ramsey Numbers

F. R. K. Chung
BELL LABORATORIES
MURRAY HILL, NEW JERSEY

ABSTRACT

Let $r(k)$ denote the least integer n such that for any graph G on n vertices either G or its complement \bar{G} contains a complete graph K_k on k vertices. In this paper, we prove the following lower bound for the Ramsey number $r(k)$ by explicit construction: $r(k) \geq \exp(c(\log k)^{4/3} / [(\log \log k)^{1/3}])$ for some constant $c > 0$.

For an integer k , the Ramsey number $r(k)$ is defined to be the least integer n such that for any graph G on n vertices, either G or its complement \bar{G} contains a complete graph K_k on k vertices. The theory of Ramsey numbers has been extensively studied in the past. However, relatively few results for $r(k)$ have yet been found. With respect to exact values, we only know $r(3) = 6$ and $r(4) = 18$ (see [1, 8]). A lower bound 42 for $r(5)$ was proved (but unpublished) by S. Lin and, independently, by J. P. Burling. An upper bound 55 for $r(5)$ was given in [11]. Thus we have

$$42 \leq r(5) \leq 55.$$

For general k , the following upper bound for $r(k)$ is still the best known so far.*

$$r(k) \leq c \binom{2k-2}{k-1}$$

for a suitable constant c .

P. Erdős [3] has proved the following lower bound by probabilistic

* The widely quoted upper bound $c \frac{\log \log k}{\log k} \binom{2k-2}{k-1}$ by J. Yackel [12] seems now in question [13].

arguments:

$$r(k) \geq k2^{k/2} \left(\frac{1}{e\sqrt{2}} + o(1) \right).$$

J. Spencer [10] improved the above bound by a factor of 2 also by nonconstructive methods. Erdős [4] has asked whether one can find an explicit construction for a graph G on $2^{k/2}$ vertices such that neither G nor its complement \bar{G} contains K_k . This problem, however, falls into an interesting category of problems which have the property that for any large n the existence of a "good configuration" is assured by probabilistic methods, (in fact most of the configurations are good), but we cannot explicitly find even one "good configuration."

H. L. Abbott [1] gives a recursive construction which shows that $r(k) \geq ck^{c'}$, where $c' = \log 41 / \log 4 = 2.679 \dots$. Nagy [9] gives a construction which shows that $r(k) \geq ck^3$. P. Frankl [7] shows constructively that

$$r(k) \geq ck^m$$

for any m and some constant c .

In this note, we will give the constructive lower bound:

$$r(k) \geq \exp [c(\log k)^{4/3} / \log \log k]^{1/3}, \quad \text{for some constant } c.$$

In other words, we present an explicit construction of a graph G on n vertices such that neither G nor its complement \bar{G} contains a complete graph on $\exp [c(\log n)^{3/4} (\log \log n)^{1/4}]$ vertices.

The basic ideas of this lower bound are due to P. Frankl [7]. We will tighten up some loose ends in [7] and give a self-contained proof of the following theorem on intersecting families (except for the use of a result of P. Erdős and R. Rado).

Theorem 1. For integers x, y, z, w, p, u , with $x \geq 2z > 0$, $p \leq xy + z + w$, $p \leq xu + w$ we define

$$L(x, y, z, w) = \{xy' + z' + w : 0 \leq y' < y, 0 \leq z' < z\}.$$

Let \mathbf{F} be a family of distinct p -subsets of $X = \{1, \dots, n\}$ such that for any two sets $F_1, F_2 \in \mathbf{F}$ we have $|F_1 \cap F_2| \in L(x, y, z, w)$. Then we have

$$|\mathbf{F}| \leq n^{u+y+z} p^{2p(u+y)}.$$

The proof of Theorem 1 is based on a recursive argument using Δ -systems. First we need some definitions.

A family of sets, S_1, \dots, S_n , is said to be a Δ -system if $S_i \cap S_j = S_i \cap S_j$.

for any $i \neq j, i' \neq j'$. In this case, $D = S_i \cap S_j, i \neq j$, is called the *kernel* of this Δ -system.

Theorem (Erdős and Rado [6]). For any t sets, S_1, \dots, S_t , with $|S_i| \leq s$ for $1 \leq i \leq t$, there exists a Δ -system consisting of $r + 1$ S_i 's provided

$$t > s! r^{s+1} \left(1 - \frac{1}{2! r} - \frac{2}{3! r^2} \dots - \frac{s-1}{s! r^{s-1}} \right).$$

Let \mathbf{F} be a family of distinct p -subsets of X such that for any $F_1, F_2 \in \mathbf{F}$ we have $|F_1 \cap F_2| \in L(x, y, z, w)$. If $|\mathbf{F}| \geq p^{2p}$, then \mathbf{F} contains a Δ -system with $p + 1$ subsets. Let D_1 be a subset of X with maximal cardinality such that D_1 is the kernel of a Δ -system consisting of $p + 1$ sets in \mathbf{F} . We define $\mathbf{F}_1 = \{F \in \mathbf{F} : D_1 \subset F\}$. If $|\mathbf{F} - \mathbf{F}_1| \geq p^{2p}$, then in a similar manner we consider D_2 which is the kernel of maximal cardinality of a Δ -system consisting of $p + 1$ sets in $\mathbf{F} - \mathbf{F}_1$ and we define $\mathbf{F}_2 = \{F \in \mathbf{F} - \mathbf{F}_1 : D_2 \subset F\}$. After a finite number of steps, we have found $D_1, \dots, D_t, \mathbf{F}_1, \dots, \mathbf{F}_t$ and $\mathbf{F} - \bigcup_{i=1}^t \mathbf{F}_i = \mathbf{F}_{t+1}$ contains fewer than p^{2p} sets.

We note that $\mathbf{F}_i, i = 1, \dots, t$, contains no more than np^{2p-1} sets (otherwise there are at least $p^{2(p-|D_i|)}$ sets in \mathbf{F}_i all of which contain a common element in $X - D_i$; this would contradict the maximality of D_i). We also note that for a Δ -system F_1, \dots, F_{p+1} with kernel D_i , we know that $F_{i'} - D_i, i' = 1, \dots, p + 1$, are pairwise disjoint. Thus, for any p' -subset $X' \subset X, p' \leq p$ we have $X' \cap D_i = X' \cap F_{i'}$ for some i' . Thus $D_i \cap D_j = F_{i'} \cap F_{j'}$ and $|D_i \cap D_j| \in L(x, y, z, w)$.

We will prove Theorem 1 by induction on p . It is easy to see that Theorem 1 holds for $p = 0$. For $p > 0$, we let $\mathbf{E}_i = \{F : F \in \mathbf{F}_i \text{ and } |D_i| = p - i\}$ for $0 < i \leq p$. Let $\mathbf{E}_{i_0} = \mathbf{E}$ have the property that $|\mathbf{E}_{i_0}| \geq |\mathbf{E}_i|$ for any i . Therefore we have

$$p |\mathbf{E}| \geq |\mathbf{F}| - p^{2p}.$$

Suppose $i_0 < z$. We define $\mathbf{X}_A = \{D_i : D_i \cup A \in \mathbf{E} \text{ and } D_i \cap A = \emptyset\}$ for $A \subset X$ and $|A| = i_0$. It is easy to see that for $D_i, D_j \in \mathbf{X}_A$ we have $|D_i \cap D_j| \in L(x, y, z - i_0, w)$. Therefore by the induction assumptions, we have

$$|\mathbf{X}_A| \leq n^{u+y+z-i_0} (p - i_0)^{2(p-i_0)(u+y)}$$

and

$$|\mathbf{F}| \leq p^{2p} + p \sum_A |\mathbf{X}_A| \leq p^{2p} + \binom{n}{i_0} n^{u+y+z-i_0} p^{(2p-1)(u+y)} \leq n^{u+y+z} p^{2p(u+y)}.$$

So, we may assume $i_0 \geq z$. We consider $\mathbf{D} = \{D_i : |D_i| = p - i_0, 1 \leq i \leq t\}$.

We have

$$|\mathbf{E}| \leq \sum_{|D_i|=p-i_0} |\mathbf{F}_i| \leq np^{2p-1} |\mathbf{D}|.$$

It suffices to show that

for any $D_i, D_j \in \mathbf{D}$ we have $|D_i \cap D_j| \in L(x, \bar{y}, z, w)$,

$$\bar{p} = p - i_0 \leq x\bar{y} + z + w, \quad \bar{p} \leq x\bar{u} + w \quad \text{and} \quad \bar{u} + \bar{y} \leq u + y - 1, \quad (1)$$

since by the induction assumptions we have

$$|\mathbf{D}| \leq n^{u+y+z-1} (p-i_0)^{2(p-i_0)(u+y-1)}$$

and

$$|\mathbf{F}| \leq p^{2p} + np^{2p} n^{u+y+z-1} (p-z)^{2(p-z)(u+y-1)} \leq n^{u+y+z} p^{2p(u+y)}.$$

It is straightforward to verify (1) by considering the following two cases.

Case 1. $x(y-1) + 2z + w < p \leq xy + z + w$, we can choose \bar{u}, \bar{y} satisfying $\bar{u} \leq y \leq u-1, \bar{y} \leq y$.

Case 2. $p \leq x(y-1) + 2z + w$. We can choose \bar{u}, \bar{y} satisfying $\bar{u} \leq u, \bar{y} \leq y-1$.

This completes the proof of Theorem 1.

Now, for any integer k , we construct a graph G such that the vertex set $V(G)$ consists of all p -subsets of $X = \{1, \dots, n\}$ and for $v_1, v_2 \in V(G)$, v_1 is adjacent to v_2 iff $|v_1 \cap v_2| \in \{2xx' + x'' : 0 \leq x', x'' < x\}$ where we choose

$$\begin{aligned} n &= \lceil \exp [(\log k)^{2/3} (\log \log k)^{1/3} / 3^{1/3}] \rceil, \\ x &= \left\lceil \frac{(3 \log k)^{1/3}}{4(\log \log k)^{1/3}} \right\rceil, \\ p &= 2x^2, \end{aligned} \quad (2)$$

with $\lceil y \rceil$ denoting the least integer greater than or equal to y . It follows Theorem 1 that neither G nor its complement \bar{G} contains any complete graph on k vertices and G has at least $\exp [c(\log k)^{4/3} / 8(\log \log k)^{1/3}]$ vertices. Therefore, we have the following result.

Theorem 2. The graph G constructed as above has at least $\exp [c(\log k)^{4/3} / (\log \log k)^{1/3}]$ vertices, for some constant c , and has the property that neither G nor its complement \bar{G} contains K_k . Therefore we have the constructive lower bound:

$$r(k) \geq \exp [c(\log k)^{4/3} / (\log \log k)^{1/3}].$$

REMARK 1. We note that the graph G we constructed by (2) contains a complete subgraph on $\exp [c'(\log k)^{4/3} / (\log \log k)^{1/3}]$ vertices for large k .

Therefore by using the above construction, the bound in Theorem 2 can not be improved asymptotically. Some new ideas will be needed in order to give a constructive lower bound of $(1 + \epsilon)^k$ for Ramsey number $r(k)$ for a fixed $\epsilon > 0$.

REMARK 2. Let $r_t(k)$ denote the least integer n such that if every edge of K_n is colored by one of t colors, then there exists a monochromatic K_k . It has been shown in [2] that $r_t(3) \geq c(3 + \delta)^t$ where $\delta = 0.103 \dots$ is the positive root of $x^3 + 6x^2 + 9x - 1 = 0$ and $c = 50\delta^2$. By a recursive construction used in [4], it can be shown that $r_t(k) \geq \exp[c't(\log k)^{4/3}/(\log \log k)^{1/3}]$ for sufficiently large k and some positive constant c' .

References

- [1] H. L. Abbott, Lower bounds for some Ramsey numbers. *Discrete Math.* 2 (1972) 289–293.
- [2] F. R. K. Chung, Ramsey numbers in multi-colors. Dissertation, University of Pennsylvania, 1974.
- [3] P. Erdős, Some remarks on the theory of graphs. *Bull. Amer. Math. Soc.* 53 (1947) 292–294.
- [4] P. Erdős, Problems and results in chromatic graph theory. in *Proof Techniques in Graph Theory*. Edited by F. Harary. Academic Press, London (1969) 27–35.
- [5] P. Erdős and J. Spencer, *Probabilistic Methods in Combinatorics*. Academic Press, New York (1974).
- [6] P. Erdős and R. Rado, Intersection theorems for systems of \mathcal{A} sets. *J. London Math. Soc.* 35 (1960) 85–90.
- [7] P. Frankl, A constructive lower bound for some Ramsey numbers. *Ars Combinatoria* 3 (1977) 297–302.
- [8] R. E. Greenwood and A. M. Gleason, Combinatorial relations and chromatic graphs. *Canad. J. Math.* 7 (1955) 1–7.
- [9] Zs. Nagy, A constructive estimation of the Ramsey numbers. *Mat. Lapok* 23 (1975) 301–302.
- [10] J. Spencer, Ramsey's theorem—A new lower bound. *J. Combinatorial Theory* 18 (1975) 108–115.
- [11] K. Walker, An upper bound for the Ramsey number $M(5, 4)$. *J. Combinatorial Theory* 11 (1971) 1–10.
- [12] J. Yackel, Inequalities and asymptotic bounds for Ramsey numbers. *J. Combinatorial Theory* 13 (1972) 56–68.
- [13] E. Levine and J. Winn, personal communication.