

Math 120A
August 29, 2022

Question 1 Given $R > 0$, let $\gamma = \{z \in \mathbb{C} \mid |z| = R\}$. Then,

A. γ can be parametrized by $z(t) = Re^{it}$ with $0 \leq t < 2\pi$.

B.
$$\int_{\gamma} |z^2| dz = \int_0^{2\pi} R^2 \cdot R ie^{it} dt = R^3 e^{it} \Big|_{t=0}^{2\pi} = 0.$$

C.
$$\int_{\gamma} |z^2| |dz| = \int_0^{2\pi} R^2 \cdot R dt = 2\pi R^3.$$

*D. All of the above.

E. None of the above. Complex line integrals can't be real.

Question 2 To develop a power series $\sum_{k=0}^{\infty} a_k(z-i)^k$ centered at $z_0 = i$ for $f(z) = \frac{1}{z-1}$, one could write $f(z) = -\frac{1}{1-z} = -(1-z)^{-1}$ and:

- A. Compute $f^{(k)}(z) = -k!(1-z)^{-(k+1)} = \frac{k!}{(1-z)^{k+1}}$ so that $f(z) = \sum_{k=0}^{\infty} \frac{f^{(k)}(i)}{k!}(z-i)^k = \sum_{k=0}^{\infty} -\frac{1}{(1-i)^{k+1}}(z-i)^k$.
- B. Write $f(z) = -\frac{1}{(1-i)-(z-i)} = -\frac{1}{1-i} \cdot \frac{1}{1-\left(\frac{z-i}{1-i}\right)}$ so that $f(z) = -\frac{1}{1-i} \sum_{k=0}^{\infty} \left(\frac{z-i}{1-i}\right)^k = \sum_{k=0}^{\infty} -\frac{1}{(1-i)^{k+1}}(z-i)^k$.
- *C. Either **A** or **B**. Both are great ways to develop the power series.
- D. Neither **A** nor **B**. $f(z) = \frac{1}{z-1}$ is not analytic on any disk centered at $z_0 = i$.
- E. This selection has intentionally been left blank.

Question 3 The function $f(z) = \frac{1}{z} + \frac{1}{z^5}$ can be written $f(z) = \frac{z^4+1}{z^5}$.

We can conclude that

- A. $\frac{1}{z} + \frac{1}{z^5}$ is the Laurent series of f for $|z| > 0$.
- B. $f(z)$ has four simple zeros: $z \in \left\{ e^{i\frac{\pi}{4}}, e^{i\frac{3\pi}{4}}, e^{i\frac{5\pi}{4}}, e^{i\frac{7\pi}{4}} \right\}$.
- C. $f(z)$ has a simple zero at ∞ .
- D. $g(w) = f(1/w) = w(1 + w^4)$ has a simple zero at $w = 0$.
- *E. All of the above.

Question 4 $\text{Log}(z)$ is not analytic at $z_0 = 0$. $z_0 = 0$ is called

- A. an isolated singularity of $\text{Log}(z)$
- *B. a branch point of $\text{Log}(z)$
- C. an essential singularity of $\text{Log}(z)$
- D. **A** and **B**
- E. **A** and **C**

Question 5 $f(z)$ has an isolated singularity at ∞ if

- A. $f(z)$ is analytic outside some bounded set.
- B. there is $R > 0$ such that $f(z)$ is analytic for $|z| > R$.
- C. $g(w) = f(1/w)$ has an isolated singularity at $w = 0$.
- D. **A** and **B**; they are the same.
- *E. **A**, **B**, and **C**; they are equivalent.

Question 6 $f(z)$ has a pole of order N at infinity if

- A. $P_\infty(z)$, the principal part of $f(z)$ at ∞ , is a polynomial of degree N ; i.e., $P_\infty(z) = b_N z^N + b_{N-1} z^{N-1} + \cdots + b_1 z + b_0$.
- B. $g(w) = f(1/w)$ has a pole of order N at $w = 0$.
- C. $g(w) = f(1/w)$ has a zero of order N at $w = 0$.
- *D. **A and B**
- E. **A and C**