

Math 120A  
August 24, 2023

**Question 1** Cauchy's integral formula states that a function  $f(z)$  analytic on a bounded domain  $D$  and smooth on  $D \cup \partial D$  satisfies

$$f(z) = \frac{1}{2\pi i} \int_{\partial D} \frac{f(w)}{w - z} dw, \quad z \in D.$$

This is important because

- A. the integral can be differentiated with respect to  $z$ .
- B. by induction, one sees that for every  $z \in D$ ,  
$$f^{(m)}(z) = \frac{m!}{2\pi i} \int_{\partial D} \frac{f(w)}{(w - z)^{m+1}} dw$$
 for all integers  $m \geq 0$ .
- C. one can conclude that a function  $f(z)$  that has one complex derivative at each  $z \in D$  must have complex derivatives of all orders at each  $z \in D$ .
- \*D. All of the above, and it's absolutely amazing.
- E. None of the above, Cauchy's integral formula is as useless as it is aesthetically pleasing.

**Question 2** Let  $\gamma$  be the curve  $|z| = 2$  with positive (counterclockwise) orientation. Then the integral  $\int_{\gamma} \frac{z^n}{z-3} dz$

- \*A. is equal to 0 by Cauchy's theorem.
- B. is equal to  $3^n$  by the Cauchy integral theorem.
- C. is equal to  $2\pi i 3^n$  by the Cauchy integral theorem.
- D. is undefined because  $\frac{z^n}{z-3}$  is undefined at  $z = 3$ .
- E. none of the above.

**Question 3** Let  $\gamma$  be the curve  $|z| = 2$  with positive (counterclockwise) orientation. Then the integral  $\int_{\gamma} \frac{z^n}{z-1} dz$

- A. is equal to 0 by Cauchy's theorem.
- B. is equal to 1 by the Cauchy integral theorem.
- \*C. is equal to  $2\pi i$  by the Cauchy integral theorem.
- D. is undefined because  $\frac{z^n}{z-1}$  is undefined at  $z = 1$ .
- E. none of the above.

**Question 4** Let  $\gamma$  be the curve  $|z| = 2$  with positive (counterclockwise) orientation. Then the integral  $\int_{\gamma} \frac{z^n}{z+2} dz$

- A. is equal to 0 by Cauchy's theorem.
- B. is equal to  $(-2)^n$  by the Cauchy integral theorem.
- C. is equal to  $2\pi i (-2)^2$  by the Cauchy integral theorem.
- \*D. is undefined because  $\frac{z^n}{z+2}$  is undefined at  $z = -2$ .
- E. none of the above.

**Question 5** The functions  $f_k : [0, 1] \rightarrow \mathbb{R}$  given by  $f_k(x) = x^k$

- A. are all continuous.
- B. converge pointwise to a discontinuous function.
- C. converge uniformly to a discontinuous function.
- \*D. **A** and **B**.
- E. all of the above; if they converge uniformly, they also converge pointwise.