

4. SOME EXAMPLES

We present some examples of varieties, mainly surfaces, with interesting Mori cones.

Definition 4.1. Let $\mathcal{C} \subset V \simeq \mathbb{R}^n$ be a subset of a finite dimensional real vector space. We say that $\mathcal{C}$ is a **cone** (respectively **convex subset**) if whenever α and $\beta \in \mathcal{C}$ then

$$\lambda\alpha + \mu\beta \in \mathcal{C} \quad \text{for all} \quad \lambda \geq 0, \mu \geq 0,$$

(respectively such that $\lambda + \mu = 1$). We say that $\mathcal{C}$ is **strictly convex** if $\mathcal{C}$ contains no positive dimensional linear subspaces.

We say that $R \subset \mathcal{C}$ is a **ray** of a cone $\mathcal{C}$ if $R = \mathbb{R}^+\alpha$, for some non-zero vector $\alpha \in \mathcal{C}$. We say that R is an **extremal ray** if whenever $\beta + \gamma \in R$, where β and $\gamma \in \mathcal{C}$, then β and $\gamma \in R$.

It follows by Kleiman's criteria that the closed cone of curves of a projective variety is strictly convex.

Lemma 4.2. Let S be a smooth projective surface and let $\alpha = \sum a_i[C_i]$ and $\beta = \sum b_j[D_j]$ be two cycles, where $a_i > 0$ and $b_j > 0$, so that α and β belong to the cone of curves.

If $\alpha \cdot \beta < 0$ then $\Sigma = C_i = D_j$ for some i and j , where $\Sigma^2 < 0$. If i and j are the only two indices with this property then $a_i b_j \Sigma^2 \leq \alpha \cdot \beta$.

Proof. Clear, since

$$\alpha \cdot \beta = \sum a_i b_j C_i \cdot D_j,$$

and $C_i \cdot D_j \geq 0$ unless $C_i = D_j$. □

Lemma 4.3. Let S be a smooth projective surface. Let

- (1) If $R = \mathbb{R}^+\alpha \subset \overline{\text{NE}}(S)$ is an extremal ray of the closed cone of curves of S then $\alpha^2 \leq 0$.
- (2) If C is an irreducible curve such that $C^2 < 0$ then $R = \mathbb{R}^+[C]$ is extremal.

Proof. We will first show that $R = \mathbb{R}^+\alpha$ is never extremal if $\alpha^2 > 0$. Let H be an ample divisor. Then $H \cdot \alpha > 0$ by Kleiman's criteria. Pick a small neighbourhood U of $\alpha \in V$ such that

- $\beta^2 > 0$
- $\beta \cdot H > 0$,

for all $\beta \in U$. Suppose that $\beta \in U$ is rational. Pick $k \in \mathbb{N}$ such that $D = k\beta$ is integral. By Asymptotic Riemann-Roch,

$$\begin{aligned} h^0(S, \mathcal{O}_S(mD)) + h^0(S, \mathcal{O}_S(K_S - mD)) &= h^0(S, \mathcal{O}_S(mD)) + h^2(S, \mathcal{O}_S(mD)) \\ &\geq \chi(S, \mathcal{O}_S(mD)) > 0, \end{aligned}$$

for large m . Thus either $|mD|$ or $|K_S - mD|$ is non-empty for large m . But $(K_S - mD) \cdot H < 0$ for large m so that $|K_S - mD|$ is empty and so $|mD|$ is non-empty. Thus $[D] \in \text{NE}(S)$ and so $U \subset \overline{\text{NE}}(S)$. In particular R is not extremal.

Now suppose that C is an irreducible curve such that $C^2 \leq 0$. Then we may write

$$[C] = \sum \beta_i$$

where the β_i generate extremal rays of $\overline{\text{NE}}(S)$. As

$$0 > C^2 = [C] \cdot \left(\sum \beta_i \right).$$

it follows that $[C] \cdot \beta_i < 0$ for some i . Since β_i is a limit of $\beta(j) \in \text{NE}(S)$, (4.2) implies that $\beta_i = \lambda[C] + \beta'$, where $\beta' \in \overline{\text{NE}}(S)$ and $\lambda > 0$. As β_i generates an extremal ray, it follows that $\mathbb{R}^+ \beta_i = \mathbb{R}^+ [C] = R$ is extremal. $\square$

Let $\pi: X \rightarrow C$ be a $\mathbb{P}^r$ -bundle over a smooth curve. We recall the classification of such bundles. We have that $S = \mathbb{P}(E)$, for some rank $r + 1$ vector bundle over C and the two $\mathbb{P}^r$ -bundles $S_i = \mathbb{P}(E_i)$ are isomorphic over C if and only if there is a line bundle L and an isomorphism of vector bundles $E_1 \otimes L \simeq E_2$.

Now the Picard group of X has rank two, and it is generated by the class of a line in a fibre and the class of any section. Thus $\overline{\text{NE}}(X) \subset \mathbb{R}^2$. Taking a compact slice, we get a closed interval, so that topologically the situation is an open and closed book. To get a complete description, we have two rays R_1 and R_2 and it suffices to determine generators for each ray.

First suppose that $C = \mathbb{P}^1$, so that $X = S$ is a rational surface, a $\mathbb{P}^1$ -bundle over $\mathbb{P}^1$. In this case any rank two vector bundle E has the form $\mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(b)$ by a Theorem of Grothendieck. We may normalise so that $E = \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-n)$. The resulting surface is denoted $\mathbb{F}_n$. Let f denote the class of a fibre and e the class of a section of minimal self-intersection $-n$. Then

$$\overline{\text{NE}}(X) = \mathbb{R}^+ f + \mathbb{R}^+ e.$$

Note that in this case there is a perfect correspondence between faces of the Mori cone and contraction morphisms. There are four faces. The extremal ray spanned by f corresponds to the map down to $\mathbb{P}^1$. The contracted curves are the fibres of π . The extremal ray spanned by the curve of negative self-intersection corresponds to a birational morphism $\phi: S \rightarrow T$, where $T \subset \mathbb{P}^{n+1}$ is the cone over a rational normal curve $\Sigma \subset \mathbb{P}^n$. The only curve contracted by ϕ is Σ the curve of negative self-intersection $-n$. If H is a hyperplane in $\mathbb{P}^{n+1}$ then $D = \phi^* H$ is zero

on Σ and is a curve of self-intersection n . The case $n = 0$ is slightly special. In this case $S = \mathbb{P}^1 \times \mathbb{P}^1$, $e^2 = 0$ and ϕ is projection to the other factor.

The other two faces are the origin, which corresponds to the identity (no curves are contracted) and the whole of $H_2(S, \mathbb{R})$ which corresponds to the map down to a point (all curves are contracted).

But now suppose that C has higher genus. We need to say something about all sections and multisections of π . Fortunately in characteristic zero we only need to keep track of the sections. We recall some of the theory of vector bundles:

Definition 4.4. Let $f: E \rightarrow C$ be a vector bundle over a curve. The **slope** of E is the rational number

$$\mu(E) = \frac{\deg E}{\operatorname{rk} E}.$$

We say that E is **stable** (respectively **semistable**) if for all quotient vector bundles $E \rightarrow F$ of E , we have

$$\mu(E) < \mu(F) \quad (\text{respectively} \quad \mu(E) \leq \mu(F)).$$

We say that E is **unstable** if it is not semistable. We say that F **destabilises** E if F is a quotient $E \rightarrow F$ of E and

$$\mu(F) < \mu(E).$$

The maximal **destabilising quotient** is a quotient vector bundle with the smallest slope and the largest rank amongst quotients with the same slope.

Example 4.5. Suppose that

$$E = \bigoplus_{i=0}^r L_i,$$

is a direct sum of line bundles.

If $\deg L_i = d_i$, then

$$\mu(E) = \frac{d_0 + d_1 + \cdots + d_r}{r+1} \quad \text{and} \quad \mu(L_i) = d_i.$$

As $F = L_i$ is a quotient of E , E is never stable and it is semistable if and only if $\deg L_i$ is independent of i . Let $m = \min_i d_i$. If $m \neq \mu(E)$, then the maximal destabilising quotient is

$$\bigoplus_{i: d_i = m} L_i.$$

Lemma 4.6. *Let E be a vector bundle over a smooth curve C .*

*Then E is semistable (respectively stable) if and only if $E' = f^*E$ is semistable (respectively stable) for all covers $f: C' \rightarrow C$.*

Proof. We first do the semistable case.

One direction is clear; if $E \rightarrow F$ destabilises E then $E' \rightarrow F' = f^*F$ destabilises E' .

Suppose that E' is not semistable. By what we already proved, passing to a finite cover of C' we may assume that f is Galois, with Galois group $G \subset \text{Aut}(C')$. Let $E' \rightarrow F'$ be a maximal destabilising subsheaf. Then F' is canonical, whence invariant under the action of $G \subset \text{Aut}(C')$. But this means that $F' = f^*F$ for some vector bundle F and F destabilises E .

We now turn to the stable case. As before, one direction is clear and as result for the other direction we reduce to the case that f is Galois and E' is not stable. In this case let F' be the socle of E' , the largest quotient of E' with the same slope as E' . Then the socle is invariant under G and so descends to a quotient of E with the same slope. But then E is semistable but not stable. $\square$

Remark 4.7. *It is not true that if V is an arbitrary vector bundle on C' which is invariant under G then $V = f^*W$, for some vector bundle W on C .*

In fact this can fail even for line bundles. For example, suppose let $f: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be the standard double cover, given by $[S : T] \rightarrow [S^2 : T^2]$. This map is Galois, with Galois group $G = \mathbb{Z}_2 \subset \text{Aut}(\mathbb{P}^1)$ where the action is given by $[S : T] \rightarrow [S : -T]$. Then the line bundle $L = \mathcal{O}_{\mathbb{P}^1}(1)$ is invariant, as it is sent to another line bundle of degree one and all such line bundles are isomorphic. However the degree of any line bundle pulled back under f must have even degree.

We need the fact F' is a quotient of $E' = f^*E$. In this case the action of G on F' is compatible with the action on E' so that we can construct F as a quotient of E .

Lemma 4.8. *Let C be a smooth curve and let $\pi: X = \mathbb{P}(E) \rightarrow C$ be a $\mathbb{P}^r$ -bundle over C . Suppose that $r > 1$.*

The following are equivalent:

- (1) *E is stable (respectively semistable).*
- (2) *$-K_{X/C} \cdot \Sigma > 0$ (respectively ≥ 0) for all curves $\Sigma \subset X$, where $K_{X/C} = K_X - \pi^*K_C$ is the relative canonical divisor.*

Example 4.9. *Let C be a curve of genus at least two. Suppose that E is a general rank two stable vector bundle, which admits an exact*

sequence

$$0 \longrightarrow \mathcal{O}_C \longrightarrow E \longrightarrow L \longrightarrow 0,$$

where L is a line bundle of positive degree $d \leq g$ (such exist by general theory).

Let $S = \mathbb{P}(E)$. If E is chosen generically (see for example [2]) then there is a section C_0 of minimal self-intersection d .

Suppose that Σ is a multi-section of S , so that $f: D = \Sigma \longrightarrow C$ is dominant. I claim that $\Sigma^2 > 0$. Let $Y = \mathbb{P}(f^*E)$. Then

$$f^*\Sigma = \Sigma_0 + \Sigma_1,$$

where Σ_0 is a section of $Y \longrightarrow D$. But then

$$\Sigma^2 = \Sigma_0 \cdot f^*\Sigma \geq \Sigma_0^2.$$

As f^*E is stable we may replace X by Y and Σ by Σ_0^2 , so that we may assume that Σ is a section of $X \longrightarrow C$. But then

$$(K_X + \Sigma) \cdot \Sigma = K_\Sigma = \pi^*K_C.$$

Thus

$$\Sigma^2 = -K_{X/C} \cdot \Sigma > 0,$$

by assumption. Consider $\overline{\text{NE}}(S)$. One edge is given by f the class of a fibre. What about the other edge? Suppose that this is generated by α . All curves other than F have positive self-intersection. By (4.3) we must have $\alpha^2 = 0$. Thus $\text{NE}(S)$ is not a closed cone.

Rescaling we may suppose that $\alpha = \sigma - af$, where σ is the class of C_0 and $a \geq 0$. We have

$$\begin{aligned} 0 &= \alpha^2 \\ &= (C_0 - aF)^2 \\ &= C_0^2 - 2aC_0 \cdot F + a^2F^2 \\ &= d - 2a. \end{aligned}$$

Thus the divisor $D = 2\Sigma - dF = -K_{X/C}$ is a divisor which intersects every curve positively, but which is not ample. This gives an example in characteristic zero where Kleiman's criteria is sharp.

Example 4.10. Let $S = E \times E$ the product of two general elliptic curves.

Then $\rho(S) = 3$ (it might be larger if E is not general). Let f_i be the class of a fibre and let δ be the class of the diagonal. Suppose that

$\delta = a_1 f_1 + a_2 f_2$. Then

$$\begin{aligned} a_i &= (a_1 f_1 + a_2 f_2) \cdot f_{2-i} \\ &= \delta \cdot f_{2-i} \\ &= 1. \end{aligned}$$

But

$$\begin{aligned} 0 &= \deg K_E \\ &= (K_S + \Delta) \cdot \Delta \\ &= K_S \cdot \Delta + \delta^2 \\ &= \delta^2 \\ &= (f_1 + f_2)^2 \\ &= 2. \end{aligned}$$

Thus f_1 and f_2 and δ define independent classes, which actually span the Néron-Severi group.

On the other hand let $D \geq 0$ be a $\mathbb{Q}$ -Cartier divisor. Then $D^2 \geq 0$, as can be seen by acting by a general translation. Thus $\overline{\text{NE}}(S)$ is half of a classical cone. There are uncountably many extremal rays, and at most countably many contractions. Most rays are not rational.

Example 4.11. *Let S be obtained from $\mathbb{P}^2$ by blowing up nine points $p_1, p_2, \dots, p_9$.*

Suppose first that these points are the nine points of the intersection of two general smooth cubics. Then S is the total space of the pencil, and there is a morphism $f: S \rightarrow \mathbb{P}^1$ whose fibres are the elements of the pencil. The nine exceptional divisors $E_1, E_2, \dots, E_9$ are then sections of this fibration. The generic fibre C of f is an elliptic curve (over the function field $\mathbb{C}(\mathbb{P}^1) = \mathbb{C}(t)$) and the nine sections define nine points $e_1, e_2, \dots, e_9$. Since the pencil is general it follows that these points generate a subgroup of C , isomorphic to $\mathbb{Z}^8$. This subgroup then corresponds to a subgroup of the automorphism group of S over the base. The orbit of the nine exceptional divisors gives infinitely many exceptional divisors. Each exceptional divisor generates an extremal ray of the closed cone of curves.

What is worse, -1 -curves persist under small deformations. If we therefore perturb these nine points to nine very general points, infinitely many of these -1 -curves survive. The resulting surface does not have any automorphisms, and yet its Mori cone is still very complicated.

Let us end these series of examples with Zariski's famous example:

Example 4.12. *Pick a smooth cubic curve C in $\mathbb{P}^2$ and let S be the blow up of $f: S \rightarrow \mathbb{P}^2$ at ten very general points $p_1, p_2, \dots, p_{10}$ of C .*

Let $E_1, E_2, \dots, E_{10}$ be the ten exceptional divisors and let Σ be the strict transform of C .

Then Σ is a curve of self-intersection $9 - 10 = -1$. By a result due to Artin, there is a contraction morphism $\pi: S \rightarrow T$ contracting Σ , where T is a normal algebraic space (or if you will, an analytic space). I claim that T is not a projective variety.

Suppose it were. Then T would have an ample divisor D . But I claim it has no Cartier divisors at all. We have

$$\pi^*D \in A_1(S) = \mathbb{Z}^{11} = \mathbb{Z}[f^*H] + \sum_{i=1}^{10} \mathbb{Z}[E_i],$$

where H is a line in $\mathbb{P}^2$. So

$$\pi^*D \sim d\pi^*H - \sum a_i E_i.$$

Now

$$\pi^*D \cdot \Sigma = D \cdot \pi_*\Sigma = 0.$$

Thus

$$3d = \sum a_i.$$

Moreover $\pi^*D|_{\Sigma}$ would be linearly equivalent to zero. Thus there would be some curve B of degree d in $\mathbb{P}^2$ such that

$$B|_C \sim \sum b_i p_i.$$

But this contradicts the fact that our ten points of C are general. Indeed the points $p_1, p_2, \dots, p_{10}$ determine points $\mathcal{O}_C(p_i - q)$ in $\text{Pic}^0(C) \simeq C \simeq S^1 \times S^1$, where the last isomorphism is in the category of topological groups. If the points are general enough then we get no unexpected relations.

Note that there are then plenty of nef divisors D on S which are zero on C but which are not semiample (since if D were semiample, it would descend to T).

Definition 4.13. *A **stable n -pointed curve of genus g** is a connected curve of arithmetic genus g , with only nodes as singularities, with n marked points contained in the smooth locus, such that the normalisation of every component isomorphic to $\mathbb{P}^1$ has three special points (either a node or a marked point).*

The moduli space of genus g , n -pointed stable curves $\overline{M}_{g,n}$ is a projective variety whose points are in natural correspondence with isomorphism classes of stable n -pointed curves of genus g . Note that

$\overline{M}_{1,1} \simeq \overline{M}_{0,4} \simeq \mathbb{P}^1$. Indeed, the second isomorphism is given by the j -invariant, and the first is a consequence since an elliptic curve double covers $\mathbb{P}^1$ over 4 points.

These gives finitely many rational curves in $\overline{M}_{g,n}$, which we call **vital curves**. Indeed take a stable curve of genus $g-1$ with n points (respectively a stable curve of genus g with $n-3$ points) whose components are copies of $\mathbb{P}^1$ with 3 labelled points, together with one component with only two labelled points. Now attach an elliptic to the special component (or a copy of $\mathbb{P}^1$ with three marked points). The resulting curve is a point of $\overline{M}_{g,n}$. Varying the moduli of the elliptic curve (or of the four points), gives a curve $C \subset \overline{M}_{g,n}$.

Conjecture 4.14 (Faber, Fulton, Mumford). *$\overline{M}_{g,n}$ is spanned by the classes of the vital curves.*

Theorem 4.15 (Gibney, Keel, Morrison: [1]). *To prove (4.14) it suffices to prove the case when $g = 0$.*

REFERENCES

- [1] A. Gibney, S. Keel, and I. Morrison, *Towards the ample cone of $\overline{M}_{g,n}$* , J. Amer. Math. Soc. **15** (2002), no. 2, 273–294.
- [2] R. Hartshorne, *Ample subvarieties of algebraic varieties*, Notes written in collaboration with C. Musili. Lecture Notes in Mathematics, Vol. 156, Springer-Verlag, Berlin, 1970.