

3. RING HOMOMORPHISMS AND IDEALS

Definition 3.1. Let $\phi: R \rightarrow S$ be a function between two rings. We say that ϕ is a **ring homomorphism** if for every a and $b \in R$,

$$\begin{aligned}\phi(a + b) &= \phi(a) + \phi(b) \\ \phi(a \cdot b) &= \phi(a) \cdot \phi(b),\end{aligned}$$

and in addition $\phi(1) = 1$.

Note that this gives us a category, the category of rings. The objects are rings and the morphisms are ring homomorphisms. Just as in the case of groups, one can define automorphisms.

Example 3.2. Let $\phi: \mathbb{C} \rightarrow \mathbb{C}$ be the map that sends a complex number to its complex conjugate. Then ϕ is an automorphism of \mathbb{C} . In fact ϕ is its own inverse.

Let $\phi: R[x] \rightarrow R[x]$ be the map that sends $f(x)$ to $f(x + 1)$. Then ϕ is an automorphism. Indeed the inverse map sends $f(x)$ to $f(x - 1)$.

By analogy with groups, we have

Definition 3.3. Let $\phi: R \rightarrow S$ be a ring homomorphism.

The **kernel** of ϕ , denoted $\text{Ker } \phi$, is the inverse image of zero.

As in the case of groups, a very natural question arises. What can we say about the kernel of a ring homomorphism? Since a ring homomorphism is automatically a group homomorphism, it follows that the kernel is a normal subgroup. However since a ring is an abelian group under addition, in fact all subgroups are automatically normal.

Definition-Lemma 3.4. Let R be a ring and let I be a subset of R . We say that I is an **ideal** of R and write $I \triangleleft R$ if I is an additive subgroup of R and for every $a \in I$ and $r \in R$, we have

$$ra \in I \quad \text{and} \quad ar \in I.$$

Let $\phi: R \rightarrow S$ be a ring homomorphism and let I be the kernel of ϕ . Then I is an ideal of R .

Proof. We have already seen that I is an additive subgroup of R .

Suppose that $a \in I$ and $r \in R$. Then

$$\begin{aligned}\phi(ra) &= \phi(r)\phi(a) \\ &= \phi(r)0 \\ &= 0.\end{aligned}$$

Thus ra is in the kernel of ϕ . Similarly for ar . □

As before, given an additive subgroup H of R , we let R/H denote the group of left cosets of H in R .

Proposition 3.5. *Let R be a ring and let I be an ideal of R , such that $I \neq R$.*

Then R/I is a ring. Furthermore there is a natural ring homomorphism

$$u: R \longrightarrow R/I$$

which sends a to $a + I$.

Proof. As I is an ideal, and addition in R is commutative, it follows that R/I is a group, with the natural definition of addition inherited from R . Further we have seen that u is a group homomorphism. It remains to define a multiplication in R/I .

Given two left cosets $a + I$ and $b + I$ in R/I , we define multiplication in the obvious way,

$$(a + I)(b + I) = ab + I.$$

In fact this is forced by requiring that u be a ring homomorphism.

As before the problem is to check that this is well-defined. Suppose that $a' + I = a + I$ and $b' + I = b + I$. Then we may find i and j in I such that $a' = a + i$ and $b' = b + j$. We have

$$\begin{aligned} a'b' &= (a + i)(b + j) \\ &= ab + aj + ib + ij. \end{aligned}$$

As I is an ideal, $ia + bj + ij \in I$. It follows that $a'b' + I = ab + I$ and multiplication is well-defined. The rest is easy to check. \square

As before the quotient of a ring by an ideal is a categorical quotient.

Theorem 3.6. *Let R be a ring and I an ideal not equal to all of R . Let $u: R \longrightarrow R/I$ be the natural map. Then u is universal amongst all ring homomorphisms whose kernel contains I .*

That is, suppose $\phi: R \longrightarrow S$ is any ring homomorphism, whose kernel contains I . Then there is a unique ring homomorphism $\psi: R/I \longrightarrow S$, which makes the following diagram commute,

$$\begin{array}{ccc} R & \xrightarrow{\phi} & S \\ u \downarrow & \nearrow & \\ R/I & & \end{array}$$

Proof. Since ϕ is a group homomorphism the existence and uniqueness of the induced map ψ is clear; this follows by invoking the universal property of R/I as a categorical group quotient. From there it is straightforward to check that ψ is a ring homomorphism. \square

Theorem 3.7 (Isomorphism Theorem). *Let $\phi: R \rightarrow S$ be a homomorphism of rings. Suppose that ϕ is onto and let I be the kernel of ϕ .*

Then S is isomorphic to R/I .

Example 3.8. *Let $R = \mathbb{Z}$. Fix a non-zero integer n and let I consist of all multiples of n . It is easy to see that I is an ideal of \mathbb{Z} . The quotient, \mathbb{Z}/I is \mathbb{Z}_n the ring of integers modulo n .*

Definition-Lemma 3.9. *Let R be a commutative ring and let $a \in R$ be an element of R .*

The set

$$I = \langle a \rangle = \{ ra \mid r \in R \},$$

*is an ideal and any ideal of this form is called **principal**.*

Proof. We first show that I is an additive subgroup.

Suppose that x and y are in I . Then $x = ra$ and $y = sa$, where r and s are two elements of R . In this case

$$\begin{aligned} x + y &= ra + sa \\ &= (r + s)a. \end{aligned}$$

Thus I is closed under addition. Further $-x = -ra = (-r)a$, so that I is closed under inverses. It follows that I is an additive subgroup.

Now suppose that $x \in I$ and that $s \in R$. Then

$$\begin{aligned} sx &= s(ra) \\ &= (sr)a \in I. \end{aligned}$$

It follows that I is an ideal. \square

Definition-Lemma 3.10. *Let R be a ring. We say that $u \in R$ is **invertible**, if u has a multiplicative inverse.*

Let I be an ideal of a ring R . If I contains an invertible element, then $I = R$.

Proof. Suppose that $u \in I$ is an invertible element of R . Then $vu = 1$, for some $v \in R$. It follows that

$$1 = vu \in I.$$

Pick $a \in R$. Then

$$a = a \cdot 1 \in I. \quad \square$$

Proposition 3.11. *Let R be a division ring. Then the only ideals of R are the zero ideal and the whole of R .*

In particular, if $\phi: R \rightarrow S$ is any ring homomorphism then ϕ is injective.

Proof. Let I be an ideal, not equal to $\{0\}$. Pick $u \in I$, $u \neq 0$. As R is a division ring, it follows that u is invertible. But then $I = R$.

Now let $\phi: R \rightarrow S$ be a ring homomorphism and let I be the kernel. Then I cannot be the whole of R , so that $I = \{0\}$. But then ϕ is injective. \square

Example 3.12. *Let X be a set and let R be a ring.*

Let F denote the set of functions from X to R . We have already seen that F forms a ring, under pointwise addition and multiplication.

Let Y be a subset of X and let I be the set of those functions from X to R whose restriction to Y is zero.

Then I is an ideal of F . Indeed I is clearly non-empty as the zero function is an element of I . Given two functions f and g in F , whose restriction to Y is zero, then clearly the restriction of $f + g$ to Y is zero. Finally, suppose that $f \in I$, so that f is zero on Y and suppose that g is any function from X to R . Then gf is zero on Y . Thus I is an ideal.

Now consider F/I . I claim that this is isomorphic to the space of functions G from Y to R . Indeed there is a natural map from F to G which sends a function to its restriction to Y ,

$$f \longrightarrow f|_Y.$$

It is clear that the kernel is I . Thus the result follows by the Isomorphism Theorem. Of course this gives an easy way to check I is an ideal.

As a special case, one can take $X = [0, 1]$ and $R = \mathbb{R}$. Let $Y = \{1/2\}$. Then the space of maps from Y to \mathbb{R} is just a copy of \mathbb{R} .

Example 3.13. *Let R be the ring of Gaussian integers, that is, those complex numbers of the form $a + bi$, where a and b are integers.*

Let I be the subset of R consisting of those numbers such $2|a$ and $2|b$. I claim that I is an ideal of R . In fact suppose that $a + bi \in I$ and $c + di \in I$. Then

$$(a + bi) + (c + di) = (a + c) + (b + d)i.$$

As a and c are even, then so is $a + c$ and similarly as b and d are even, then so is $b + d$.

Thus I is an additive subgroup. On the other hand, if $a + bi \in I$ and $c + di \in R$ then

$$(c + di)(a + bi) = (ac - bd) + (bc + ad)i.$$

By assumption a and b are even. It is easy to see that $ac - bd$ and $bc + ad$ are even, so that the product is in I and so I is an ideal.

In fact

$$I = \langle 2 \rangle.$$