

Math 280B, Winter 2005

SLLN for Martingales

Everything that follows takes place on a probability space $(\Omega, \mathcal{F}, \mathbf{P})$ equipped with a filtration $\{\mathcal{F}_n : n = 0, 1, 2, \dots\}$, with $\mathcal{F}_n \subset \mathcal{F}$ for all n .

1. Square-integrable martingales. A martingale $M = (M_n)$ is said to be *square integrable* if $\mathbf{E}[M_n^2] < \infty$ for each n . In this case M_n^2 is a submartingale, with Doob decomposition

$$M_n^2 = M_0^2 + X_n + \langle M \rangle_n, \quad n \geq 0,$$

where (X_n) is a martingale with $X_0 = 0$ and the (predictable) *quadratic variation* process $\langle M \rangle$ is defined by $\langle M \rangle_0 = 0$ and

$$(1) \quad \langle M \rangle_n := \sum_{k=1}^n \mathbf{E}[(\Delta M_k)^2 | \mathcal{F}_{k-1}], \quad n \geq 1.$$

A variation of a stopping time argument used in class proves the following result. We write $\langle M \rangle_\infty := \lim_n \langle M \rangle_n$, and $\{M_n \rightarrow\}$ for the event on which (M_n) converges to a real-valued limit.

2. Theorem. *Let $M = (M_n)$ be a square-integrable martingale with quadratic variation $\langle M \rangle$ as defined in (1).*

(a) $\{\langle M \rangle_\infty < \infty\} \subset \{M_n \rightarrow\}$, a.s.

(b) If $\mathbf{E}[\sup_n (\Delta M_n)^2] < \infty$, then $\{\langle M \rangle_\infty < \infty\} = \{M_n \rightarrow\}$, a.s.

Proof. (a) For $b > 0$ define $T_b := \inf\{n : \langle M \rangle_{n+1} > b\}$. Then T_b is a stopping time (because $\langle M \rangle$ is predictable), and $\langle M \rangle_{T_b} \leq b$. The stopped process $X_{T_b \wedge n}$ is a martingale, so

$$\mathbf{E}[M_{T_b \wedge n}^2] = \mathbf{E}[X_0] + \mathbf{E}[\langle M \rangle_{T_b \wedge n}] = \mathbf{E}[M_0^2] + \mathbf{E}[\langle M \rangle_{T_b \wedge n}] \leq \mathbf{E}[M_0^2] + b.$$

That is, the stopped martingale $(M_{T_b \wedge n})$ is L^2 -bounded. Thus, $(M_{T_b \wedge n})$ converges a.s., so (M_n) converges a.s. on the event $\{T_b = \infty\}$. Varying b over the positive integers, we see that (M_n) converges a.s. on the event $\cup_{b \in \mathbf{N}} \{T_b = \infty\} = \{\langle M \rangle_\infty < \infty\}$.

(b) Define, for $b > 0$, stopping times $S_b := \inf\{n : M_n^2 > b\}$, and note that $|M_{S_b-1}| \leq \sqrt{b}$. Consequently, on the event $\{S_b < \infty\}$,

$$\begin{aligned} |\Delta(M_{S_b}^2)| &= |M_{S_b}^2 - M_{S_b-1}^2| \leq (\Delta M_{S_b})^2 + 2|M_{S_b-1}| \cdot |\Delta M_{S_b}| \\ &\leq (\Delta M_{S_b})^2 + 2\sqrt{b} \cdot |\Delta M_{S_b}|. \end{aligned}$$

Thus,

$$\mathbf{E}[|\Delta(M_{S_b}^2)|; S_b < \infty] \leq C + 2\sqrt{b} + \sqrt{C} < \infty,$$

where $C := \mathbf{E}[\sup_n (\Delta M_n)^2] < \infty$. Now, because $M_n^2 - \langle M \rangle_n$ is a martingale,

$$\mathbf{E}[\langle M \rangle_{S_b \wedge n}] = \mathbf{E}[M_{S_b \wedge n}^2] - \mathbf{E}[M_0^2] \leq \mathbf{E}[M_{S_b \wedge n}^2] \leq b^2 + \mathbf{E}[|\Delta(M_{S_b}^2)|; S_b < \infty] < \infty,$$

and so, by Fatou's lemma,

$$\mathbf{E}[\langle M \rangle_{S_b}] \leq b^2 + \mathbf{E}[|\Delta(M_{S_b}^2)|; S_b < \infty] < \infty.$$

Thus $\langle M \rangle_{S_b} < \infty$ a.s. In particular, $\langle M \rangle_\infty < \infty$ on $\{S_b = \infty\}$. Varying $b > 0$ we see that $\langle M \rangle_\infty < \infty$ on the event $\cup_{b=1}^\infty \{S_b = \infty\} = \{\sup_n M_n^2 < \infty\} \supset \{M_n \rightarrow\}$. \square

As a corollary of Theorem 2 we have the following Strong Law of Large Numbers for square-integrable martingales.

3. Theorem. *Let $M = (M_n)$ be a square-integrable martingale with quadratic variation $\langle M \rangle$ as defined in (1). Then*

$$\frac{M_n}{\langle M \rangle_n} \rightarrow 0 \quad \text{a.s. on the event } \{\langle M \rangle_\infty = \infty\}.$$

Proof. On $\{\langle M \rangle_\infty = \infty\}$, we have

$$\frac{\langle M \rangle_n}{1 + \langle M \rangle_n} \rightarrow 1 \text{ as } n \rightarrow \infty.$$

It therefore suffices to show that, a.s. on $\{\langle M \rangle_\infty = \infty\}$,

$$(2) \quad \frac{M_n}{1 + \langle M \rangle_n} \rightarrow 0.$$

In view of Kronecker's lemma, to show (2) it suffices to show that

$$K_n := \sum_{k=1}^n \frac{\Delta M_k}{1 + \langle M \rangle_k}$$

converges as $n \rightarrow \infty$, a.s. on $\{\langle M \rangle_\infty = \infty\}$. But (K_n) is a square-integrable martingale with

$$\langle K \rangle_n = \sum_{k=1}^n \frac{\mathbf{E}[(\Delta M_k)^2 | \mathcal{F}_{k-1}]}{(1 + \langle M \rangle_k)^2} \leq \sum_{k=1}^n (\Delta M_k)^2 = \langle M \rangle_n.$$

By part (a) of Theorem 2, $\{\langle K \rangle_\infty < \infty\} \subset \{K_n \rightarrow\}$ (a.s.). To conclude, we must verify that $\langle K \rangle_\infty < \infty$. To this end notice that

$$\langle K \rangle_\infty = \sum_{k=1}^{\infty} \frac{\Delta\beta_k}{\beta_k^2},$$

where $\beta_k := 1 + \langle M \rangle_k > 0$, so that $\Delta\beta_k \geq 0$. Because

$$-\Delta\left(\frac{1}{\beta_k}\right) = -\left(\frac{1}{\beta_k} - \frac{1}{\beta_{k-1}}\right) = \frac{\Delta\beta_k}{\beta_k\beta_{k-1}} \geq \frac{\Delta\beta_k}{\beta_k^2},$$

we have

$$\sum_{k=1}^n \frac{\Delta\beta_k}{\beta_k^2} \leq -\sum_{k=1}^n \Delta(1/\beta_k) = \beta_0^{-1} - \beta_n^{-1} = 1 - \beta_n^{-1} \leq 1.$$

It follows that $\langle K \rangle_\infty \leq 1$ a.s., and we are done. \square

The SLLN in hand, it is quite natural to look for martingale results analogous to the CLT. The following basic martingale CLT is due to B.M. Brown [*Annals of Math. Stat.* **42** (1971) 59–66], to which you are referred for the proof.

4. Theorem. *Let $M = (M_n)$ be a square-integrable martingale with quadratic variation $\langle M \rangle$ as defined in (1). Define $s_n^2 := \mathbf{E}[\langle M \rangle_n]$. Assume that*

$$\frac{\langle M \rangle_n}{s_n^2} \xrightarrow{P} 1$$

and that the Lindeberg-type condition

$$s_n^{-2} \sum_{k=1}^n \mathbf{E}[(\Delta M_k)^2; (\Delta M_k)^2 \geq \epsilon \cdot s_n] \xrightarrow{P} 0, \quad \forall \epsilon > 0,$$

is satisfied. Then $M_n/s_n \Rightarrow \mathcal{N}(0, 1)$.