

Note on a paper of Chen & Tweedie

In a recent paper [1], P.-D. CHEN and R.L. TWEEDIE make use of the natural embedding of a separable and separated measurable space in the compact cube $\{0, 1\}^{\mathbb{N}}$. The point of this note is to show that a well-known representation of functions measurable with respect to the σ -algebra generated by a sequence of functions can be substituted for the rather intricate arguments found in [1].

Let (X, \mathcal{X}) be a measurable space such that \mathcal{X} is countably generated and separated. The latter term means that if x and y are distinct points of X then there exists $A \in \mathcal{X}$ such that $x \in A$ and $y \in A^c$.

Let I denote the cube $\{0, 1\}^{\mathbb{N}}$ endowed with its natural (compact, metrizable) topology, and let \mathcal{I} denote the class of Borel subsets of I . Let $\{A_1, A_2, \dots\}$ be a sequence of elements of \mathcal{X} such that $\sigma\{A_n : n \in \mathbb{N}\} = \mathcal{X}$. Define a mapping $T : X \rightarrow I$ by the formula $T(x) = (1_{A_1}(x), 1_{A_2}(x), \dots)$. Clearly T is injective and \mathcal{X}/\mathcal{I} -measurable. If we endow the image $Y := T(X)$ with the trace σ -algebra $\mathcal{Y} := \mathcal{I} \cap Y$, then T is an isomorphism of the measurable spaces (X, \mathcal{X}) and (Y, \mathcal{Y}) . In general, $Y \notin \mathcal{I}$; indeed, $Y \in \mathcal{I}$ if and only if (X, \mathcal{X}) is a measurable Lusin space (*i.e.*, isomorphic to a Borel subspace of some compact metrizable space).

Put $Z := \bar{Y}$. Then Z is compact, and the trace σ -algebra $\mathcal{Z} := \mathcal{I} \cap Z$ coincides with the class of Borel subsets of Z . Of course, $\mathcal{Y} = \mathcal{Z} \cap Y$.

Lemma. *Let (F, \mathcal{F}) be a Lusin space and let $f : X \rightarrow F$ be \mathcal{X}/\mathcal{F} -measurable. Then there exists an \mathcal{Z}/\mathcal{F} -measurable function $\hat{f} : Z \rightarrow F$ such that $f = \hat{f} \circ T$.*

Proof. The assertion is trivial if F is countable. If F is uncountable, then it is Borel isomorphic to the unit interval $[0, 1]$. Since both the hypotheses and conclusions are isomorphism invariant we can assume without loss of generality that $F = [0, 1]$. But in this case the assertion is a standard consequence of the monotone class theorem. \square

If P is a probability measure on (X, \mathcal{X}) , let $Q := T_*P$ denote the image of P under T , viewed as a measure on (Z, \mathcal{Z}) :

$$Q(H) := P(T^{-1}H), \quad \forall H \in \mathcal{Z}.$$

The class of images $\mathcal{Q} := \{T_*P : P \in \mathcal{P}(E)\}$ does not exhaust $\mathcal{P}(Z)$.* In fact, $Q \in \mathcal{P}(Z)$ lies in \mathcal{Q} if and only if the Q outer measure of Y is 1.

Say that $A \in \mathcal{Z}$ is Q -negligible provided $Q(A) = 0$ for all $Q \in \mathcal{Q}$. Likewise, \mathcal{Z} -measurable functions f and g are Q -equivalent ($f \sim g$) provided $\{f \neq g\}$ is Q -negligible. Clearly $f \sim g$ if and only if $\{f \neq g\} \subset Z \setminus Y$. In particular, the function \hat{f} provided by the Lemma is uniquely determined modulo Q -equivalence.

* $\mathcal{P}(Z)$ denotes the class of probability measures on (Z, \mathcal{Z}) , etc.

Let us now see how the Lemma provides an easy way to “lift” a Markov kernel on (X, \mathcal{X}) to a Markov kernel on (Z, \mathcal{Z}) . By the term “Markov kernel” (on (X, \mathcal{X}) , say) we mean a mapping $(x, A) \mapsto K(x, A)$ from $X \times \mathcal{X}$ to $[0, 1]$ such that $x \mapsto K(x, A)$ is \mathcal{X} -measurable for each $A \in \mathcal{A}$ and $A \mapsto K(x, A)$ is an element of $\mathcal{P}(X)$ for all $x \in X$. Here is the main result of section 3 of [1]:

Theorem. *Let K be a Markov kernel on (X, \mathcal{X}) . Then there exists a Markov kernel \hat{K} on (Z, \mathcal{Z}) such that $K(x, T^{-1}H) = \hat{K}(T(x), H)$ for all $x \in X$ and all $H \in \mathcal{Z}$. If \bar{K} is another such “lifting” of K , then $\{z \in Z : \bar{K}(z, \cdot) \neq \hat{K}(z, \cdot)\}$ is \mathcal{Q} -negligible.*

Proof. For each fixed $x \in X$, the mapping $\mathcal{Z} \ni H \mapsto K(x, T^{-1}H)$ defines an element $K_*(x, \cdot)$ of $\mathcal{P}(Z)$ (in fact, an element of \mathcal{Q}). Since Z is compact metrizable, so is $\mathcal{P}(Z)$. By the Lemma (applied to the function $f : x \mapsto K_*(x, \cdot)$, which is a measurable mapping of X into $\mathcal{P}(Z)$) there is a measurable function $\hat{f} : Z \rightarrow \mathcal{P}(Z)$ such that $f = \hat{f} \circ T$. Evidently, \hat{f} determines a Markov kernel \hat{K} on (Z, \mathcal{Z}) by the formula $\hat{K}(z, H) = [\hat{f}(z)](H)$. In particular, $\hat{K}(T(x), H) = [\hat{f}(T(x))](H) = [f(x)](H) = K(x, T^{-1}H)$, for all $H \in \mathcal{Z}$ and $x \in X$. The uniqueness assertion is obvious. \square

References

- [1] P.-D. CHEN and R.L. TWEEDIE: Orthogonal measures and absorbing sets for Markov chains, *Math. Proc. Camb. Phil. Soc.* **121** (1997) 101–113).