

**SIXTY-FIFTH ANNUAL
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION**

Saturday, December 4, 2004

Examination A

A1. Basketball star Shanille O'Keal's team statistician keeps track of the number, $S(N)$, of successful free throws she has made in her first N attempts of the season. Early in the season, $S(N)$ was less than 80% of N , but by the end of the season, $S(N)$ was more than 80% of N . Was there necessarily a moment in between when $S(N)$ was exactly 80% of N ?

A2. For $i = 1, 2$, let T_i be a triangle with side lengths a_i, b_i, c_i , and area A_i . Suppose that $a_1 \leq a_2, b_1 \leq b_2, c_1 \leq c_2$, and that T_2 is an acute triangle. Does it follow that $A_1 \leq A_2$?

A3. Define a sequence $\{u_n\}_{n=0}^{\infty}$ by $u_0 = u_1 = u_2 = 1$, and thereafter by the condition that

$$\det \begin{bmatrix} u_n & u_{n+1} \\ u_{n+2} & u_{n+3} \end{bmatrix} = n!$$

for all $n \geq 0$. Show that u_n is an integer for all n . (By convention, $0! = 1$.)

A4. Show that for any positive integer n there is an integer N such that the product $x_1 x_2 \cdots x_n$ can be expressed identically in the form

$$x_1 x_2 \cdots x_n = \sum_{i=1}^N c_i (a_{i1} x_1 + a_{i2} x_2 + \cdots + a_{in} x_n)^n$$

where the c_i are rational numbers and each a_{ij} is one of the numbers, $-1, 0, 1$.

A5. An $m \times n$ checkerboard is colored randomly: each square is independently assigned red or black with probability $1/2$. We say that two squares, p and q , are in the same connected monochromatic region if there is a sequence of squares, all of the same color, starting at p and ending at q , in which successive squares in the sequence share a common side. Show that the expected number of connected monochromatic regions is greater than $mn/8$.

A6. Suppose that $f(x, y)$ is a continuous real-valued function on the unit square $0 \leq x \leq 1, 0 \leq y \leq 1$. Show that

$$\begin{aligned} \int_0^1 \left(\int_0^1 f(x, y) dx \right)^2 dy + \int_0^1 \left(\int_0^1 f(x, y) dy \right)^2 dx \\ \leq \left(\int_0^1 \int_0^1 f(x, y) dx dy \right)^2 + \int_0^1 \int_0^1 [f(x, y)]^2 dx dy. \end{aligned}$$

**SIXTY-FIFTH ANNUAL
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION**

Saturday, December 4, 2004

Examination B

B1. Let $P(x) = c_n x^n + c_{n-1} x^{n-1} + \cdots + c_0$ be a polynomial with integer coefficients. Suppose that r is a rational number such that $P(r) = 0$. Show that the n numbers

$$c_n r, c_n r^2 + c_{n-1} r, c_n r^3 + c_{n-1} r^2 + c_{n-2} r, \dots, c_n r^n + c_{n-1} r^{n-1} + \cdots + c_1 r$$

are integers.

B2. Let m and n be positive integers. Show that

$$\frac{(m+n)!}{(m+n)^{m+n}} < \frac{m!}{m^m} \cdot \frac{n!}{n^n}.$$

B3. Determine all real numbers $a > 0$ for which there exists a nonnegative continuous function $f(x)$ defined on $[0, a]$ with the property that the region

$$R = \{(x, y) : 0 \leq x \leq a, 0 \leq y \leq f(x)\}$$

has perimeter k units and area k square units for some real number k .

B4. Let n be a positive integer, $n \geq 2$, and put $\theta = 2\pi/n$. Define points $P_k = (k, 0)$ in the xy -plane, for $k = 1, 2, \dots, n$. Let R_k be the map that rotates the plane counterclockwise by the angle θ about the point P_k . Let R denote the map obtained by applying, in order, R_1 , then R_2 , \dots , then R_n . For an arbitrary point (x, y) , find, and simplify, the coordinates of $R(x, y)$.

B5. Evaluate

$$\lim_{x \rightarrow 1^-} \prod_{n=0}^{\infty} \left(\frac{1+x^{n+1}}{1+x^n} \right)^{x^n}.$$

B6. Let \mathcal{A} be a non-empty set of positive integers, and let $N(x)$ denote the number of elements of \mathcal{A} not exceeding x . Let \mathcal{B} denote the set of positive integers b that can be written in the form $b = a - a'$ with $a \in \mathcal{A}$ and $a' \in \mathcal{A}$. Let $b_1 < b_2 < \cdots$ be the members of \mathcal{B} , listed in increasing order. Show that if the sequence $b_{i+1} - b_i$ is unbounded, then $\lim_{x \rightarrow \infty} N(x)/x = 0$.