

**SEVENTIETH ANNUAL  
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION**

Saturday, December 5, 2009

Examination A

**A1.** Let  $f$  be a real-valued function on the plane such that for every square  $ABCD$  in the plane,  $f(A) + f(B) + f(C) + f(D) = 0$ . Does it follow that  $f(P) = 0$  for all points  $P$  in the plane?

**A2.** Functions  $f, g, h$  are differentiable on some open interval around 0 and satisfy the equations and initial conditions

$$\begin{aligned}f' &= 2f^2gh + \frac{1}{gh}, & f(0) &= 1, \\g' &= fg^2h + \frac{4}{fh}, & g(0) &= 1, \\h' &= 3fgh^2 + \frac{1}{fg}, & h(0) &= 1.\end{aligned}$$

Find an explicit formula for  $f(x)$ , valid in some open interval around 0.

**A3.** Let  $d_n$  be the determinant of the  $n \times n$  matrix whose entries, from left to right and then from top to bottom, are  $\cos 1, \cos 2, \dots, \cos n^2$ . (For example,

$$d_3 = \begin{vmatrix} \cos 1 & \cos 2 & \cos 3 \\ \cos 4 & \cos 5 & \cos 6 \\ \cos 7 & \cos 8 & \cos 9 \end{vmatrix}.$$

The argument of  $\cos$  is always in radians, not degrees.) Evaluate  $\lim_{n \rightarrow \infty} d_n$ .

**A4.** Let  $S$  be a set of rational numbers such that

- (a)  $0 \in S$ ;
- (b) If  $x \in S$  then  $x + 1 \in S$  and  $x - 1 \in S$ ; and
- (c) If  $x \in S$  and  $x \notin \{0, 1\}$ , then  $1/(x(x - 1)) \in S$ .

Must  $S$  contain all rational numbers?

**A5.** Is there a finite abelian group  $G$  such that the product of the orders of all its elements is  $2^{2009}$ ?

**A6.** Let  $f : [0, 1]^2 \rightarrow \mathbf{R}$  be a continuous function on the closed unit square such that  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  exist and are continuous on the interior  $(0, 1)^2$ . Let  $a = \int_0^1 f(0, y) dy$ ,  $b = \int_0^1 f(1, y) dy$ ,  $c = \int_0^1 f(x, 0) dx$ ,  $d = \int_0^1 f(x, 1) dx$ . Prove or disprove: There must be a point  $(x_0, y_0)$  in  $(0, 1)^2$  such that

$$\frac{\partial f}{\partial x}(x_0, y_0) = b - a \quad \text{and} \quad \frac{\partial f}{\partial y}(x_0, y_0) = d - c.$$

**SEVENTIETH ANNUAL  
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION**

Saturday, December 5, 2009

Examination B

---

**B1.** Show that every positive rational number can be written as a quotient of products of factorials of (not necessarily distinct) primes. For example,

$$\frac{10}{9} = \frac{2! \cdot 5!}{3! \cdot 3! \cdot 3!}.$$

**B2.** A game involves jumping to the right on the real number line. If  $a$  and  $b$  are real numbers and  $b > a$ , the cost of jumping from  $a$  to  $b$  is  $b^3 - ab^2$ . For what real numbers  $c$  can one travel from 0 to 1 in a finite number of jumps with total cost exactly  $c$ ?

**B3.** Call a subset  $S$  of  $\{1, 2, \dots, n\}$  *mediocre* if it has the following property: Whenever  $a$  and  $b$  are elements of  $S$  whose average is an integer, that average is also an element of  $S$ . Let  $A(n)$  be the number of mediocre subsets of  $\{1, 2, \dots, n\}$ . [For instance, every subset of  $\{1, 2, 3\}$  except  $\{1, 3\}$  is mediocre, so  $A(3) = 7$ .] Find all positive integers  $n$  such that  $A(n+2) - 2A(n+1) + A(n) = 1$ .

**B4.** Say that a polynomial with real coefficients in two variables,  $x, y$ , is *balanced* if the average value of the polynomial on each circle centered at the origin is 0. The balanced polynomials of degree at most 2009 form a vector space  $V$  over  $\mathbf{R}$ . Find the dimension of  $V$ .

**B5.** Let  $f : (1, \infty) \rightarrow \mathbf{R}$  be a differentiable function such that

$$f'(x) = \frac{x^2 - (f(x))^2}{x^2((f(x))^2 + 1)} \quad \text{for all } x > 1.$$

Prove that  $\lim_{x \rightarrow \infty} f(x) = \infty$ .

**B6.** Prove that for every positive integer  $n$ , there is a sequence of integers  $a_0, a_1, \dots, a_{2009}$  with  $a_0 = 0$  and  $a_{2009} = n$  such that each term after  $a_0$  is either an earlier term plus  $2^k$  for some nonnegative integer  $k$ , or of the form  $b \bmod c$  for some earlier positive terms  $b$  and  $c$ . [Here  $b \bmod c$  denotes the remainder when  $b$  is divided by  $c$ , so  $0 \leq (b \bmod c) < c$ .]