

ON GRAPHS WHICH CONTAIN ALL SMALL TREES, II.

F.R.K. CHUNG — R.L. GRAHAM — N. PIPPENGER

INTRODUCTION

Let  $\mathcal{T}_n$  denote the class of all trees\* with  $n$  edges and denote by  $s(\mathcal{T}_n)$  the minimum number of edges a graph  $G$  can have which contains all  $T \in \mathcal{T}_n$  as subgraphs. In a previous paper [2], two of the authors established the following bounds on  $s(\mathcal{T}_n)$ :

$$(1) \quad \frac{1}{2} n \log n < s(\mathcal{T}_n) < n^{1 + \frac{1}{\log \log n}}$$

where  $n$  is taken sufficiently large. In this note, we strengthen the upper bound on  $s(\mathcal{T}_n)$  considerably. In addition we also consider the same problem in the case that  $G$  is restricted to be a tree, with  $s_{\mathcal{T}}(\mathcal{T}_n)$  denoting the corresponding minimum number of edges. Surprisingly, we show that  $s_{\mathcal{T}}(\mathcal{T}_n)$  does *not* grow exponentially in  $n$ , answering a question in [2]. It is annoying, however, that at present we cannot even show that  $s_{\mathcal{T}}(\mathcal{T}_n)$  must exceed  $n^{2+\epsilon}$  for large  $n$ .

\*The reader may consult [1] or [3] for any undefined graph-theoretic terminology.

## W-SUBTREES OF A TREE

Before establishing new bounds on  $s(\mathcal{T}_n)$  and  $s_{\mathcal{T}}(\mathcal{T}_n)$ , we first require a result concerning the decomposition of trees.

Let  $W$  be a nonempty set of vertices of a tree  $T$ . By a  $W$ -subtree of  $T$ , we mean a subtree  $T'$  of  $T$  consisting of one of the components  $C$  formed from  $T$  by the removal of all the vertices of  $W$ , except for those vertices of  $W$  adjacent to some vertex of  $C$  (and the edges joining them).

**Example.**




Fig. 1

As usual, we let  $\|G\|$  denote the number of edges of a graph  $G$ .

**Lemma.** *Let  $w$  be a nonnegative integer. Then if  $\alpha$  is sufficiently large, any tree  $T$  with at least  $\alpha + 1$  edges has a subset of vertices  $W$  with  $|W| \leq w + 1$  so that for some set  $\mathcal{C}$  of  $W$ -subtrees of  $T$  we have*

$$(2) \quad \alpha < \sum_{T' \in \mathcal{C}} \|T'\| \leq \left(1 + \left(\frac{2}{3}\right)^w\right) \alpha.$$

**Proof.** For  $w = 0$ , this is a result in [2]. Assume  $w = 1$ . We know that if  $\alpha$  is large enough then for some vertex  $u$  there is a set  $\mathcal{C}(u)$  of  $\{u\}$ -subtrees of  $T$  such that

$$(3) \quad \alpha < \sum_{T' \in \mathcal{C}(u)} \|T'\| \leq 2\alpha.$$

If

$$\sum_{T' \in \mathcal{C}(u)} \|T'\| \leq \frac{5}{3}\alpha,$$

then the lemma holds for  $w = 1$ . Hence, we may assume

$$\frac{5}{3}\alpha < \sum_{T' \in \mathcal{C}(u)} \|T'\| \leq 2\alpha.$$

Let  $T_1$  be the subtree of  $T$  formed by taking the union of all  $T' \in \mathcal{C}(u)$ . Again, for  $\alpha$  sufficiently large, there exists a vertex  $v$  of  $T_1$  so that for some set  $\mathcal{C}(v)$  of  $\{v\}$ -subtrees of  $T_1$ , we have

$$\frac{\alpha}{3} \sum_{T'' \in \mathcal{C}(v)} \|T''\| \leq \frac{2\alpha}{3}.$$

Consider the set  $\mathcal{C}'(v)$  all of  $\{v\}$ -subtrees of  $T_1$  which are *not* in  $\mathcal{C}(v)$ . Then

$$\alpha < \sum_{T' \in \mathcal{C}'(v)} \|T'\| \leq \frac{5}{3}\alpha.$$

However, a  $\{v\}$ -subtree of  $T_1$  is a  $\{u, v\}$ -subtree of  $T$ . This proves the lemma for the case  $w = 1$ . The inductive proof of (2) for general  $w$  follows very similar lines and will not be given. ■

## AN UPPER BOUND ON $s(\mathcal{F}_n)$

### Theorem 1.

$$s(\mathcal{F}_n) = O(n \log n (\log \log n)^2).$$

**Proof.** For  $p \geq 0$ , let us define the graph  $G_{w,p}$  as follows.  $G_{w,0} = K_{w+1}$ , the complete graph on  $w+1$  vertices. For  $p > 0$ ,  $G_{w,p}$  will denote the graph formed from  $K_{w+1}$  and two disjoint copies of  $G_{w,p-1}$ , by placing an edge between each vertex of  $K_{w+1}$  and each vertex of each of the copies of  $G_{w,p-1}$  (see Figure 2).




Fig. 2

Simple inductive arguments show that  $|G_{w,p}| = O(w2^p)$  and  $\|G_{w,p}\| = O(w^2 p 2^p)$  (where  $|G|$  denotes the number of vertices in  $G$ ). It is also not difficult to see that  $G_{w,p}$  contains all trees with at most

$$\left( \frac{2 + \left(\frac{2}{3}\right)^w}{1 + \left(\frac{2}{3}\right)^w} \right)^p$$

edges. For  $p = 1$ , the expression is less than 2 and the claim is trivial. For  $p > 1$ , application of the preceding Lemma with

$$\alpha = \frac{1}{1 + \left(\frac{2}{3}\right)^w} \left( \frac{2 + \left(\frac{2}{3}\right)^w}{1 + \left(\frac{2}{3}\right)^w} \right)^{p-1},$$

guarantees a set  $W$  of  $w + 1$  vertices (which may be assigned to the vertices of  $K_{w+1}$  in  $G_{w,p}$ ) and a decomposition of the  $W$ -subtrees into two classes, each having at most

$$\left( \frac{2 + \left(\frac{2}{3}\right)^w}{1 + \left(\frac{2}{3}\right)^w} \right)^{p-1}$$

edges (which may be assigned to the two copies of  $G_{w,p-1}$  in  $G_{w,p}$ ).

If we now choose  $q = \left\lceil \frac{\log 2n}{\log 2} \right\rceil$  and  $w = \left\lceil \frac{\log q}{\log \frac{3}{2}} \right\rceil$  we find that

$$\|G_{w,q}\| = O(n \log n (\log \log n)^2).$$

Furthermore, a simple calculation shows that

$$\left( \frac{2 + \left(\frac{2}{3}\right)^w}{1 + \left(\frac{2}{3}\right)^w} \right)^q \geq 2^q \left( 1 - \frac{1}{2} \left(\frac{2}{3}\right)^w \right)^q \geq 2^{q-1} \geq n,$$

so that  $G_{w,q}$  contains as subgraphs all trees with at most  $n$  edges. ■

### TREES CONTAINING ALL SMALL TREES

We next turn our attention to the case in which  $G$  is restricted to be a tree. As mentioned in the introduction, it was asked in [2] whether or not  $s_{\mathcal{T}}(\mathcal{T}_n)$ , the corresponding minimum number of edges in this case, must grow exponentially in  $n$ . This is settled by Theorem 2.

Before presenting this result, we first list the values of  $s(\mathcal{T}_n)$  for  $n \leq 7$ . We also show trees which produce these values (see Fig. 3). The corresponding proofs for these results are straightforward (using degree sequence considerations) and are omitted.

$n$	$s_{\mathcal{T}}(\mathcal{T}_n)$
1	1
2	2
3	4
4	6
5	9
6	13
7	17

Table 1




Fig. 3

**Theorem 2.**

$$s_{\mathcal{F}}(\mathcal{F}_n) \leq \frac{2\sqrt{2}}{n} \exp \frac{\log^2 n}{2 \log 2}$$

for  $n$  sufficiently large.

**Proof.** Let us consider a family of rooted trees  $\bar{G}(x)$  with a root at some vertex of degree 1 which contains as subgraphs all rooted trees on at most  $x$  edges which have a root at some vertex of degree 1. For  $1 \leq k < n$ , let  $\bar{G}(\frac{n-1}{k})$  have as its root  $r_k$ . Form the graph  $\bar{G}(n)$  (as shown in Fig. 4) by identifying all the  $r_k$  as a single vertex  $r^*$  and adjoining a root  $r_n$  of degree 1 to  $r^*$ . We note that  $\bar{G}(x) = \bar{G}(n)$  where  $n$  is the integral part of  $x$ .




Fig. 4

It is easy to see that if  $\bar{f}$  satisfies

$$(4) \quad \bar{f}(x) \geq \sum_{k=1}^{\lfloor x \rfloor} \bar{f}\left(\frac{x-1}{k}\right),$$

for sufficiently large  $x$  then

$$(5) \quad \|\bar{G}(n)\| \leq \bar{f}(n).$$

We claim that it will suffice to have  $\bar{f}$  satisfy

$$(6) \quad \bar{f}(x) \geq \bar{f}(x-1) + 2\bar{f}\left(\frac{x+1}{2}\right)$$

in order for (4) to hold. For (6) implies

$$\begin{aligned} \bar{f}(x) &\geq \bar{f}(x-1) + 2\bar{f}\left(\frac{x-1}{2}\right) \geq \\ &\geq \bar{f}(x-1) + 2\bar{f}\left(\frac{x-1}{2}\right) + 4\bar{f}\left(\frac{x-3}{4}\right) \geq \\ &\geq \bar{f}(x-1) + 2\bar{f}\left(\frac{x-1}{2}\right) + 4\bar{f}\left(\frac{x-1}{4}\right) + 8\bar{f}\left(\frac{x+7}{8}\right) \geq \\ &\quad \vdots \\ &\geq \sum_{2^k < x} 2^{k-1} \bar{f}\left(\frac{x-1}{2^k}\right) \geq \sum_{k=1}^{\lfloor x \rfloor} \bar{f}\left(\frac{x-1}{k}\right). \end{aligned}$$

A straightforward computation now shows that the choice

$$\bar{f}(x) = e^{\frac{\log^2 x}{2 \log 2}}$$

satisfies (6) for  $x$  sufficiently large.

Let  $G(x)$  be a graph as shown in Figure 5.

It is immediate that  $G(x)$  contains all  $T \in \mathcal{T}_n$  as subgraphs and we have

$$s_{\mathcal{T}}(\mathcal{T}_n) \leq \|G(x)\| \leq \frac{2\sqrt{2}}{n} \cdot \exp\left(\frac{(\log n)^2}{2 \log 2}\right).$$

This proves the theorem. ■

Let  $s_{\mathcal{T}}^*(\mathcal{T}_n)$  be the minimum number of edges a *rooted* tree can have which contains all *rooted* trees of  $n$  edges as subgraphs. Of course, the inequality

$$s_{\mathcal{T}}(\mathcal{T}_n) \leq s_{\mathcal{T}}^*(\mathcal{T}_n)$$

is immediate. In fact, we now show that if  $s_{\mathcal{T}}(\mathcal{T}_n)$  grows polynomially in  $n$ , then so does  $s_{\mathcal{T}}^*(\mathcal{T}_n)$ .






Fig. 5

**Theorem 3.**

$$s_{\mathcal{T}_n} \leq s_{\mathcal{T}_n}^* \leq s_{\mathcal{T}_n} \cdot (s_{\mathcal{T}_n} + 1).$$

**Proof.** Let  $G_n$  be a tree with  $s_{\mathcal{T}_n}$  edges which contains all  $T \in \mathcal{T}_n$  as subgraphs. Let  $G_n(v)$ ,  $v \in G_n$ , be a rooted tree which has the same structure as  $G_n$  and which has  $v$  as its root. Now, form the rooted tree  $H_n$  (as shown in Fig. 6) by identifying all the roots  $v$  in  $G_n(v)$  for  $v \in G_n$ .

It is easily verified that  $H_n$  contains all rooted trees with  $n$  edges and satisfies

$$s_{\mathcal{T}_n}^* \leq \|H_n\| \leq s_{\mathcal{T}_n} (s_{\mathcal{T}_n} + 1).$$

This proves the theorem. ■




Fig. 6

### CONCLUDING REMARKS

As remarked earlier, the best known lower bound for  $s(\mathcal{T}_n)$  is  $\frac{1}{2} n \log n$  which is not too far from the upper bound of  $O(n \log n (\log \log n)^2)$  of Theorem 1. Perhaps the lower bound is the correct order of magnitude. Unfortunately, the only lower bound presently known for  $s_{\mathcal{F}}(\mathcal{T}_n)$  is rather weak. By considering the possible locations of the vertices of degree 1 of the  $T \in \mathcal{T}_n$ , it can be argued that

$$s_{\mathcal{F}}(\mathcal{T}_n) > cn^2$$

for some  $c > 0$ . It seems likely that

$$\frac{s_{\mathcal{F}}(\mathcal{T}_n)}{n^k} \rightarrow \infty$$

for any fixed  $k$ .

## REFERENCES

- [1] C. Berge, *Graphs and hypergraphs*, North-Holland, Amsterdam, 1973.
- [2] F.R.K. Chung – R.L. Graham, On graphs which contain all small trees (to appear in *Jour. Comb. Th.*, (B)).
- [3] F. Harary, *Graph theory*, Addison-Wesley, New York, 1969.

F.R.K. Chung – R.L. Graham

Bell Laboratories, Murray Hill, New Jersey.

N. Pippenger

IBM Thomas J. Watson Research Center, Yorktown Heights, New York.