

## Note

# On Subsets of Abelian Groups with No 3-Term Arithmetic Progression

P. FRANKL\*

*C.N.R.S., Paris, France*

R. L. GRAHAM

*AT&T Bell Laboratories  
600 Mountain Avenue, Murray Hill, New Jersey 07974*

AND

V. RÖDL\*

*FJFI CVUT, Prague, Czechoslovakia*

*Communicated by the Managing Editors*

Received October 25, 1986

A short proof of the following result of Brown and Buhler is given: For any  $\varepsilon > 0$  there exists  $n_0 = n_0(\varepsilon)$  such that if  $A$  is an abelian group of odd order  $|A| > n_0$  and  $B \subseteq A$  with  $|B| > \varepsilon|A|$ , then  $B$  must contain three distinct elements  $x, y, z$  satisfying  $x + y = 2z$ . © 1987 Academic Press, Inc.

### 1. INTRODUCTION

Let  $N$  denote the set of positive integers, and for  $n \in N$ , let  $[n]$  denote the set  $\{1, 2, \dots, n\}$ . A well-known theorem of Roth [R] asserts that if  $P \subseteq N$  contains no 3-term arithmetic progression, then  $P$  has upper density zero. That is, for every  $\varepsilon > 0$ ,  $|P \cap [n]| < \varepsilon n$  holds for all sufficiently large  $n$ .

Brown and Buhler [BB1] proved the following generalization of Roth's result.

\* This work was performed while the authors were visiting AT&T Bell Laboratories.

**THEOREM 1.** *For every  $\varepsilon > 0$  there exists  $n_0 = n_0(\varepsilon)$  with the following property. Suppose  $A$  is an abelian group of odd order,  $|A| > n_0$ . Then every subset  $B \subset A$  with  $|B| > \varepsilon|A|$  contains three distinct elements  $x, y, z$  with  $x + y = 2z$ .*

For a finite set  $X$ , define  $\binom{X}{k} := \{F \subseteq X : |F| = k\}$ . A family  $\mathbf{F} \subseteq \binom{X}{k}$  is called a  $k$ -graph. It is called *linear* if  $|F \cap G| \leq 1$  holds for all distinct  $F, G \in \mathbf{F}$ . Three distinct edges,  $F, G, H$  of a linear  $k$ -graph are said to form a *triangle* if the three intersections  $F \cap G, G \cap H, H \cap F$  are all non-empty and distinct.

**THEOREM 2 (Ruzsa–Szemerédi [RS]).** *Suppose that  $\mathbf{F}$  is a linear 3-graph on  $n$  vertices which contains no triangle. Then  $|\mathbf{F}| = o(n^2)$ .*

For a simple proof of Theorem 2, see [EFR]. Here we show that Theorem 1 follows easily from Theorem 2.

## 2. PROOF OF THEOREM 1

Suppose  $A$  is an abelian group of odd order and  $B \subseteq A$  contains no three distinct elements  $x, y, z$  with  $x + y = 2z$ . Define  $X = A \times [3]$  to be the  $3|A|$ -element set with general element  $(a, i)$ ,  $a \in A$ ,  $1 \leq i \leq 3$ . Now define a 3-graph  $\mathbf{F}$  as

$$\mathbf{F} := \{ \{ (a, 1), (a + b, 2), (a + 2b, 3) \} : a \in A, b \in B \}.$$

Clearly,  $|\mathbf{F}| = |A| |B|$ . Also,  $\mathbf{F}$  is linear since any two elements of an edge uniquely determine the edge.

Suppose now to the contrary that  $\mathbf{F}$  contains a triangle, say

$$\{ (a_i, 1), (a_i + b_i, 2), (a_i + 2b_i, 3) \}, \quad i = 1, 2, 3.$$

By symmetry, we may assume that

$$a_1 = a_2, \quad a_1 + b_1 = a_3 + b_3, \quad a_2 + 2b_2 = a_3 + 2b_3.$$

However, these equations imply

$$2b_2 - 2b_3 = a_3 - a_2 = a_3 - a_1 = b_1 - b_3,$$

i.e.,

$$2b_2 = b_1 + b_3.$$

By the choice of  $B$ , this implies  $b_1 = b_2 = b_3$  and thus,  $a_1 = a_2 = a_3$ , a contradiction. Thus,  $\mathbf{F}$  contains no triangle.

Hence, by Theorem 2,

$$|\mathbf{F}| = |A| |B| = o(|A|^2),$$

i.e.,

$$|B| = o(|A|) \quad \text{as desired.}$$

*Remark.* The same proof can be used in the case when  $A$  is a  $d$ -dimensional affine space over  $\text{GF}(2^t)$ ,  $t \geq 2$ . For the definition of edges in the proof, one replaces  $a + 2b$  by  $a + \gamma b$  where  $\gamma \neq 0$ , 1 is an arbitrary element of  $\text{GF}(2^t)$ . The conclusion then becomes:  $B$  contains three points on a line.

### 3. SOME LOWER BOUNDS

The most important special cases of Theorem 1 are when  $A$  is a cyclic group (corresponding to Roth's theorem) and when  $A$  is an affine space  $A(d, q)$  of dimension  $d$  over  $\text{GF}(q)$ .

In both cases, stronger theorems are known. Szemerédi's theorem [S] asserts that sets with positive upper density contain arithmetic progressions of arbitrary length, while a recent result of Furstenberg and Katznelson [FK] implies that for any  $\varepsilon > 0$  and any prime power  $q$  there exists  $d_0 = d_0(\varepsilon, q)$  so that the following is true: Every subset  $B \subseteq A(d, q)$  with  $|B| > \varepsilon q^d$ ,  $d > d_0$ , contains all the points of some line in  $A(d, q)$ .

In view of [BB2] this implies the same statement if we replace lines by planes, spaces, etc.

Let  $a_q(d)$  denote the maximum of  $|B|$  where  $B \subseteq A(d, q)$  contains no line. In the case of the integers, Behrend [B] showed that for every  $\delta > 0$  and  $n > n_0(\delta)$  there exists  $B \subseteq [n]$  with  $|B| > n^{1-\delta}$  so that  $B$  contains no 3-term arithmetic progression. We do not know if the corresponding statement holds for affine spaces.

**PROBLEM.** Is it true that for every  $\delta > 0$ ,  $q \geq 3$  and  $d > d_0(\delta, q)$ , there exists  $B \subseteq A(d, q)$  which contains no line and satisfies  $|B| > (q - \delta)^d$ .

It is easy to construct such a  $B$  with  $|B| = (q - 1)^d$ ; simply take

$$B = \{(b_1, \dots, b_d) : b_i \in \text{GF}(q) \setminus \{0\}, i = 1, \dots, d\}.$$

To improve on this bound note that if  $x \in \text{GF}(q)$  is fixed and  $\{(b_1^{(i)}, \dots, b_d^{(i)}), i = 1, \dots, q\}$  forms a line then the sets  $F_i = \{j : b_j^{(i)} = x\}$  form a *sunflower* of size  $q$ , that is,  $F_1 \cap \dots \cap F_q = F_j \cap F_{j'}$  holds for all  $1 \leq j < j' \leq q$ .

Let  $f_q(d, r)$  denote the maximum of  $|\mathbf{F}|$  where  $\mathbf{F} \subseteq \binom{[d]}{r}$ , and  $\mathbf{F}$  contains no sunflower of size  $q$ .

PROPOSITION. For all positive integers  $n, r, d$ , one has

$$a_q(n) \geq f_q(n, r)(q-1)^{n-r} \quad \text{and} \quad a_q(nd) \geq a_q(n)^d. \quad (2)$$

*Proof.* Let  $\mathbf{F} \subseteq \binom{[n]}{r}$  be a family without sunflowers of size  $q$  which satisfies  $|\mathbf{F}| = f_q(n, r)$ .

Fix an element  $x \in GF(q)$ . For  $\mathbf{b} = (b_1, \dots, b_n) \in A(n, q)$ , define  $F(\mathbf{b}) := \{j: b_j = x\}$  and

$$B := \{\mathbf{b} \in A(n, q): F(\mathbf{b}) \in \mathbf{F}\}.$$

Then  $B$  contains no line. To prove the second assertion, one simply notes that if  $B$  contains no line then

$$B \oplus \dots \oplus B \subseteq A(n, q) \oplus \dots \oplus A(n, q) = A(dn, q)$$

contains no line either.

If we knew the value of  $f_q(n, r)$ , then probably we could get fairly good lower bounds on  $a_q(n)$ .

Although this problem goes back to Erdős and Rado [ER], very little is known about  $f_q(n, r)$ .

For  $q$  odd,  $n = 2q$  and  $r = 2$ , one can take two disjoint complete graphs on  $q$  vertices each. This shows  $f_q(2q, 2) \geq q(q-1)$ . Actually one has equality, but we do not need it. Using (1) we obtain

$$a_q(2dq) \geq (q-1)^{2dq} \left( \frac{q}{q-1} \right)^d.$$

Using the fact that there is a collection of 300 6-element subsets of [18] without a sunflower of size three, one obtains  $a_3(18) \geq 300 \cdot 12^{12}$  and thus  $a_3(d) \geq (2.179)^d$  for  $d > d_0$ .

## REFERENCES

- [B] F. A. BEHREND, On sets of integers which contain no three elements in arithmetic progression, *Proc. Nat. Acad. Sci.* **23** (1946), 331–332.
- [BB1] T. C. BROWN AND J. P. BUHLER, A density version of a geometric Ramsey theorem, *J. Combin. Theory A* **32** (1982), 20–34.
- [BB2] T. C. BROWN AND J. P. BUHLER, Lines imply spaces in density Ramsey Theory, *J. Combin. Theory A* **36** (1984), 214–220.
- [EFR] P. ERDŐS, P. FRANKL, AND V. RÖDL, The asymptotic number of graphs not containing a fixed subgraph and a problem for hypergraphs having no exponent, *Graphs and Combin.* **2** (1986), 113–121.

- [ER] P. ERDŐS AND R. RADO, Intersection theorems for systems of sets, *J. London Math. Soc.* **35** (1960), 85–90.
- [FK] H. FÜRSTENBERG AND Y. KATZNELSON, An ergodic Szemerédi theorem for  $\mathbb{P}$ -systems and combinatorial theory, *J. Anal. Math.* **45** (1985), 117–168.
- [R] K. F. ROTH, On certain sets of integers, *J. London Math. Soc.* **28** (1953), 104–109.
- [RS] I. Z. RUZSA AND E. SZEMERÉDI, Triple systems with no six points carrying three triangles, *Colloq. Math. Soc. Janos Bolyai* **18** (1978), 939–945.
- [S] E. SZEMERÉDI, On sets of integers containing no  $k$  elements in arithmetic progression, *Acta Arith.* **27** (1975) 199–245.