

1.2 Algebraic Structures of Subsets (II.4 in Driver)

Start with a sample space Ω (any set).

Definition: A collection $\mathcal{F} \subseteq 2^\Omega$ is a _____ if

1. $\Omega \in \mathcal{F}$, $\emptyset \in \mathcal{F}$

2. If $E \in \mathcal{F}$, then $E^c \in \mathcal{F}$.

3. If $E_1, \dots, E_n \in \mathcal{F}$, then $\bigcup_{j=1}^n E_j \in \mathcal{F}$

4. If $E_1, \dots, E_n \in \mathcal{F}$, then $\bigcap_{j=1}^n E_j \in \mathcal{F}$

If, instead of 3, we have the stronger

3'. If $\{E_j\}_{j=1}^\infty$ is a countable set of events in \mathcal{F} , then $\bigcup_{j=1}^\infty E_j \in \mathcal{F}$

then we call \mathcal{F} a _____.

Examples

E.g. $\mathcal{F} = 2^\Omega$

E.g. $\mathcal{F} = \{\emptyset, \Omega\}$

E.g. $\Omega = \{1, 2, 3\}$ $\mathcal{F} = \{\emptyset, \{1\}, \{2, 3\}, \Omega\}$

E.g. $\mathcal{F} = \{B \subseteq \Omega : B \text{ is countable or } B^c \text{ is countable}\}$

Lemma: If I is any index set and $\{\mathcal{F}_i : i \in I\}$ are σ -fields over Ω , then $\bigcap_{i \in I} \mathcal{F}_i$ is a σ -field.

Pf. 1. Ω

2. $E \in \bigcap_i \mathcal{F}_i$

3. $\{E_n\}_{n=1}^\infty \in \bigcap_i \mathcal{F}_i$


Prop: Let $\mathcal{E} \subseteq 2^\Omega$ be any collection of subsets of Ω .

There is a unique smallest σ -field $\sigma(\mathcal{E})$ that contains \mathcal{E} . It is called the σ -field _____ by \mathcal{E} .

Pf.

///

E.g. $\Omega = \{1, 2, 3\}$, $\mathcal{E} = \{\emptyset, \{1, 2\}, \Omega\}$



Exercise

Let $\mathcal{E}_1, \mathcal{E}_2 \subseteq 2^\Omega$. Show that $\sigma(\mathcal{E}_1) = \sigma(\mathcal{E}_2)$ iff:

The Borel σ -Field

Let X be a topological space (e.g. Euclidean space \mathbb{R}^d).

The **Borel σ -field** $\mathcal{B}(X)$ is the σ -field generated by the **open** subsets in X .

$$\mathcal{B}(X) = \sigma\{\text{open subsets of } X\}$$

Events in $\mathcal{B}(X)$ are called **Borel sets**.

Fun Fact



For $X = \mathbb{R}^d$, every open set U is a countable union of open balls

$$U = \bigcup_{i=1}^{\infty} B(x_i, r_i)$$

$$\therefore \mathcal{B}(\mathbb{R}^d) = \sigma\{\text{open balls in } \mathbb{R}^d\}$$

$$d=1: \mathcal{B}(\mathbb{R}) = \sigma\{(a,b) : a < b\} =$$