

Recall that a sequence $\{\mu_n\}_{n \in \mathbb{N}}$ of probability measures on a common measurable metric space (C, \mathcal{B}) is **tight** if, for each $\varepsilon > 0$,
 \exists compact $K_\varepsilon \subseteq C$ s.t. $\mu_n(K_\varepsilon^c) < \varepsilon \quad \forall n \in \mathbb{N}$.

Let $(X_n(t))_{t \in [0,1]}$ be a sequence of continuous stochastic processes in S .

Their laws are $P_n \in \text{Prob}(C([0,1], S), \mathcal{B}(C([0,1], S))$

When is such a sequence tight?

First question: what do compact sets in $C([0,1], S)$ look like?

Theorem: (Arzela-Ascoli) Let S be a complete metric space w Heine-Borel property (closed bounded sets are compact). Then a subset $K \subseteq C([0,1], S)$, does

is compact iff it is closed, pointwise bounded,

and **equicontinuous**: $\exists \beta: [0,1] \rightarrow \mathbb{R}_+$ s.t. $d(x_0, w(s)) \leq \beta(s) \quad \forall s \in [0,1]$.

$\forall t_0 \in [0,1], w \in K, \varepsilon > 0 \quad \exists \delta = \delta(t_0, \varepsilon) > 0$ s.t.

$\forall s \in [0,1], |s - t_0| < \delta \Rightarrow d_S(w(s), w(t_0)) < \varepsilon$.

Lemma: Let $N < \infty$, $\alpha > 0$. Suppose $W \subseteq C([0, 1], S)$ satisfies
 $d_S(w(s), w(t)) \leq N|s-t|^\alpha \quad \forall s, t \in [0, 1]$,
and $d_S(w(x_0), x_0) \leq N$ for some $x_0 \in S$.

Then W is uniformly equicontinuous and uniformly bounded.

Pf. Equicontinuous: $\delta = (\varepsilon/N)^{1/\alpha}$

$$\forall s, t \quad |s-t| < \delta \Rightarrow d_S(w(s), w(t)) \leq N|s-t|^\alpha < N\delta^\alpha = \varepsilon.$$

Bounded: $\forall w \in W, t \in [0, 1]$

$$\begin{aligned} d(x_0, w(t)) &\leq d(x_0, w(b)) + d(w(b), w(t)) \\ &\leq N + \underbrace{N|t-b|^\alpha}_{\leq N} \leq 2N. \quad // \end{aligned}$$

Note: the d_S -closure of an equicontinuous / pointwise bounded set is equicontinuous / pointwise bounded.

Theorem: (Kolmogorov's Tightness Criteria)

Let S be a complete metric space with the Heine-Borel property.

Let $(X_n(t))_{t \in [0,1], n \in \mathbb{N}}$ be a sequence of continuous stochastic processes in S . Suppose $\exists \varepsilon, C > 0$ and $p \geq 1 + \varepsilon$ s.t.

$$\sup_n \mathbb{E}[d_S(X_n(s), X_n(t))^p] \leq C |s-t|^{1+\varepsilon} \quad \forall s, t \in [0,1],$$

and $\{X_n(0)\}_{n \in \mathbb{N}}$ is uniformly bounded w high probability
i.e. is tight.

$$\lim_{N \rightarrow \infty} \sup_n P(d_S(X_n(0), x_0) > N) = 0 \text{ for some } x_0 \in S.$$

$X_n(0) \in \overline{B_N(x_0)}^c$

Then the laws $\{P_n\}_{n \in \mathbb{N}} \subset \text{Prob}(C([0,1], S))$

form a tight sequence of probability measures

on path space.

$$\text{Pf. Recall } \Delta_k(w) = \max_{1 \leq j \leq 2^k} d(w(\frac{j-1}{2^k}), w(\frac{j}{2^k}))$$

$$K_\alpha(w) = 2^{1+\alpha} \sum_{k=0}^{\infty} 2^{\alpha k} \Delta_k(w)$$

$$K_\alpha(X_n) = K_\alpha(t \mapsto X_n(t))$$

By Kolmogorov's Continuity Criteria, for any $\alpha \in (0, \varepsilon/p)$, $K_\alpha(X_n) \in L^p$:

$$\mathbb{E}[K_\alpha(X_n)^p] \leq \frac{C \cdot 2^{p(1+\alpha)}}{(1 - 2^{\alpha - \varepsilon/p})^p} =: M < \infty.$$

Fix any $\alpha \in (0, \varepsilon/p)$, and define

$$W_N^\alpha = \{w \in C([0, 1], S) : K_\alpha(w) \leq N \text{ and } d(w(0), x_0) \leq N\}$$

$$\begin{aligned} \text{Then } P_n((W_N^\alpha)^c) &= P(X_n \notin W_N^\alpha) \\ &= P(K_\alpha(X_n) > N \text{ or } d(X_n(0), x_0) > N) \\ &\leq P(K_\alpha(X_n) > N) + P(d(X_n(0), x_0) > N) \end{aligned}$$

$$\begin{aligned} \text{Markov's Ineq: } &\leq \underbrace{\frac{1}{N^p} \mathbb{E}[K_\alpha(X_n)^p]}_{\xrightarrow[N \rightarrow \infty]{\rightarrow 0}} \leq \underbrace{\sup_m P(d(X_m(0), x_0) > N)}_{\xrightarrow[N \rightarrow \infty]{\rightarrow 0}}. \end{aligned}$$

Thus $\sup_n P_n((W_N^\alpha)^c) \rightarrow 0$ as $N \rightarrow \infty$.

Now set $K_N^\alpha = \overline{W_N^\alpha}$. Then $W_N^\alpha \subseteq K_N^\alpha \therefore (K_N^\alpha)^c \subseteq (W_N^\alpha)^c$

$$\sup_n P_n((K_N^\alpha)^c) \leq \sup_n P_n((W_N^\alpha)^c) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since W_N^α is uniformly equicontinuous and bounded (by the lemma), it follows that $\{K_N^\alpha\}_{N \in \mathbb{N}}$ are all compact. $\therefore \{P_n\}_{n \in \mathbb{N}}$ is tight. $\//\//$