

BROWN MEASURE SUPPORT AND THE FREE MULTIPLICATIVE BROWNIAN MOTION

BRIAN C. HALL AND TODD KEMP

ABSTRACT. The free multiplicative Brownian motion b_t is the large- N limit of Brownian motion B_t^N on the general linear group $GL(N; \mathbb{C})$. We prove that the Brown measure for b_t —which is an analog of the empirical eigenvalue distribution for matrices—is supported on the closure of a certain domain Σ_t in the plane. The domain Σ_t was introduced by Biane in the context of the large- N limit of the Segal–Bargmann transform associated to $GL(N; \mathbb{C})$.

We also consider a two-parameter version, $b_{s,t}$: the large- N limit of a related family of diffusion processes on $GL(N; \mathbb{C})$ introduced by the second author. We show that the Brown measure of $b_{s,t}$ is supported on the closure of a certain planar domain $\Sigma_{s,t}$, generalizing Σ_t , introduced by Ho.

In the process, we introduce a new family of spectral domains related to any operator in a tracial von Neumann algebra: the L_n^p -spectrum for $n \in \mathbb{N}$ and $p \geq 1$, a subset of the ordinary spectrum defined relative to potentially-unbounded inverses. We show that, in general, the support of the Brown measure of an operator is contained in its L_2^2 -spectrum.

CONTENTS

1. Introduction	1
2. Preliminaries	4
2.1. Lie group Brownian motions and their large- N limits	4
2.2. The domains Σ_t	6
2.3. Brown measure	8
3. Free Segal–Bargmann transform	10
3.1. Using free probability	10
3.2. As an integral operator	10
3.3. From the generalized Segal–Bargmann transform	11
4. An outline of the proof of Theorem 1.1	12
5. The two-parameter case	13
5.1. Brownian motions	13
5.2. Segal–Bargmann transform	14
5.3. The domains $\Sigma_{s,t}$	15
5.4. The main result	16
6. Proofs	16
6.1. A general result on the support of the Brown measure	16
6.2. Computing the L_2^2 spectrum	19
Acknowledgments	23
References	23

1. INTRODUCTION

One of the core theorems in random matrix theory is the *circular law*. Suppose C^N is an $N \times N$ complex matrix whose entries are independent centered normal random variables of variance $\frac{1}{N}$. Then

Kemp’s research is supported in part by NSF CAREER Award DMS-1254807 and NSF Grant DMS-1800733.

the empirical eigenvalue distribution of C^N (the random probability measure placing points of equal mass at the eigenvalues) converges almost surely to the uniform probability measure on the unit disk as $N \rightarrow \infty$. This theorem is due to Ginibre [16] and C^N is often called a *Ginibre ensemble*. The circular law has been incrementally generalized to its strongest form where the entries are independent but can have any distribution with two finite moments [1, 17, 18, 34].

We can recast this as a theorem about *matrix-valued Brownian motion*. In a finite-dimensional real Hilbert space, there is a canonical Brownian motion, constructed by adding independent standard real Brownian motions in all the directions of any orthonormal basis. (See Section 2.1.1 below.) Let us regard the space $M_N(\mathbb{C})$ of complex $N \times N$ matrices as a real vector space of dimension $2N^2$, and equip it with the inner product

$$\langle X, Y \rangle_N = N \operatorname{Re} \operatorname{Trace}(X^*Y). \quad (1.1)$$

Then the associated Brownian motion C_t^N has the same law as the Ginibre ensemble, scaled by a factor of \sqrt{t} . (The factor N in front of the Hilbert–Schmidt inner product is the correct choice to give the entries variance of order $1/N$.) Hence, matrix-valued Brownian motion C_t^N has eigenvalues that concentrate uniformly in the disk of radius \sqrt{t} as $N \rightarrow \infty$.

In this paper, we are interested in the Brownian motion B_t^N on the general linear group $\operatorname{GL}(N; \mathbb{C})$. One nice geometric way to define this object is by the *rolling map*. The tangent space to the identity in $\operatorname{GL}(N; \mathbb{C})$ (i.e., the Lie algebra of this Lie group) is all of $M_N(\mathbb{C})$. Take the Brownian paths in the tangent space, and roll them onto the group; this yields the paths of B_t^N . Since the paths are not smooth this rolling is accomplished by a *stochastic* differential equation for B_t^N in terms of C_t^N (cf. (2.2)). In particular, for small time, B_t^N and C_t^N are “close”, and so it is natural to expect that the eigenvalues of B_t^N should follow a deformation of the circular law of radius \sqrt{t} .

For a normal matrix A , the eigenvalues are encoded in the matrix moments $\{\operatorname{Trace}[A^k(A^*)^\ell]\}_{k, \ell \in \mathbb{N}}$. However, the ensemble C_t^N is almost surely non-normal for any $t > 0$; in fact, a stronger statement is true: with probability 1, C_t^N is non-normal for all $t > 0$ [27, Proposition 4.15]). The lack of normality presents significant hurdles to understanding the limit behavior of its eigenvalues, whose connection to matrix moments is quite a bit more subtle. Nevertheless, the *mixed* moments of C_t^N and $(C_t^N)^*$ (i.e., traces of all words in these non-commuting matrices) do have a meaningful large- N limit: in the language of free probability, the ensemble C_t^N converges in **-distribution* to an operator c_t , cf. [35] (see Section 2.1.2).

This *circular Brownian motion* c_t , living in noncommutative probability space, does not have eigenvalues, and is not normal, so it does not have a spectral resolution. But there is an analog known as the **Brown measure** μ_{c_t} [9]. Each operator a in a tracial von Neumann algebra has an associated Brown measure μ_a , which is a probability measure supported in the spectrum of a in \mathbb{C} . If a is normal, μ_a is the usual spectral measure inherited from the spectral theorem; if A is an $N \times N$ matrix, its Brown measure is simply its empirical eigenvalue distribution. We discuss the Brown measure in general in Section 2.3 below. Girko’s proof [17] of the general circular law begins by proving that the Brown measure of c_1 is uniform on the unit disk, and then shows that the empirical eigenvalue distribution of C^N actually converges to the Brown measure of the large- N limit.

Meanwhile, the Brownian motion B_t^N on the group $\operatorname{GL}(N; \mathbb{C})$ also has a large- N limit in terms of **-distribution*: an operator b_t known as the **free multiplicative Brownian motion**. It was introduced by Biane [4, 5] and conjectured to be the large- N limit of B_t^N ; this conjecture was proven by the second author in [27]. The first step in understanding the large- N behavior of the eigenvalues of B_t^N is to determine the Brown measure μ_{b_t} of b_t . It is a probability measure supported in the spectrum of b_t ; but b_t is a complicated object, and in particular its spectrum is completely unknown.

In this paper, we identify a closed set $\bar{\Sigma}_t$ (see Section 2.2) which contains the support of the Brown measure μ_{b_t} . The region Σ_t was introduced by Biane in [5] in the context of the *Segal–Bargmann transform* (or “Hall transform”) associated to the unitary group $U(N)$ and its complexification $\operatorname{GL}(N; \mathbb{C})$ (cf. [20]). Biane introduced a *free Hall transform* \mathcal{G}_t , which he understood as a sort of large- N limit of the

Hall transform for $U(N)$. The transform \mathcal{G}_t is an integral operator which maps functions on the unit circle to a space \mathcal{A}_t of holomorphic functions on the region $\Sigma_t \subset \mathbb{C}$, whose definition falls out of the complex analysis used in Biane's proofs.

The meaning of the region Σ_t and its relation to the free multiplicative Brownian motion b_t have remained mysterious. One clue to its origin comes from the holomorphic functional calculus. Using the metric properties of \mathcal{G}_t , Biane showed that one can make sense of $F(b_t)$, as a possibly unbounded operator, for any $F \in \mathcal{A}_t$. Now, if the spectrum of b_t were contained in Σ_t , properties of the holomorphic functional calculus would show that $F(b_t)$ is a *bounded* operator for all F in $\mathcal{H}(\Sigma_t)$ and thus for all F in \mathcal{A}_t , which is (presumably) not the case. On the other hand, the fact that $F(b_t)$ can be defined at all — even as an unbounded operator — suggests that the spectrum of b_t is at least contained in the *closure* of Σ_t . Such a results would then imply that the support of the Brown measure of b_t is contained in $\overline{\Sigma}_t$. The latter statement is the main theorem of this paper.

Theorem 1.1. *For all $t > 0$, the support of the Brown measure μ_{b_t} of the free multiplicative Brownian motion b_t is contained in $\overline{\Sigma}_t$.*

We expect that the large- N limit of the empirical eigenvalue distribution of the Brownian motion B_t^N on $GL(N; \mathbb{C})$ coincides with the Brown measure μ_{b_t} of the free multiplicative Brownian motion. If that is the case, the eigenvalues of B_t^N should concentrate in $\overline{\Sigma}_t$ for large N ; this claim is very clearly supported by numerical evidence. Figure 1 shows simulations of B_t^N with $N = 2000$ and four different values of t , plotted along with the domains Σ_t . (The domain for $t \geq 4$ has a small hole around the origin, as can be seen in the bottom two images in the figure.)

We also consider a two-parameter version $b_{s,t}$ of the free multiplicative Brownian motion and show that its Brown measure is supported on the closure of a certain domain $\Sigma_{s,t}$, introduced by Ho [26]. These domains similarly arise in the large- N limit of the two-parameter Segal–Bargmann transform in the Lie group setting. The precise statement and proof can be found in Section 5.

The strategy we employ to prove Theorem 1.1 is of independent interest, as it provides a new restriction on the support of the Brown measure of an operator, in terms of a family of spectral domains associated to the operator. Let (\mathcal{A}, τ) be a tracial von Neumann algebra and let $a \in \mathcal{A}$. As noted, the Brown measure μ_a is supported in the spectrum of a ; in fact, its support set may be a strict subset of the spectrum. Recall that the spectrum of a is the complement of the resolvent set of a , which is the set of $\lambda \in \mathbb{C}$ for which $a - \lambda$ is invertible (meaning that $(a - \lambda)^{-1}$ is a bounded operator). The rich structure of \mathcal{A} and the trace τ give a natural but heretofore unstudied generalization of these spaces: we may ask that the inverse $(a - \lambda)^{-1}$ exist but not necessarily be bounded, instead insisting that it is in $L^p(\mathcal{A}, \tau)$ for some $p \geq 1$. (The $p = \infty$ case coincides with the usual resolvent set.) What is more, given the often bizarre algebraic properties of non-normal operators (which can, for example, be nilpotent), we may ask that some power $(a - \lambda)^n$ have an inverse in $L^p(\mathcal{A}, \tau)$. (Unless a is normal, this is *not* equivalent to $a - \lambda$ having an inverse in $L^{np}(\mathcal{A}, \tau)$.) The set of $\lambda \in \mathbb{C}$ for which $(a - \lambda)^n$ has an inverse in L^p (and for which certain uniform local bounds hold) is called the L_n^p -**resolvent set** of a , and its complement is $\text{spec}_n^p(a)$, the L_n^p -**spectrum** of a .

Our key observation leading to the proof of Theorem 1.1 is the following new description of (a closed set containing) the support of the Brown measure.

Theorem 1.2. *For any operator a in a tracial von Neumann algebra, the support of the Brown measure μ_a is contained in $\text{spec}_2^2(a)$ (which is, itself, a subset of the spectrum of a).*

A detailed discussion of these generalized spectral domains, and the proof of Theorem 1.2, can be found in Section 6.1. Once this result is established, we use Biane's "free Hall transform" \mathcal{G}_t to show that $\text{spec}_2^2(b_t) = \overline{\Sigma}_t$, which proves that the support of μ_{b_t} is contained in $\overline{\Sigma}_t$, establishing Theorem 1.1.

A more detailed version of this outline of the proof is contained in Section 4, (cf. Theorems 4.1 and 4.2); the complete proofs can then be found in Section 6.




FIGURE 1. Simulations of the eigenvalues of B_t^N shown with the domain Σ_t , for $N = 2000$ and $t = 2, t = 3.9, t = 4$, and $t = 4.1$

2. PRELIMINARIES

In this section, we provide background on the objects in the statement of our main theorem—the free multiplicative Brownian motion, the Brown measure, and the domains Σ_t .

2.1. Lie group Brownian motions and their large- N limits.

2.1.1. *Lie group Brownian motions.* Let H be a finite-dimensional real Hilbert space. The *Brownian motion on H* , W_t^H , is the diffusion process defined by

$$W_t^H = \sum_{j=1}^d B_t^j e_j \quad (2.1)$$

where $\{e_1, \dots, e_d\}$ is an orthonormal basis for H , and $\{B_t^j\}_{j=1}^d$ are i.i.d. standard (real) Brownian motions. The law of this process is invariant under rotations, and hence does not depend on which orthonormal basis is chosen.

Let $G \subset \text{GL}(N; \mathbb{C})$ be a matrix Lie group (in $M_N(\mathbb{C})$), and let $\mathfrak{g} \subset M_N(\mathbb{C})$ be its Lie algebra. A choice of inner product on \mathfrak{g} induces a left-invariant Riemannian metric on G . As a Riemannian manifold, then, G has a well-defined Brownian motion: the diffusion with infinitesimal generator given by half

the Laplacian. In the Lie group setting there is a simple description of the Brownian motion B_t^G in terms of the Brownian motion W_t^g on the Lie algebra (as in (2.1)):

$$dB_t^G = B_t^G \circ dW_t^g, \quad B_0^G = I. \quad (2.2)$$

The \circ denotes the Stratonovich stochastic integral. This Stratonovich SDE can be converted to an Itô SDE; the form of the resulting equation depends on the structure of the group (cf. [29, p. 116]).

For our purposes, the two relevant Lie groups are the unitary group $U(N)$ whose Lie algebra is $\mathfrak{u}(N) = M_N^{\text{s.a.}}(\mathbb{C})$ (self-adjoint matrices), and the general linear group $GL(N; \mathbb{C})$ whose Lie algebra consists of all complex matrices $\mathfrak{gl}(N; \mathbb{C}) = M_N(\mathbb{C})$. (In the unitary case, we follow the physicists' convention; mathematicians typically use skew-self-adjoint matrices.) Using the inner product (1.1) on both these Lie algebras, we obtain Brownian motions which we will denote thus:

$$W_t^{\mathfrak{u}(N)} = X_t^N \quad \text{and} \quad W_t^{\mathfrak{gl}(N; \mathbb{C})} = C_t^N.$$

To be more explicit, C_t^N is the *Ginibre Brownian motion*, which has i.i.d. entries that are all complex Brownian motions of variance t/N , and X_t^N is the *Wigner Brownian motion*, which is Hermitian with i.i.d. upper triangular entries, with complex Brownian motions above the diagonal and real Brownian motions on the diagonal, each of variance t/N .

The Brownian motions on the groups, which we denote $B_t^{\mathfrak{u}(N)} = U_t^N$ and $B_t^{\mathfrak{gl}(N; \mathbb{C})} = B_t^N$, then satisfy Stratonovich SDEs given by (2.2). These equations can be written in Itô form as follows:

$$dU_t^N = iU_t^N dX_t^N - \frac{1}{2}U_t^N dt \quad \text{and} \quad dB_t^N = B_t^N dC_t^N \quad (2.3)$$

(both started at the identity matrix). These defining SDEs play the role of the rolling map described in the introduction.

2.1.2. The large- N limits. The four processes X_t^N , C_t^N , U_t^N , and B_t^N all have large- N limits in the sense of free probability theory. (For a thorough introduction to free probability theory and its connection to random matrix theory, the reader is directed to [30] and [31].) The limits are one-parameter families of operators x_t , c_t , u_t , and b_t all living in a noncommutative probability space (\mathcal{B}, τ) . (More precisely, \mathcal{B} is a finite von Neumann algebra and τ is a faithful, normal, tracial state.) The sense of convergence is *almost sure convergence of the finite-dimensional noncommutative distributions*, defined as follows.

Definition 2.1. Let A_t^N be a sequence of $M_N(\mathbb{C})$ -valued stochastic processes (all defined on the same sample space). Let (\mathcal{A}, τ) be a noncommutative probability space, and let $a_t \in \mathcal{A}$ for each $t > 0$. We say A_t^N **converges to a_t in finite-dimensional noncommutative distributions** if, for each $n \in \mathbb{N}$ and times $t_1, \dots, t_n \geq 0$, and each noncommutative polynomial P in $2n$ indeterminates,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \text{Trace} [P(A_{t_1}^N, \dots, A_{t_n}^N, (A_{t_1}^N)^*, \dots, (A_{t_n}^N)^*)] = \tau [P(a_{t_1}, \dots, a_{t_n}, a_{t_1}^*, \dots, a_{t_n}^*)] \quad \text{almost surely.}$$

The limit of X_t^N was identified by Voiculescu [35]; it is known as **free additive Brownian motion** x_t , and can be constructed on a Fock space. From here, one can derive the C_t^N case by noting that $C_t^N = \frac{1}{\sqrt{2}}(X_t^N + iY_t^N)$ where Y_t^N is an independent copy of X_t^N . Given the independence and rotational invariance, it follows by standard results on asymptotic freeness that the large- N limit of C_t^N can be represented as $c_t = \frac{1}{\sqrt{2}}(x_t + iy_t)$ where $\{x_t, y_t\}$ are *freely* independent free additive Brownian motions; we call c_t a **free circular Brownian motion**.

Since the unitary and general linear Brownian motions U_t^N and B_t^N are defined as solutions of SDEs involving X_t^N and C_t^N , good candidates for their large- N limits are given by *free* SDEs involving x_t and c_t . (Free stochastic analysis was introduced in [7] and further developed in [8, 28]; the reader may also consult the background sections of [11, 27] for succinct summaries of the relevant concepts.) The **free unitary Brownian motion** u_t and **free multiplicative Brownian motion** b_t are defined as solutions to the free SDEs

$$du_t = iu_t dx_t - \frac{1}{2}u_t dt \quad \text{and} \quad db_t = b_t dc_t \quad (2.4)$$

(both started at 1), mirroring the matrix SDEs of (2.3).

The free unitary Brownian motion u_t was introduced by Biane in [4], wherein he also showed that it is the large- N limit of the unitary Brownian motion U_t^N as a process (as in Definition 2.1). In particular, in the case of a single time t ($n = 1$ in the definition, $t_1 = t$), since $u_t^* = u_t^{-1}$, the statement is simply that the trace moments $\frac{1}{N} \text{Trace}[(U_t^N)^k]$, $k \in \mathbb{Z}$, converge almost surely as $N \rightarrow \infty$ to $\tau[u_t^k]$. The numbers $\nu_k(t) := \tau[u_t^k]$, meanwhile, are the moments of a probability measure ν_t on the unit circle. Biane computed these limit moments, which had already appeared in work of Singer [33] in Yang–Mills theory on the plane in an asymptotic regime. They are given by

$$\nu_k(t) := \int_{\mathbb{C}^*} \omega^k \nu_t(d\omega) = e^{-\frac{|k|}{2}t} \sum_{j=0}^{|k|-1} \frac{(-t)^j}{j!} |k|^{j-1} \binom{|k|}{j} \quad (2.5)$$

for $n \in \mathbb{Z} \setminus \{0\}$, with $\nu_0(t) = 1$.

From here, using complex analytic techniques, Biane completely determined the measure ν_t . It is supported on an arc in the unit circle (symmetric about 1) that is a proper subset for $t < 4$ and is fully supported on the circle for $t \geq 4$:

$$\text{supp } \nu_t = \left\{ e^{i\theta} : |\theta| \leq \frac{1}{2} \sqrt{t(4-t)} + \arccos \left(1 - \frac{t}{2} \right) \right\}. \quad (2.6)$$

Biane also gave an implicit description of the measure ν_t , which has a real analytic density on the interior of its support, but we do not need this description presently.

The key to analyzing ν_t was determining a certain analytic transform of ν_t in the unit disk \mathbb{D} . Let

$$\psi_t(z) := \int_{\partial\mathbb{D}} \frac{\omega z}{1 - \omega z} \nu_t(d\omega), \quad z \in \mathbb{D}$$

denote the moment generating function (with no constant term). The function ψ_t has a continuous extension to the closed disk $\overline{\mathbb{D}}$; this is tantamount to the fact, as Biane proved, that ν_t possesses a continuous density on $\partial\mathbb{D}$. Of greater computational use is the following function:

$$\chi_t(z) := \frac{\psi_t(z)}{1 + \psi_t(z)} \quad (2.7)$$

which also has a continuous extension to $\overline{\mathbb{D}}$. In fact, χ_t is one-to-one on the open disk, and its inverse is analytic, with the following simple explicit formula:

$$f_t(z) := \chi_t^{(-1)}(z) = z e^{\frac{t}{2} \frac{1+z}{1-z}}. \quad (2.8)$$

It is from this identity that the explicit formulas (2.5) and (2.6) are derived.

Biane also introduced the free multiplicative Brownian motion process b_t in [5], where he conjectured that it is the large- N limit of the $\text{GL}(N; \mathbb{C})$ Brownian motion B_t^N . Given the non-normality of the matrices and operators involved, this turned out to be a difficult problem that took nearly two decades to solve; the second author proved this in [27]. Now, the “holomorphic” moments $\tau[b_t^k]$ are easily shown to have the value 1 for all k (see also (5.6)). But since the process b_t is not normal, these moments do not determine much of the noncommutative distribution. The free SDE (2.4) that defines b_t allows for any mixed moment in b_t and b_t^* to be computed (iteratively) given enough patience; see [27, Proposition 1.8] for some notable examples. There is, at present, no known simple description of the full noncommutative distribution of this complicated object. Its spectrum is also unknown.

2.2. The domains Σ_t . In this section, we describe the family of domains $\Sigma_t \subset \mathbb{C}$, introduced by Biane in [5], which enter into the statement of our main result. They arose in the context of the free Segal–Bargmann transform (see Section 3), which connects u_t and b_t . For this reason, they are related to the inverse shifted moment generating series f_t (2.8) of the spectral measure ν_t of the free unitary Brownian motion.




FIGURE 2. The domains Σ_t with $t = 3$ and $t = 4.05$, with the unit circle shown for comparison

It is easily verified that if $|z| = 1$, then $|f_t(z)| = 1$. There are, however, points z with $|z| \neq 1$ for which $|f_t(z)|$ is nevertheless equal to 1.

Proposition 2.2. *For all $t > 0$, consider the set*

$$E_t = \overline{\{z \in \mathbb{C} \mid |z| \neq 1, |f_t(z)| = 1\}}$$

and define Σ_t to be the connected component of the complement of E_t containing 1. Then Σ_t is bounded for all $t > 0$, Σ_t is simply connected for $t \leq 4$, and Σ_t is doubly connected for $t > 4$. In all cases, we have

$$\partial\Sigma_t = E_t.$$

These properties of the region Σ_t were proved by Biane in [5]; see especially pp. 273–274. See also [26, Section 4.2]. The closure in the definition of the set E_t is needed to fill in the points (at most two of them) where $\partial\Sigma_t$ intersects the unit circle. Figure 2 shows the domain Σ_t with $t = 3$ and $t = 4.05$, with the unit circle shown for comparison. Figure 3 then shows the transitional case $t = 4$ in more detail. In all cases, 1 is in Σ_t and 0 is not in $\overline{\Sigma}_t$.




FIGURE 3. The region Σ_t with $t = 4$ (left) and a detail thereof (right)

An important property of the region Σ_t , which follows from the just-cited results of Biane [5], involves the support arc of the spectral measure ν_t of free unitary Brownian motion. This result is crucial to the proof of our main theorem.

Proposition 2.3. *For all $t > 0$, the function f_t maps $\mathbb{C} \setminus \overline{\Sigma}_t$ injectively onto $\mathbb{C} \setminus \text{supp } \nu_t$.*

This is a typical “slit plane” conformal map; see Figure 4.




FIGURE 4. The map f_t takes $\mathbb{C} \setminus \overline{\Sigma}_t$ (left) injectively onto $\mathbb{C} \setminus \text{supp } \nu_t$ (right). Shown for $t = 3$

2.3. Brown measure. We work in the context of a sufficiently rich noncommutative probability space: a tracial von Neumann algebra.

Definition 2.4. *A **tracial von Neumann algebra** is a finite von Neumann algebra \mathcal{A} together with a faithful, normal, tracial state $\tau : \mathcal{A} \rightarrow \mathbb{C}$.*

Recall that a *state* τ is norm-one linear functional taking non-negative elements to non-negative real numbers. (Such a functional necessarily satisfies $\tau(1) = \|\tau\| = 1$.) A state τ is called *faithful* if $\tau(a^*a) > 0$ for all $a \neq 0$, it is called *normal* if it is continuous with respect to the strong operator topology, and it is called *tracial* if $\tau(ab) = \tau(ba)$ for all $a, b \in \mathcal{A}$.

Let (\mathcal{A}, τ) be a noncommutative probability space. For each element a of \mathcal{A} , it is possible to define a probability measure μ_a on \mathbb{C} called the **Brown measure** of a , which should be interpreted as something like an empirical eigenvalue distribution for the operator a . The definitions and properties stated in this section may be found in Brown’s original paper [9] and in [30, Chapter 11].

We first recall the notion of the **Fuglede–Kadison determinant** of a [14, 15], denoted $\Delta(a)$, which is most easily defined by a limiting process:

$$\log \Delta(a) = \lim_{\varepsilon \rightarrow 0} \frac{1}{2} \tau[\log(a^*a + \varepsilon)]. \quad (2.9)$$

In general, $\log \Delta(a)$ may have the value $-\infty$, in which case, $\Delta(a) = 0$. If, for example, $\mathcal{A} = M_N(\mathbb{C})$ and τ is the normalized trace, then $\Delta(a) = |\det a|^{1/N}$, where $\det a$ is the ordinary determinant of a .

For a tracial von Neumann algebra (\mathcal{A}, τ) , the Brown measure of an element $a \in \mathcal{A}$ is then defined as

$$\mu_a = \frac{1}{4\pi} \nabla_\lambda^2 \log(\Delta(a - \lambda)),$$

where ∇_λ^2 is the Laplacian with respect to λ , computed in the distributional sense. It can be shown that this distributional Laplacian is represented by a probability measure on the plane.

Proposition 2.5. *Let a be an element of \mathcal{A} and let μ_a be its Brown measure. Then the following results hold.*

- (1) *The measure μ_a is a probability measure on the plane.*

- (2) The support of μ_a is contained in spectrum of a , but the two sets do not coincide in general.
(3) For all non-negative integers n ,

$$\int z^n d\mu_a(z) = \tau[a^n],$$

and if a is invertible, the same result holds for all integers n .

Using the limiting formula (2.9) for the Flugede–Kadison determinant, we may give a limiting formula for the Brown measure. With the notation

$$a_\lambda := a - \lambda,$$

we have

$$\mu_a = \frac{1}{4\pi} \lim_{\varepsilon \rightarrow 0} \{ \nabla_\lambda^2 \tau[\log(a_\lambda^* a_\lambda + \varepsilon)] d^2 \lambda \}, \quad (2.10)$$

where $d^2 \lambda$ is the two-dimensional Lebesgue measure on \mathbb{C} and the limit is in the weak sense. Furthermore, the Laplacian on the right-hand side of (2.10) can be computed explicitly [30, Section 11.5], giving still another formula for the Brown measure:

$$\mu_a = \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0} \{ \varepsilon \tau[(a_\lambda^* a_\lambda + \varepsilon)^{-1} (a_\lambda a_\lambda^* + \varepsilon)^{-1}] d^2 \lambda \}. \quad (2.11)$$

The following result follows easily from (2.11).

Corollary 2.6. *Suppose the quantity*

$$\tau[(a_\lambda^* a_\lambda + \varepsilon)^{-1} (a_\lambda a_\lambda^* + \varepsilon)^{-1}] \quad (2.12)$$

is bounded uniformly for all $\varepsilon > 0$ and all λ in a neighborhood of some value λ_0 . Then λ_0 does not belong to the support of the Brown measure μ_a .

In particular, if λ_0 belongs to the resolvent set of a , it is not hard to see that the quantity (2.12) has a finite limit as $\varepsilon \rightarrow 0$, for all λ in a neighborhood of λ_0 , so that the corollary applies. Thus, Corollary 2.6 implies Point 2 of Proposition 2.5. Although the support of the Brown measure can be a proper subset of the spectrum, there are many interesting examples in which the two sets coincide.

We close this section by noting two important special cases of the Brown measure.

- When $\mathcal{A} = M_N(\mathbb{C})$ and τ is the normalized trace, the Brown measure of a matrix A is its empirical eigenvalue distribution. That is,

$$\mu_A = \frac{1}{N} \sum_{j=1}^n \delta_{\lambda_j},$$

where $\lambda_1, \dots, \lambda_N$ are the eigenvalues of A , listed with their algebraic multiplicity.

- For a normal element a of \mathcal{A} , Brown measure coincided with the spectral measure: μ_a is just the composition of the projection-valued spectral resolution with τ :

$$\mu_a(V) = \tau(E^a(V)), \quad V \in \text{Borel}(\mathbb{C}),$$

where E^a is the projection-valued measure associated to a by the spectral theorem.

Since the free multiplicative Brownian motion b_t is neither normal nor finite-dimensional, neither of the preceding cases applies.

3. FREE SEGAL–BARGMANN TRANSFORM

Recall from the Section 2.1 that the law ν_t of free unitary Brownian motion is a probability measure on the unit circle that represents the limiting empirical eigenvalue distribution for Brownian motion in the unitary group. In [5], Biane introduced a “free Hall transform” \mathcal{G}_t that maps $L^2(\partial\mathbb{D}, \nu_t)$ into $\mathcal{H}(\Sigma_t)$, the space of holomorphic functions on the domain Σ_t . In this section, we recall both the original construction of \mathcal{G}_t given by Biane and a realization given by the authors together with Driver [13] and Cébron [10]. The transform \mathcal{G}_t will be a crucial tool in the proof of our main theorem.

3.1. Using free probability. Let u_t be a free unitary Brownian motion and b_t a free multiplicative Brownian that is freely independent from u_t , both living in a tracial von Neumann algebra (\mathcal{B}, τ) . In Biane’s approach, the map \mathcal{G}_t is characterized by the requirement that for each Laurent polynomial p , we have

$$(\mathcal{G}_t p)(b_t) = \tau[p(b_t u_t)|b_t], \quad (3.1)$$

where $\tau[\cdot|b_t]$ is the conditional trace with respect to the algebra generated by b_t . (Compare Theorem 8 and the computations following it in [5].) If, for example, p is the polynomial $p(u) = u^2$, then it is not hard to compute (cf. [30, p. 55]) that

$$\begin{aligned} \tau[b_t u_t b_t u_t | b_t] &= \tau(u_t)^2 b_t^2 + (\tau(u_t^2) - \tau(u_t)^2) \tau(b_t) b_t \\ &= e^{-t}(b_t^2 - t b_t), \end{aligned}$$

where we have used the following moment formulas: $\tau(u_t) = e^{-t}$, $\tau(u_t^2) = e^{-t}(1 - t)$, and $\tau(b_t) = 1$. (The moments of u_t are computed in Definition 2 and Lemma 1 of [4], while the moment of b_t is the $s = t$ case of (5.6).) Thus, in this case, $\mathcal{G}_t p$ is also a polynomial, given by

$$(\mathcal{G}_t p)(b) = e^{-t}(b^2 - t b). \quad (3.2)$$

As explained in Section 3.3, the map \mathcal{G}_t can be viewed as the large- N limit of the generalized Segal–Bargmann transform over $U(N)$ introduced by the first author in [20]. The motivation for Biane’s definition of \mathcal{G}_t is the stochastic approach to the generalized Segal–Bargmann transform developed by Gross and Malliavin [19].

3.2. As an integral operator. Using the subordination method developed in [6], Biane realized \mathcal{G}_t as an integral operator mapping $L^2(\partial\mathbb{D}, \nu_t)$ into $\mathcal{H}(\Sigma_t)$. Explicitly,

$$(\mathcal{G}_t f)(z) = \int_{\partial\mathbb{D}} f(\omega) \frac{|1 - \chi_t(\omega)|^2}{(z - \chi_t(\omega))(z^{-1} - \overline{\chi_t(\omega)})} d\nu_t(\omega), \quad z \in \Sigma_t, \quad (3.3)$$

where χ_t was defined in (2.7). (See [5, Theorem 8] and the computations that follow it, along with Proposition 13.) Here, for $\omega \in \partial\mathbb{D}$, $\chi_t(\omega)$ denotes the value of the unique continuous extension of χ_t to $\overline{\mathbb{D}}$; in other words, it is the limiting value of $\chi_t(\zeta)$ as $\zeta \in \mathbb{D}$ approaches ω from *inside* the unit disk. Note that for $\omega \in \partial\mathbb{D}$ both $\chi_t(\omega)$ and $1/\overline{\chi_t(\omega)}$ lie on the boundary of Σ_t , so that the integrand in (3.3) is a holomorphic function of z for z in the interior of Σ_t . Biane showed that, for $t \neq 4$, the map \mathcal{G}_t is injective, so that it is possible to identify $L^2(\partial\mathbb{D}, \nu_t)$ with its image:

$$\mathcal{A}_t := \text{Image}(\mathcal{G}_t) \subset \mathcal{H}(\Sigma_t).$$

Now, let us define $L_{\text{hol}}^2(b_t, \tau)$ to be the closure in the noncommutative L^2 space $L^2(\mathcal{B}, \tau)$ of the space of elements of the form $p(b_t)$, where p is a Laurent polynomial in one variable. That is to say, $L_{\text{hol}}^2(b_t, \tau)$ is the closure of the span of the positive and negative integer powers of b_t , *not* including any powers of b_t^* . (Biane used a slightly different definition that is easily seen to be equivalent to this one.) Biane showed that for $t \neq 4$, there is a bijection between \mathcal{A}_t and $L_{\text{hol}}^2(b_t, \tau)$ uniquely determined by the condition that for each Laurent polynomial p , we have

$$p \mapsto p(b_t).$$

Note that for $t \neq 4$, the space $L^2_{\text{hol}}(b_t, \tau)$ is identified with the space \mathcal{A}_t of holomorphic functions. Nevertheless, the noncommutative L^2 norm on $L^2_{\text{hol}}(b_t, \tau)$ does not correspond to an L^2 norm on \mathcal{A}_t with respect to any measure on Σ_t . (It is, instead, the Hilbert space norm induced by a certain reproducing kernel on \mathcal{A}_t which is induced by the integral kernel of \mathcal{G}^t , cf. (3.3).)

For a general $f \in \mathcal{A}_t$, we will write the corresponding element of $L^2_{\text{hol}}(b_t, \tau)$ suggestively as $f(b_t)$ and think of the map $f \mapsto f(b_t)$ as variant of the usual holomorphic functional calculus. That is to say, we think of the map from \mathcal{A}_t to $L^2_{\text{hol}}(b_t, \tau)$ as “evaluation on b_t .” Note, however, that elements of $L^2_{\text{hol}}(b_t, \tau)$ are in general unbounded operators.

Theorem 3.1 (Biane’s Free Hall Transform). *For all $t > 0$ with $t \neq 4$, the map \mathcal{G}_t is a unitary isomorphism from $L^2(\partial\mathbb{D}, \nu_t)$ to \mathcal{A}_t , where the norm on \mathcal{A}_t is defined by identification with $L^2_{\text{hol}}(b_t, \tau)$. In particular, we have*

$$\|f\|_{L^2(\partial\mathbb{D}, \nu_t)} = \|(\mathcal{G}_t f)(b_t)\|_{L^2_{\text{hol}}(b_t, \tau)}$$

for all $f \in L^2(\partial\mathbb{D}, \nu_t)$.

When $t = 4$, the preceding theorem is not known to hold, because it is not known that \mathcal{G}_t is injective. But one still has a theorem, as follows. One considers at first the map $p \mapsto \mathcal{G}_t p$ on polynomials and then constructs a map from the space of polynomials into $L^2_{\text{hol}}(b_t, \tau)$ by mapping p to $(\mathcal{G}_t p)(b_t)$. This map is isometric for all $t > 0$ and it extends to a unitary map of $L^2(\partial\mathbb{D}, \nu_t)$ onto $L^2_{\text{hol}}(b_t, \tau)$; see Section 6.2 for details.

In light of the preceding discussion, we expect that, at least for $t \neq 4$, the spectrum of b_t will *not* be contained in Σ_t . After all, if such a containment held, the operator $f(b_t)$, $f \in \mathcal{A}_t \subset \mathcal{H}(\Sigma_t)$ would presumably be computable by the holomorphic functional calculus, in which case, $f(b_t)$ would be a bounded operator. But actually, *every* element of $L^2_{\text{hol}}(b_t, \tau)$ arises as $f(b_t)$ for some $f \in \mathcal{A}_t$, and the elements of $L^2_{\text{hol}}(b_t, \tau)$ are in general unbounded operators.

On the other hand, since we are able to define $f(b_t)$ for any $f \in \mathcal{A}_t$, at least as an unbounded operator, it seems reasonable to expect that the spectrum of Σ_t is contained in the *closure* of Σ_t . Our main result, that the Brown measure of b_t is supported in $\overline{\Sigma}_t$, is a step toward establishing this claim; compare Proposition 2.5.

3.3. From the generalized Segal–Bargmann transform. In 1994, the first author introduced a generalized Segal–Bargmann transform for compact Lie groups [20]. In the case of the unitary group $U(N)$, the transform, which we denote presently as \mathcal{G}_t^N , maps $L^2(U(N), \rho_t)$ to the space of *holomorphic* functions in $L^2(\text{GL}(N; \mathbb{C}), \gamma_t)$. (Note: in [20] and follow-up work such as [13], the transform was often denoted B_t ; to avoid clashing with our present notation B_t^N for the Brownian motion on $\text{GL}(N; \mathbb{C})$, we use \mathcal{G}_t^N instead for the Segal–Bargmann transform here.) Here ρ_t and γ_t are *heat kernel measures*—that is, the distributions at time t of Brownian motions on the respective groups, starting at the identity. The transform is defined as

$$\mathcal{G}_t^N f = (e^{t\Delta/2} f)_{\mathbb{C}}, \tag{3.4}$$

where Δ is the Laplacian on $U(N)$, $e^{t\Delta/2}$ is the associated (forward) heat operator, and $(\cdot)_{\mathbb{C}}$ denotes the holomorphic extension of a sufficiently nice function from $U(N)$ to $\text{GL}(N; \mathbb{C})$. See also [22] for more information. The transform can easily be “boosted” to map matrix-valued functions on $U(N)$ to holomorphic matrix-valued functions on $\text{GL}(N; \mathbb{C})$ (by acting component-wise; i.e., via $\mathcal{G}_t^N \otimes \mathbf{1}_{M_N(\mathbb{C})}$).

A stochastic approach to the transform was developed by Gross and Malliavin in [19]; this approach played an important role in Biane’s paper [5]. See also [23, 24] for further development of the ideas in [19]. Let U_t^N and B_t^N be independent Brownian motions in $U(N)$ and $\text{GL}(N; \mathbb{C})$ (cf. (2.3)), and let f be a function on $U(N)$ that admits a holomorphic extension (also denoted f) to $\text{GL}(N; \mathbb{C})$. Then we have

$$\mathbb{E}[f(B_t^N U_t^N) | B_t^N] = (\mathcal{G}_t^N f)(B_t^N). \tag{3.5}$$

This result, by itself, is not deep. After all, in the finite-dimensional case, the conditional expectation can be computed as an expectation with respect to U_t^N , with B_t^N treated as a constant. Since U_t^N is

distributed as a heat kernel on $U(N)$, the left-hand side of (3.5) becomes a convolution of f with the heat kernel, giving the heat kernel in the definition (3.4) of the transform \mathcal{G}_t^N .

The crucial next step in [19] is to regard U_t^N and B_t^N as functionals of Brownian motions in the Lie algebra, by solving the relevant versions of the stochastic differential equation (2.2). Using this idea, Gross and Malliavin are able to deduce the properties of the *generalized* Segal–Bargmann from the previously known properties of the *classical* Segal–Bargmann transform for an infinite-dimensional linear space, namely the path space in the Lie algebra of $U(N)$. (We are glossing over certain technical distinctions; the preceding description is actually closer to [24, Theorem 18].) The expression (3.5) was the motivation for Biane’s formula (3.1) in the free case, and just as in [19], Biane was able to obtain properties of the transform \mathcal{G}_t from the corresponding linear case.

In [5], Biane conjectured, with an outline of a proof, that the free Hall transform \mathcal{G}_t can be realized using the large- N limit of \mathcal{G}_t^N . This conjecture was then verified independently by the authors and Driver in [13] and by Cébron in [10]; see also the expository paper [25].

The limiting process is as follows. Consider the transform \mathcal{G}_t^N on matrix-valued functions of the form $f(U)$, where f is a function on the unit circle and $f(U)$ is computed by the functional calculus. If, for example, f is the function $f(u) = u^2$ on the circle, then we can consider the associated matrix-valued function $f(U) = U^2$ on the unitary group $U(N)$. For any fixed N , the transformed function $\mathcal{G}_t^N(f)$ on $GL(N; \mathbb{C})$ will no longer be of functional-calculus type. Nevertheless, *in the large- N limit*, \mathcal{G}_t^N will map $f(U)$ to the functional-calculus function $(\mathcal{G}_t f)(Z)$, $Z \in GL(N; \mathbb{C})$.

Specifically, if p is a Laurent polynomial, then $\mathcal{G}_t p$ is also a Laurent polynomial, and (abusing notation slightly)

$$\mathcal{G}_t^N(p(U)) = (\mathcal{G}_t p)(Z) + O(1/N^2), \quad Z \in GL(N; \mathbb{C}),$$

where $O(1/N^2)$ denotes a term whose norm is bounded by a constant times $1/N^2$. See [13, Theorem 1.11] and [10, Theorem 4]. In particular, if $f(U) = U^2$, then in light of (3.2), we have

$$(\mathcal{G}_t^N f)(Z) = e^{-t}(Z^2 - tZ) + O(1/N^2), \quad Z \in GL(N; \mathbb{C}).$$

(See also Example 3.5 and the computations on p. 2592 of [13].)

4. AN OUTLINE OF THE PROOF OF THEOREM 1.1

As we pointed out in Proposition 2.5, the Brown measure of an operator a is supported on the spectrum of a . We strengthen this result, as follows. Given a noncommutative probability space (\mathcal{A}, τ) , we can construct the noncommutative L^2 space $L^2(\mathcal{A}, \tau)$, which is the completion of \mathcal{A} with respect to the noncommutative L^2 inner product, $\langle a, b \rangle = \tau(b^* a)$. It makes sense to multiply an element of the noncommutative L^2 space $L^2(\mathcal{A}, \tau)$ by an element of \mathcal{A} itself, and the result is again in $L^2(\mathcal{A}, \tau)$. We say that $a \in \mathcal{A}$ **has an inverse in L^2** if there exists some $b \in L^2(\mathcal{A}, \tau)$ such that $ab = ba = 1$.

Theorem 4.1. *Let (\mathcal{A}, τ) be a noncommutative probability space and let λ_0 be in \mathbb{C} . Suppose that $(a - \lambda)^2$ has an inverse—denoted $(a - \lambda)^{-2}$ —in $L^2(\mathcal{A}, \tau)$ for all λ in a neighborhood of λ_0 and that $\|(a - \lambda)^{-2}\|_{L^2(\mathcal{A}, \tau)}$ is bounded near λ_0 . Then λ_0 does not belong to the support of the Brown measure μ_a .*

Note that if $a - \lambda_0$ has a *bounded* inverse—that is, an inverse in \mathcal{A} —then $a - \lambda$ also has an inverse for all λ in a neighborhood of λ_0 , and $\|(a - \lambda)^{-1}\|_{\mathcal{A}}$ is bounded near λ_0 . In that case, we have

$$\|(a - \lambda)^{-2}\|_{L^2(\mathcal{A}, \tau)} \leq \|(a - \lambda)^{-2}\|_{\mathcal{A}} \leq \|(a - \lambda)^{-1}\|_{\mathcal{A}}^2,$$

which shows that $\|(a - \lambda)^{-2}\|_{L^2(\mathcal{A}, \tau)}$ is bounded near λ_0 . Thus, we can recover from Theorem 4.1 the result that the support of μ_a is contained in the spectrum of a . In general, however, Theorem 4.1 could apply even if $a - \lambda_0$ does not have a bounded inverse.

We now briefly indicate the proof of Theorem 4.1. Using the notation

$$a_\lambda = a - \lambda,$$

we make the following intuitive but non-rigorous estimates: for all $\varepsilon > 0$,

$$\tau[(a_\lambda^* a_\lambda + \varepsilon)^{-1} (a_\lambda a_\lambda^* + \varepsilon)^{-1}] \leq \tau[(a_\lambda^* a_\lambda)^{-1} (a_\lambda a_\lambda^*)^{-1}] = \tau[a_\lambda^{-2} (a_\lambda^{-2})^*] = \|(a - \lambda)^{-2}\|_{L^2(\mathcal{A}, \tau)}^2.$$

(The given estimate actually does hold, but its proof is quite subtle; the details are in Section 6.1.) If the hypotheses of the theorem hold, this last expression is bounded for λ near λ_0 . Corollary 2.6 then shows that λ_0 is not in the support of the Brown measure of a .

We now apply Theorem 4.1 to the case of interest to us, in which a is taken to be a free multiplicative Brownian motion b_t in a noncommutative probability space (\mathcal{B}, τ) . Recall that $L_{\text{hol}}^2(b_t, \tau)$ denotes the closure in $L^2(\mathcal{B}, \tau)$ of the space of Laurent polynomials in the element b_t .

Theorem 4.2. *For all $t > 0$, if $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_t$, then the element $(b_t - \lambda)^n$ has an inverse in $L_{\text{hol}}^2(b_t, \tau) \subset L^2(\mathcal{B}, \tau)$ for all $n = 1, 2, 3, \dots$, with local bounds on the L^2 norm of the inverse.*

When $t \neq 4$, the proof of this lemma draws on the transform \mathcal{G}_t in Theorem 3.1. We will show that the function $1/(z - \lambda)^n$ belongs to the space \mathcal{A}_t of holomorphic functions on Σ_t , at which point Theorem 3.1 tells us that there is a corresponding element $(b_t - \lambda)^{-n}$, which will be the inverse of $(b_t - \lambda)^n$. We demonstrate this key fact — that $1/(z - \lambda)^n$ belongs to the space $\mathcal{A}_t = \text{Image}(\mathcal{G}_t)$ — by explicitly constructing the preimage of $1/(z - \lambda)^n$ in $L^2(\partial\mathbb{D}, \nu_t)$. Specifically, using results from [5] or [13] about the generating function of the transform \mathcal{G}_t , we will show that

$$(\mathcal{G}_t)^{-1} \left(\frac{1}{(\cdot - \lambda)^n} \right) (\omega) = \frac{1}{(n-1)!} \left(\frac{\partial}{\partial \lambda} \right)^{n-1} \left[\frac{f_t(\lambda)}{\lambda} \frac{1}{\omega - f_t(\lambda)} \right], \quad \omega \in \text{supp}(\nu_t) \subset \partial\mathbb{D}.$$

Recall from Section 2.2 that f_t maps the complement of $\overline{\Sigma}_t$ to the complement of $\text{supp} \nu_t$. It follows that the function on the right-hand side is bounded—and therefore square integrable—on $\text{supp} \nu_t$, for all $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_t$. When $t = 4$, the proof is very similar, except that now we have to bypass the space \mathcal{A}_t and go directly from $L^2(\partial\mathbb{D}, \nu_t)$ to $L_{\text{hol}}^2(b_t, \tau)$.

The $n = 2$ case of Theorem 4.2 shows that Theorem 4.1 applies, and we conclude that the Brown measure of b_t is supported in $\overline{\Sigma}_t$.

5. THE TWO-PARAMETER CASE

In this section, we discuss the generalization of the process b_t and the Segal–Bargmann transform to the two-parameter setting $b_{s,t}$ of [13, 26, 27]; since $b_t = b_{t,t}$, we will prove the single-time theorems as stated as special cases of the general two-parameter framework. We mostly follow the notation in [26, Section 2.5].

5.1. Brownian motions. Fix positive real numbers s and t with $s > t/2$. Let $\{x_r\}_{r>0}$ and $\{y_r\}_{r>0}$ be freely independent free additive Brownian motions in a tracial von Neumann algebra (\mathcal{B}, τ) , with time-parameter denoted by r . Now define

$$w_{s,t}(r) = \sqrt{s - \frac{t}{2}} x_r + i \sqrt{\frac{t}{2}} y_r,$$

which we call a free elliptic (s, t) Brownian motion. The particular dependence of the coefficients on s and t is chosen to match the two-parameter Segal–Bargmann transform, which will be discussed below. Note: when $s = t$, $w_{t,t}(r) = \sqrt{\frac{t}{2}}(x_r + iy_r) = \sqrt{t}c_r$ (in terms of the free circular Brownian motion in Section 2.1).

We now define a “free multiplicative (s, t) Brownian motion” $b_{s,t}(r)$ as a solution to the free stochastic differential equation

$$db_{s,t}(r) = i b_{s,t}(r) dw_{s,t}(r) - \frac{1}{2}(s - t)b_{s,t}(r) dr \tag{5.1}$$

subject to the initial condition $b_{s,t}(0) = 1$. (The second term on the right-hand side of (5.1) is an Itô correction term that can be eliminated by writing the equation as a Stratonovich SDE.) We also use the notation

$$b_{s,t} = b_{s,t}(1). \quad (5.2)$$

When $s = t$, (5.1) becomes

$$db_{t,t}(r) = b_{t,t}(r) i\sqrt{t} dc_r.$$

Using the fact (from the usual Brownian scaling and rotational invariance) that the process $i\sqrt{t}c_r$ has the same law as the process c_{tr} , we see that $b_{t,t} = b_{t,t}(1)$ has the same noncommutative distribution as the free multiplicative Brownian motion b_t . On the other hand, the limiting case $(s, t) = (1, 0)$ gives a free unitary Brownian motion $b_{1,0}(r) = u_r$, cf. (2.4).

We can regard $b_{s,t}(r)$ as the large- N limit of a certain Brownian motion on the general linear group $GL(N; \mathbb{C})$ as follows. We define an inner product $\langle \cdot, \cdot \rangle_{s,t}$ on the Lie algebra $\mathfrak{gl}(N; \mathbb{C})$ by

$$\langle X_1 + iY_1, X_2 + iY_2 \rangle_{s,t} = \frac{N}{\sqrt{s-t/2}} \langle X_1, X_2 \rangle + \frac{N}{\sqrt{t/2}} \langle Y_1, Y_2 \rangle,$$

where X_1, X_2, Y_1 , and Y_2 are in the Lie algebra $\mathfrak{u}(N)$ of $U(N)$ and where the inner products on the right-hand side are the standard Hilbert–Schmidt inner product $\langle X, Y \rangle = \text{Trace}(Y^* X)$. We extend this inner product to a left-invariant Riemannian metric on $GL(N; \mathbb{C})$ and we then let

$$B_{s,t}^N(r)$$

be the Brownian motion with respect to this metric. In [27], the second author showed that the process $B_{s,t}^N(r)$ converges (in the sense of Definition 2.1) to the process $b_{s,t}(r)$, for all positive real numbers s, t , and r with $s > t/2$. (We are translating the results of [27] into the parametrizations used in [26].)

5.2. Segal–Bargmann transform. Meanwhile, the first author and Driver introduced in [12] a “two-parameter” Segal–Bargmann transform; see also [21]. In the case of the unitary group $U(N)$, the transform is a unitary map

$$\mathcal{G}_{s,t}^N : L^2(U(N), \rho_s) \rightarrow \mathcal{HL}^2(GL(N; \mathbb{C}), \gamma_{s,t}),$$

where ρ_s is the same heat kernel measure as in the one-parameter transform, but evaluated at time s , and where $\gamma_{s,t}$ is a heat kernel measure on $GL(N; \mathbb{C})$. Specifically, $\gamma_{s,t}$ is the distribution of the Brownian motion $B_{s,t}^N(r)$ at $r = 1$. The transform itself is defined precisely as in the one-parameter case:

$$\mathcal{G}_{s,t}^N f = (e^{t\Delta/2} f)_{\mathbb{C}};$$

only the inner products on the domain and range have changed. When $s = t$, the transform $\mathcal{G}_{s,t}^N$ coincides with the one-parameter transform \mathcal{G}_t^N .

In [13], the authors and Driver showed that the transform $\mathcal{G}_{s,t}^N$ has limiting properties as $N \rightarrow \infty$ similar to those of \mathcal{G}_t^N . Specifically, for each Laurent polynomial p in one variable, we showed that there is a unique Laurent polynomial $q_{s,t}$ in one variable such that (abusing notation slightly)

$$\mathcal{G}_{s,t}^N(p(U)) = q_{s,t}(Z) + O(1/N^2), \quad Z \in GL(N; \mathbb{C}).$$

As an example, if $p(u) = u^2$, then $q_{s,t}(z) = e^{-t}(z^2 - te^{-(s-t)/2}z)$, so that the transform of the matrix-valued function $F : U \mapsto U^2$ on $U(N)$ satisfies

$$(\mathcal{G}_{s,t}^N F)(Z) = e^{-t}(Z^2 - te^{-(s-t)/2}Z) + O(1/N^2), \quad Z \in GL(N; \mathbb{C}).$$

(See [13, p. 2592].)

In [26], Ho then constructed an integral transform $\mathcal{G}_{s,t}$ mapping $L^2(\partial\mathbb{D}, \nu_s)$ into a space of holomorphic functions on a certain domain $\Sigma_{s,t}$ in the plane. Ho’s transform $\mathcal{G}_{s,t}$ is uniquely determined by the fact that

$$\mathcal{G}_{s,t}(p) = q_{s,t}$$

for all Laurent polynomials p . Ho gave a description of $\mathcal{G}_{s,t}$ in terms of free probability similar to the description of Biane's transform \mathcal{G}_t given in Section 3.1, and he proved a unitary isomorphism theorem similar to Biane's result described in Theorem 3.1.

5.3. The domains $\Sigma_{s,t}$. Ho's domains have the property that f_{s-t} maps the complement of Σ_s to the complement of $\Sigma_{s,t}$. That is to say, $\Sigma_{s,t}$ is the complement of $f_{s-t}(\mathbb{C} \setminus \Sigma_s)$:

$$\Sigma_{s,t} = \mathbb{C} \setminus f_{s-t}(\mathbb{C} \setminus \Sigma_s). \quad (5.3)$$

(See Figure 5 along with [26, Figures 2 and 3].) Note that $\Sigma_{t,t}$ is the same as Σ_t . The topology of the domain $\Sigma_{s,t}$ is determined by s ; it is simply connected for $s \leq 4$ and doubly connected for $s > 4$.




FIGURE 5. The domain Σ_s with $s = 3$ (left) and the domain $\Sigma_{s,t}$ with $s = 3, t = 1$ (right). The map $f_{s-t} = f_2$ takes the complement of the domain on the left to the complement of the domain on the right

We need a two-parameter version of Proposition 2.3. To formulate the correct generalization, we first note that the function f_s satisfies

$$f_s(0) = 0; \quad f'_s(0) = e^{s/2} \neq 0.$$

Thus, f_s has a local inverse defined near zero, which we denote by χ_s . Recall from (2.6) that the support of the measure ν_s is a proper arc inside the unit circle for $s < 4$ and the whole unit circle for $s \geq 4$.

Proposition 5.1. *For all $s > 0$, χ_s can be extended uniquely to a holomorphic function on $\mathbb{C} \setminus \text{supp } \nu_s$ satisfying*

$$\chi_s(1/z) = 1/\chi_s(z). \quad (5.4)$$

Note that when $s \geq 4$, the support of ν_s is the entire unit circle, in which case $\mathbb{C} \setminus \text{supp } \nu_s$ is a disconnected set. For such values of s , the proposition is really asserting just that χ_s extends from a neighborhood of the origin to the open unit disk, at which point (5.4) serves to *define* $\chi_s(z)$ for $|z| > 1$.

For $s > t/2$, define a function $\chi_{s,t}$ by

$$\chi_{s,t} = f_{s-t} \circ \chi_s. \quad (5.5)$$

Since χ_s maps $\mathbb{C} \setminus \text{supp } \nu_s$ holomorphically to a region that does not include 1, we see that $\chi_{s,t}$ can also be defined holomorphically on $\mathbb{C} \setminus \text{supp } \nu_s$. Ho established the following result, generalizing Proposition 2.3. (See [26, Section 4.2], including the discussion following Remark 4.7.)

Proposition 5.2. *For all positive numbers s and t with $s > t/2$, define $\Sigma_{s,t}$ by (5.3). Then the function $\chi_{s,t}$ maps $\mathbb{C} \setminus \text{supp}(\nu_s)$ injectively onto the complement of $\overline{\Sigma_{s,t}}$. We denote the inverse function by $f_{s,t}$, so that*

$$f_{s,t}: \mathbb{C} \setminus \overline{\Sigma_{s,t}} \rightarrow \mathbb{C} \setminus \text{supp } \nu_s.$$

Note that, at least for sufficiently small z , we have $f_{s,t}(z) = f_s(\chi_{s-t}(z))$, by taking inverses in (5.5).

5.4. The main result. We are now ready to state the two-parameter version of Theorem 1.1.

Theorem 5.3. *Let $b_{s,t}$ be the free multiplicative Brownian motion with parameters (s, t) , as in (5.2). Then the support of the Brown measure $\mu_{b_{s,t}}$ is contained in $\overline{\Sigma}_{s,t}$.*

As in the one-parameter case, we expect that the limiting empirical eigenvalue distribution for $B_{s,t}^N := B_{s,t}^N(1)$ will also be supported in $\Sigma_{s,t}$. This is supported by simulations; see Figure 6. More generally, we expect that the empirical eigenvalue distribution of the Brownian motion $B_{s,t}^N$ in $\text{GL}(N; \mathbb{C})$ will converge almost surely to $\mu_{b_{s,t}}$ as $N \rightarrow \infty$. This question will be explored in a future paper.

Remark 5.4. While they do not determine the Brown measure, it is worth noting that the values of the holomorphic moments of $b_{s,t}$ were computed in [27]:

$$\tau(b_{s,t}^n) = \nu_n(s - t) \tag{5.6}$$

for all $n \in \mathbb{Z}$ and all $s > t/2 > 0$; here $\nu_n(r)$ are the moments of u_r , cf. (2.5). In particular, when $s = t$, since $\nu_n(0) = 1$ for all n , this recovers the fact that all holomorphic moments of b_t are 1.




FIGURE 6. Simulations of the Brownian motions $B_{s,t}^N$ for $N = 2000$ and $(s, t) = (3, 1)$ (left) and $(s, t) = (5, 3)$ (right), plotted against the domains $\Sigma_{s,t}$

6. PROOFS

In this final section, we present the complete proof of Theorem 5.3, which includes the main Theorem 1.1 as the special case $s = t$. We follow the outline of Section 4, and will therefore provide proofs of Theorem 4.1 and (a two-parameter generalization of) Theorem 4.2. We begin with the former.

6.1. A general result on the support of the Brown measure. In this subsection, we work with a general operator in a tracial von Neumann algebra (\mathcal{A}, τ) .

For $1 \leq p < \infty$, the noncommutative L^p norm on \mathcal{A} is

$$\|a\|_p = (\tau[|a|^p])^{1/p}.$$

The **noncommutative L^p space** $L^p(\mathcal{A}, \tau)$ is the completion of \mathcal{A} with respect to this norm; it can be concretely realized as a space of (largely unbounded, densely-defined) operators affiliated to \mathcal{A} .

For $a, b \in \mathcal{A}$ we have the inequality

$$\|ab\|_p \leq \|a\| \|b\|_p.$$

This shows that the operation of “left multiplication by a ” is bounded as an operator on \mathcal{A} with respect to the noncommutative L^p norm. Thus, by the bounded linear transformation theorem (e.g. [32, Theorem I.7]), left multiplication by a extends uniquely to a bounded linear map (with the same norm) of $L^p(\mathcal{A}, \tau)$ to itself. We may easily verify that

$$(ab)c = a(bc) \quad (6.1)$$

for $a, b \in \mathcal{A}$ and $c \in L^p(\mathcal{A}, \tau)$; this result holds when $c \in \mathcal{A}$ and then extends by continuity. Similar results hold for right multiplication by a .

Similarly, since

$$\|ab\|_1 \leq \|a\|_2 \|b\|_2, \quad (6.2)$$

for all $a, b \in \mathcal{A}$, the product map $(a, b) \mapsto ab$ can be extended by continuity first in a and then in b , giving a map from $L^2(\mathcal{A}, \tau) \times L^2(\mathcal{A}, \tau) \rightarrow L^1(\mathcal{A}, \tau)$ satisfying (6.2). We then observe a simple “associativity” property for the actions of \mathcal{A} on L^1 and L^2 :

$$a(bc) = (ab)c; \quad (bc)a = b(ca). \quad (6.3)$$

For any $1 \leq p < \infty$, we say that an element $a \in \mathcal{A}$ **has an inverse in L^p** if there is an element b of the noncommutative L^p space $L^p(\mathcal{A}, \tau)$ such that $ab = ba = 1$.

Definition 6.1. Let $a \in \mathcal{A}$, $n \in \mathbb{N}$, and $p \geq 1$. We say that λ_0 belongs to the L_n^p -**resolvent set** of a if $(a - \lambda)^n$ has an inverse, denoted $(a - \lambda)^{-n}$, in L^p for all λ in a neighborhood of λ_0 and $\|(a - \lambda)^{-n}\|_{L^p}$ is bounded near λ_0 . We say that λ_0 is in the L_n^p -**spectrum** of a if λ_0 is not in the L_n^p -resolvent set of a . We denote the L_n^p -spectrum of a by $\text{spec}_n^p(a)$.

Note that if $a - \lambda_0$ has a bounded inverse, then so does $a - \lambda$ for all λ sufficiently near λ_0 , and $\|(a - \lambda)^{-1}\|_{\mathcal{A}}$ —and therefore $\|(a - \lambda)^{-n}\|_{\mathcal{A}}$ —is automatically bounded near λ_0 . It follows that any point in the ordinary resolvent set of a is also in the L_n^p -resolvent set; equivalently,

$$\text{spec}_n^p(a) \subset \text{spec}(a),$$

where $\text{spec}(a)$ is the ordinary spectrum of a . Theorem 4.1 may then be restated as follows, strengthening the standard result that the Brown measure is supported on $\text{spec}(a)$.

Theorem 6.2. The support of the Brown measure μ_a of a is contained in $\text{spec}_2^2(a)$.

To prove this, the main result we need is the following.

Proposition 6.3. Suppose $a \in \mathcal{A}$ and a^2 has an inverse, denoted a^{-2} , in $L^2(\mathcal{A}, \tau)$. Then for all $\varepsilon > 0$, we have

$$\tau[(a^*a + \varepsilon)^{-1}(aa^* + \varepsilon)^{-1}] \leq \|a^{-2}\|_{L^2(\mathcal{A}, \tau)}^2. \quad (6.4)$$

Theorem 6.2 follows immediately from Proposition 6.3 together with Corollary 2.6. To prove the proposition, we need the following lemmas.

Lemma 6.4. For $a, b \in \mathcal{A}$, if a is invertible in \mathcal{A} and b is invertible in L^p , then ab and ba are invertible in L^p with inverses $b^{-1}a^{-1}$ and $a^{-1}b^{-1}$, respectively. If a and b are invertible in L^2 then ab is invertible in L^1 with inverse $b^{-1}a^{-1}$. Finally, if a is invertible in L^p then a^* is invertible in L^p with inverse $(a^{-1})^*$.

Proof. For $a, b \in \mathcal{A}$ with b invertible in L^p , we have

$$\begin{aligned} b^{-1}a^{-1}(ab) &= ((b^{-1}a^{-1})a)b = ((b^{-1}(a^{-1}a))b \\ &= b^{-1}b = 1, \end{aligned}$$

where we have used (6.1) twice, and similarly for the product in the other order. A similar argument, using both (6.1) and (6.3), verifies the second claim. Finally, the identity $(ab)^* = b^*a^*$, which holds initially for $a, b \in \mathcal{A}$, extends by continuity to the case $a \in \mathcal{A}$, $b \in L^p(\mathcal{A}, \tau)$. Thus, if a is invertible in L^p , then $a^*(a^{-1})^* = (a^{-1}a)^* = 1$ and similarly for the product in the reverse order. \square

Lemma 6.5. *Let x be a non-negative element of \mathcal{A} and suppose x has an inverse in L^1 . Then*

$$\lim_{\varepsilon \rightarrow 0^+} \tau[(x + \varepsilon)^{-1}] = \tau[x^{-1}].$$

Proof. We begin by noting that

$$\begin{aligned} x^{-1} - (x + \varepsilon)^{-1} &= x^{-1}((x + \varepsilon) - x)(x + \varepsilon)^{-1} \\ &= \varepsilon x^{-1}(x + \varepsilon)^{-1}. \end{aligned}$$

Now, $\|(x + \varepsilon)^{-1}\|$ is at most $1/\varepsilon$, by the equality of norm and spectral radius for self-adjoint elements. Thus,

$$\|x^{-1} - (x + \varepsilon)^{-1}\|_1 \leq \varepsilon \|(x + \varepsilon)^{-1}\| \|x^{-1}\|_1 \leq \|x^{-1}\|_1.$$

It follows that $\|(x + \varepsilon)^{-1}\|_1 \leq 2\|x^{-1}\|_1$ for all $\varepsilon > 0$.

Let E^x be the projection-valued measure associated to x by the spectral theorem. Then

$$\|(x + \varepsilon)^{-1}\|_1 = \int_0^\infty \frac{1}{\lambda + \varepsilon} \tau[E^x(d\lambda)] \leq 2\|x^{-1}\|_1.$$

Thus, by monotone convergence, we have

$$\int_0^\infty \frac{1}{\lambda} \tau[E^x(d\lambda)] \leq 2\|x^{-1}\|_1 < \infty.$$

Once this is established, we note that for $\lambda > 0$,

$$\left| \frac{1}{\lambda} - \frac{1}{\lambda + \varepsilon} \right| = \frac{\varepsilon}{\lambda(\lambda + \varepsilon)} \leq \frac{1}{\lambda}.$$

Thus, by dominated convergence, $1/(\lambda + \varepsilon)$ converges in $L^1((0, \infty), \tau \circ E^x)$ to $1/\lambda$ as $\varepsilon \rightarrow 0$. It follows that for any sequence $\{\varepsilon_n\}$ tending to zero, the operators $(x + \varepsilon_n)^{-1}$ form a Cauchy sequence with respect to the noncommutative L^1 norm. Since, by dominated convergence, the functions $\lambda/(\lambda + \varepsilon_n)$ converge to 1 in $L^1((0, \infty), \tau \circ E^x)$, we can easily see that the limit of $(x + \varepsilon_n)^{-1}$ in the Banach space $L^1(\mathcal{A}, \tau)$ is the inverse in L^1 of x .

We have shown, then, that the (unique) inverse in L^1 of x is the limit in L^1 of $(x + \varepsilon_n)^{-1}$. Applying τ to this result gives the claim, along any sequence $\{\varepsilon_n\}$ tending to zero. \square

Lemma 6.6. *Let $x, y \geq 0$ be positive semidefinite operators in \mathcal{A} , and suppose that they are invertible in $L^1(\mathcal{A}, \tau)$. If $x \leq y$ (i.e., $x - y$ is positive semidefinite) then $\tau(y^{-1}) \leq \tau(x^{-1})$.*

Proof. For all $\varepsilon > 0$, the elements $x + \varepsilon$ and $y + \varepsilon$ are invertible in \mathcal{A} and $x + \varepsilon \geq y + \varepsilon$. Thus, by the (reverse) operator monotonicity of the inverse [3, Prop. V.1.6], we have

$$\tau((y + \varepsilon)^{-1}) \geq \tau((x + \varepsilon)^{-1}). \quad (6.5)$$

By Lemma 6.5, the claimed result then follows. \square

Lemma 6.7. *Let $a \in \mathcal{A}$ and suppose that a^2 has an inverse in $L^2(\mathcal{A}, \tau)$. Then a and a^* have inverses in L^2 ; and hence a^*a and aa^* have inverses in $L^1(\mathcal{A}, \tau)$.*

Proof. Let $b = a^{-2}$ denote the $L^2(\mathcal{A}, \tau)$ inverse of a^2 . First, note that $a(ab) = a^2b = 1$ and $(ba)a = ba^2 = 1$. Since ab and ba are in $L^2(\mathcal{A}, \tau)$, it follows that a is both left and right invertible in $L^2(\mathcal{A}, \tau)$. Hence, a is invertible in $L^2(\mathcal{A}, \tau)$, by the usual argument and (6.1). Taking adjoints shows that a^* is also invertible in $L^2(\mathcal{A}, \tau)$. Thus $aa^*[(a^*)^{-1}a^{-1}] = 1 = [(a^*)^{-1}a^{-1}]aa^*$. Since $(a^*)^{-1}$ and a^{-1} are in $L^2(\mathcal{A}, \tau)$, their product is in $L^1(\mathcal{A}, \tau)$ by Hölder's inequality; this shows aa^* is invertible in $L^1(\mathcal{A}, \tau)$. An analogous argument holds for a^*a . \square

We are now ready for the proof of Proposition 6.3.

Proof of Proposition 6.3. We begin by arguing formally and then fill in the details. We start by noting that

$$(a^*a + \varepsilon)^{1/2}(aa^* + \varepsilon)(a^*a + \varepsilon)^{1/2} - (a^*a + \varepsilon)^{1/2}(aa^*)(a^*a + \varepsilon)^{1/2} = \varepsilon(a^*a + \varepsilon) \geq 0.$$

Thus, by Lemma 6.4 and the cyclic property of the trace, we have

$$\begin{aligned} \tau[(a^*a + \varepsilon)^{-1}(aa^* + \varepsilon)^{-1}] &= \tau[(a^*a + \varepsilon)^{-1/2}(aa^* + \varepsilon)^{-1}(a^*a + \varepsilon)^{-1/2}] \\ &\leq \tau[(a^*a + \varepsilon)^{-1/2}(aa^*)^{-1}(a^*a + \varepsilon)^{-1/2}] \\ &= \tau[(aa^*)^{-1}(a^*a + \varepsilon)^{-1}]. \end{aligned} \quad (6.6)$$

We then use the same argument again. We note that

$$a^*(a^*a + \varepsilon)a - a^*(a^*a)a = \varepsilon a^*a \geq 0, \quad (6.7)$$

from which we obtain

$$\begin{aligned} \tau[(aa^*)^{-1}(a^*a + \varepsilon)^{-1}] &= \tau[a^{-1}(a^*a + \varepsilon)^{-1}(a^*)^{-1}] \\ &= \tau[(a^*(a^*a + \varepsilon)a)^{-1}] \\ &\leq \tau[((a^*)^2a^2)^{-1}] \\ &= \tau[a^{-2}(a^{-2})^*]. \\ &= \|a^{-2}\|_{L^2(\mathcal{A}, \tau)}^2 \end{aligned} \quad (6.8)$$

Combining (6.6) and (6.8) gives the desired inequality.

To make the argument rigorous, we need to make sure that all the relevant inverses exist in $L^1(\mathcal{A}, \tau)$, so that Lemma 6.4 is applicable. The operator $(a^*a + \varepsilon)^{1/2}(aa^* + \varepsilon)(a^*a + \varepsilon)^{1/2}$ is invertible in \mathcal{A} and thus in $L^1(\mathcal{A}, \tau)$. On the other hand, by assumption a^2 is invertible in $L^2(\mathcal{A}, \tau)$. It then follows from Lemma 6.5 that aa^* is invertible in $L^1(\mathcal{A}, \tau)$, so that $(a^*a + \varepsilon)^{1/2}(aa^*)(a^*a + \varepsilon)^{1/2}$ is also invertible in $L^1(\mathcal{A}, \tau)$, by Lemma 6.4. Thus, Lemma 6.4 is applicable in (6.6).

We now consider the two terms on the left-hand side of (6.7). By Lemma 6.5, both a and a^* are invertible in L^2 ; it then follows from Lemma 6.4 that $a^*(a^*a + \varepsilon)$ is invertible in L^2 and that $a^*(a^*a + \varepsilon)a$ is invertible in L^1 . Meanwhile, a^2 is, by assumption, invertible in L^2 ; it then follows from Lemma 6.4 that $(a^*)^2$ is invertible in L^2 . Thus, using Lemma 6.4 one last time, we conclude that $(a^*)^2a^2$ is invertible in L^1 . Thus, Lemma 6.5 is also applicable in (6.8). \square

6.2. Computing the L^2_2 spectrum. We consider the free multiplicative (s, t) Brownian motion $b_{s,t}$ defined in (5.2) as an element of a noncommutative probability space (\mathcal{B}, τ) . Recall that when $s = t$, the operator $b_{s,t}$ has the same noncommutative distribution as the ordinary free multiplicative Brownian motion b_t described in Section 2.1. Our goal is to prove a two-parameter version of Theorem 4.2 from Section 4, by constructing an L^2 inverse to $(b_{s,t} - \lambda)^n$ for λ in the complement of $\overline{\Sigma}_{s,t}$, with local bounds on the norm of the inverse. This will, in particular, show that

$$\text{spec}_n^2(b_{s,t}) \subset \overline{\Sigma}_{s,t} \quad (6.9)$$

for all n . (Recall Definition 6.1.) If we specialize (6.9) to the case $n = 2$, Theorem 6.2 will then tell us that the support of the Brown measure of $b_{s,t}$ is contained in $\overline{\Sigma}_{s,t}$.

Our tool is the two-parameter ‘‘free Hall transform’’ $\mathcal{G}_{s,t}$ (cf. Section 5), which includes the one-parameter transform $\mathcal{G}_t = \mathcal{G}_{t,t}$ (cf. Section 3) as a special case. To avoid technical issues for the case $s = 4$, we introduce a variant of the transform $\mathcal{G}_{s,t}$, denoted $\mathcal{S}_{s,t}$. Define

$$L^2_{\text{hol}}(b_{s,t}, \tau)$$

to be the closure in the noncommutative L^2 norm of the space of elements of the form $p(b_{s,t})$, where p is a Laurent polynomial in one variable.

If \mathcal{P} denotes the space of Laurent polynomials in one variable, then [13, Theorem 1.13] shows that $\mathcal{G}_{s,t}$ maps \mathcal{P} bijectively onto \mathcal{P} . We then define $\mathcal{S}_{s,t}$ initially on \mathcal{P} by evaluating each polynomial $\mathcal{G}_{s,t}(p)$ on the element $b_{s,t}$:

$$\mathcal{S}_{s,t}(p) = \mathcal{G}_{s,t}(p)(b_{s,t}) \in L^2_{\text{hol}}(b_{s,t}, \tau).$$

By [26, Theorem 5.7], $\mathcal{S}_{s,t}$ maps $\mathcal{P} \subset L^2(\partial\mathbb{D}, \nu_s)$ isometrically into $L^2_{\text{hol}}(b_{s,t}, \tau)$. (Although some parts of the just-cited theorem implicitly assume that $s \neq 4$, Part 2 of the theorem does not depend on this assumption.) Furthermore, since $\mathcal{G}_{s,t}$ maps *onto* \mathcal{P} , the image of $\mathcal{S}_{s,t}$ is dense in $L^2_{\text{hol}}(b_{s,t}, \tau)$. Thus, $\mathcal{S}_{s,t}$ extends to a unitary map

$$\mathcal{S}_{s,t} : L^2(\partial\mathbb{D}, \nu_s) \rightarrow L^2_{\text{hol}}(b_{s,t}, \tau).$$

For $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$, we will then construct an inverse $(b_{s,t} - \lambda)^{-n}$ to $(b_{s,t} - \lambda)^n$ in $L^2_{\text{hol}}(b_{s,t}, \tau)$ by constructing the preimage of $(b_{s,t} - \lambda)^{-n}$ under $\mathcal{S}_{s,t}$. Note that in Section 4, we outlined the proof in the case $s = t$, with $t \neq 4$. In that case, [5, Lemma 17] allows us to identify $L^2_{\text{hol}}(\mathcal{B}, \tau)$ with the space \mathcal{A}_t , in which case, it is harmless to work with $\mathcal{G}_{s,t}$ instead of $\mathcal{S}_{s,t}$.

Recall the definition of the function $f_{s,t}$ in Proposition 5.2.

Theorem 6.8. *Define a function $r_\lambda^{s,t} \in L^2(\partial\mathbb{D}, \nu_s)$ by the formula*

$$r_\lambda^{s,t}(\omega) = \frac{f_{s,t}(\lambda)}{\lambda} \left(\frac{1}{\omega - f_{s,t}(\lambda)} \right), \quad \omega \in \text{supp}(\nu_s), \quad (6.10)$$

for all $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$. Then for all such λ , we have

$$\mathcal{S}_{s,t}(r_\lambda^{s,t}) = (b_{s,t} - \lambda)^{-1}.$$

That is to say, $\mathcal{S}_{s,t}(r_\lambda^{s,t})$ is an inverse in L^2 of $(b_{s,t} - \lambda)$.

Recall from Proposition 5.2 that $f_{s,t}(\lambda)$ is outside the support of ν_s for all λ in the complement of $\overline{\Sigma}_{s,t}$. Thus, $r_\lambda^{s,t}(\omega)$ is bounded and is therefore a ν_s -square integrable function of ω for all $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$.

When $\lambda = 0$, we interpret $r_\lambda^{s,t}$ to be the limit of the right-hand side of (6.10) as λ approaches zero, which is easily computed to be $r_0^{s,t}(\omega) = e^{t/2}\omega^{-1}$.

Proof. For the case $\lambda = 0$, we compute that $\mathcal{G}_{s,t}(\omega^{-1}) = e^{-t/2}z^{-1}$, as may be verified from the behavior of $\mathcal{G}_{s,t}$ with respect to inversion [13, Eq. (5.2)] and the recursive formula in [13, Proposition 5.2]. Thus, by definition, we have $\mathcal{S}_{s,t}(\omega^{-1}) = e^{-t/2}b_{s,t}^{-1}$ so that $\mathcal{S}_{s,t}(e^{t/2}\omega^{-1}) = b_{s,t}^{-1}$, which is the $\lambda = 0$ case of the theorem. The subsequent calculations assume $\lambda \neq 0$.

We start by considering large values of λ . For $|\lambda| > \|b_{s,t}\|$, the element $b_{s,t} - \lambda$ has a *bounded* inverse, which may be computed as a power series:

$$(b_{s,t} - \lambda)^{-1} = -\frac{1}{\lambda}(1 - b_{s,t}/\lambda)^{-1} = -\frac{1}{\lambda} \sum_{k=0}^{\infty} \lambda^{-k} (b_{s,t})^k,$$

with the series converging in the operator norm and thus also in the noncommutative L^2 norm. Applying $\mathcal{S}_{s,t}^{-1}$ term by term gives

$$\mathcal{S}_{s,t}^{-1}((b_{s,t} - \lambda)^{-1})(\omega) = -\frac{1}{\lambda} \sum_{k=0}^{\infty} \lambda^{-k} p_k^{s,t}(\omega) \quad (6.11)$$

$$= -\frac{1}{\lambda} [1 + \Pi(s, t, \omega, 1/\lambda)], \quad (6.12)$$

where $p_k^{s,t}$ is the unique polynomial such that

$$\mathcal{G}_{s,t}(p_k^{s,t})(z) = z^k, \quad k \in \mathbb{Z},$$

and where Π is the generating function in [13, Theorem 1.17].

Meanwhile, according to [13, Eq. (1.21)], we have the following implicit formula for Π :

$$\Pi(s, t, \omega, f_{s-t}(z)) = \frac{1}{1 - \omega f_s(z)} - 1.$$

(This result extends a formula of Biane in the $s = t$ case.) Then, at least for sufficiently small z , we may replace z by $\chi_{s-t}(z)$, where $\chi_{s,t}$ is the inverse function to f_{s-t} . This substitution gives

$$\Pi(s, t, \omega, z) = \frac{1}{1 - \omega f_s(\chi_{s-t}(z))} - 1. \quad (6.13)$$

Plugging this expression into (6.12) gives

$$r_\lambda^{s,t}(\omega) = -\frac{1}{\lambda} \frac{1}{1 - \omega f_s(\chi_{s-t}(1/\lambda))},$$

for λ sufficiently large, which simplifies—using (5.4) and the identity $f_s(1/z) = 1/f_s(z)$ —to the expression in (6.10).

Similarly, for $0 < |\lambda| < 1/\|b_{s,t}^{-1}\|$, we use the series expansion

$$\begin{aligned} (b_{s,t} - \lambda)^{-1} &= b_{s,t}^{-1}(1 - \lambda b_{s,t}^{-1})^{-1} \\ &= b_{s,t}^{-1} \sum_{k=0}^{\infty} \lambda^k b_{s,t}^{-k} \\ &= \frac{1}{\lambda} \sum_{k=1}^{\infty} \lambda^k b_{s,t}^{-k}. \end{aligned}$$

Since $p_{-k}^{s,t}(\omega) = p_k^{s,t}(\omega^{-1})$ (see [13, Eq. (5.2)]), we may apply $\mathcal{S}_{s,t}^{-1}$ term by term to obtain

$$\begin{aligned} \mathcal{S}_{s,t}^{-1}((b_{s,t} - \lambda)^{-1})(\omega) &= \frac{1}{\lambda} \sum_{k=1}^{\infty} \lambda^k p_k^{s,t}(\omega^{-1}) \\ &= \frac{1}{\lambda} \Pi(s, t, \omega^{-1}, \lambda). \end{aligned}$$

Using the formula (6.13) for Π (for sufficiently small λ) and simplifying gives the same expression as for the large- λ case.

Finally, for general $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$ we use an analytic continuation argument. The complement of the closure of $\Sigma_{s,t}$ has at most two connected components, the unbounded component and, for $s \geq 4$, a bounded component containing 0. Now, for all $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$, the function $r_\lambda^{s,t}$ in (6.10) is a well-defined element of $L^2(\partial\mathbb{D}, \nu_s)$, because (by Proposition 5.2) $f_{s,t}$ maps $\mathbb{C} \setminus \overline{\Sigma}_{s,t}$ into $\mathbb{C} \setminus \text{supp } \nu_s$. Furthermore, it is evident from (6.10) that $r_\lambda^{s,t}$ is a weakly holomorphic function of $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$, with values in $L^2(\partial\mathbb{D}, \nu_s)$, meaning that $\lambda \mapsto \phi(r_\lambda^{s,t})$ is a holomorphic for each bounded linear functional on $L^2(\partial\mathbb{D}, \nu_s)$.

Thus, since $\mathcal{S}_{s,t}$ is unitary (hence bounded), $\mathcal{S}_{s,t}(r_\lambda^{s,t})$ is a weakly holomorphic function of $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$, with values in $L_{\text{hol}}^2(\mathcal{B}, \tau)$. It is then easy to see that

$$(b_{s,t} - \lambda) \mathcal{S}_{s,t}(r_\lambda^{s,t}) = b_{s,t} \mathcal{S}_{s,t}(r_\lambda^{s,t}) - \lambda \mathcal{S}_{s,t}(r_\lambda^{s,t}) \quad (6.14)$$

is also weakly holomorphic. After all, applying a bounded linear functional ϕ gives

$$\phi(b_{s,t} \mathcal{S}_{s,t}(r_\lambda^{s,t})) - \lambda \phi(\mathcal{S}_{s,t}(r_\lambda^{s,t})). \quad (6.15)$$

Since multiplication by $b_{s,t}$ is a bounded linear map on $L_{\text{hol}}^2(\mathcal{B}, \tau)$, the linear functional $\phi(b_{s,t} \cdot)$ is bounded, so both terms in (6.15) are holomorphic in λ .

Meanwhile, we have shown that (6.14) is equal to 1 on a nonempty, open subset of each connected component of $\mathbb{C} \setminus \overline{\Sigma}_{s,t}$. Since, also, (6.14) is weakly holomorphic, it must be equal to 1 on all of $\mathbb{C} \setminus \overline{\Sigma}_{s,t}$. \square

We emphasize that, although for very large and small λ the standard power-series argument gives an inverse of $b_{s,t} - \lambda$ in the algebra \mathcal{B} , the analytic continuation takes place in $L^2_{\text{hol}}(\mathcal{B}, \tau)$, which is the range of the transform $\mathcal{S}_{s,t}$. Thus, for general $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$, we are guaranteed that $b_{s,t} - \lambda$ has an inverse in L^2 , but not necessarily in \mathcal{B} .

Corollary 6.9. *For all $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$ and all positive integers n , the operator $(b_{s,t} - \lambda)^n$ has an inverse $(b_{s,t} - \lambda)^{-n}$ in $L^2(\mathcal{B}, \tau)$. Specifically,*

$$(b_{s,t} - \lambda)^{-n} = \mathcal{S}_{s,t} \left(\frac{1}{(n-1)!} \left(\frac{\partial}{\partial \lambda} \right)^{n-1} r_{\lambda}^{s,t} \right), \quad (6.16)$$

where $r_{\lambda}^{s,t}$ is as in (6.10). Furthermore, $\|(b_{s,t} - \lambda)^{-n}\|_{L^2(\mathcal{B}, \tau)}$ is locally bounded on $\mathbb{C} \setminus \overline{\Sigma}_{s,t}$.

Proof. We let

$$r_{\lambda}^{s,t,n}(\omega) = \frac{1}{(n-1)!} \left(\frac{\partial}{\partial \lambda} \right)^{n-1} r_{\lambda}^{s,t}(\omega).$$

If we inductively compute the derivatives in the definition of $r_{\lambda}^{s,t,n}(\omega)$, we will find that the result is polynomial in $1/(\omega - f_{s,t}(\lambda))$, with coefficients that are holomorphic functions of $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$. Thus, for each n , the quantity $r_{\lambda}^{s,t,n}(\omega)$ is jointly continuous as a function of $\omega \in \text{supp}(\nu_t)$ and $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$. Thus, $\|r_{\lambda}^{s,t,n}\|_{L^2(\partial\mathbb{D}, \nu_s)}$ is finite and depends continuously on λ . Thus, once (6.16) is verified, the local bounds on the norm of $(b_{s,t} - \lambda)^{-n}$ will follow from the unitarity of $\mathcal{S}_{s,t}$.

We establish (6.16) by induction on n , the $n = 1$ case being the content of Theorem 6.8. Assume, then, the result for a fixed n and recall that $r_{\lambda}^{s,t,n}(\omega)$ is a polynomial in $1/(\omega - f_{s,t}(\lambda))$, with coefficients that are holomorphic functions of λ . It is then an elementary matter to see that for each fixed $\lambda \in \mathbb{C} \setminus \overline{\Sigma}_{s,t}$, the limit

$$\lim_{h \rightarrow 0} \frac{r_{\lambda+h}^{s,t,n}(\omega) - r_{\lambda}^{s,t,n}(\omega)}{h}$$

exists as a uniform limit on $\text{supp} \nu_s$, and thus also in $L^2(\partial\mathbb{D}, \nu_s)$. Applying $\mathcal{S}_{s,t}$ and using our induction hypothesis gives

$$\mathcal{S}_{s,t} \left(\frac{\partial}{\partial \lambda} r_{\lambda}^{s,t,n}(\omega) \right) = \lim_{h \rightarrow 0} \frac{1}{h} [(b_{s,t} - (\lambda + h))^{-n} - (b_{s,t} - \lambda)^{-n}], \quad (6.17)$$

where the limit is in $L^2_{\text{hol}}(b_{s,t}, \tau)$.

We now multiply both sides of (6.17) by $(b_{s,t} - \lambda)^{n+1}$, which we write as $(b_{s,t} - \lambda)(b_{s,t} - \lambda)^n$. Since multiplication by an element of \mathcal{B} is a continuous linear map on $L^2(\mathcal{B}, \tau)$, we obtain

$$\begin{aligned} & (b_{s,t} - \lambda)^{n+1} \mathcal{S}_{s,t} \left(\frac{\partial}{\partial \lambda} r_{\lambda}^{s,t,n}(\omega) \right) \\ &= \lim_{h \rightarrow 0} (b_{s,t} - \lambda) \frac{1}{h} [(b_{s,t} - \lambda)^n (b_{s,t} - (\lambda + h))^{-n} - 1]. \end{aligned} \quad (6.18)$$

In the first term on the right-hand side of (6.18), we write

$$(b_{s,t} - \lambda)^n = (b_{s,t} - (\lambda + h) + h)^n = \sum_{k=0}^n \binom{n}{k} (b_{s,t} - (\lambda + h))^{n-k} h^k. \quad (6.19)$$

Upon substituting (6.19) into (6.18), the $k = 0$ term cancels with the existing term of -1 , while the $k = 1$ term gives

$$(b_{s,t} - \lambda) \frac{1}{h} \cdot nh(b_{s,t} - (\lambda + h))^{-1} = n(b_{s,t} - \lambda)(b_{s,t} - (\lambda + h))^{-1}. \quad (6.20)$$

If we again write $(b_{s,t} - \lambda) = (b_{s,t} - (\lambda + h) + h)$, we see that (6.20) tends to $n \cdot 1$ as $h \rightarrow 0$. Finally, all terms with $k \geq 2$ will vanish in the limit, leaving us with

$$(b_{s,t} - \lambda)^{n+1} \mathcal{S}_{s,t} \left(\frac{\partial}{\partial \lambda} r_{\lambda}^{s,t,n}(\omega) \right) = n \cdot 1.$$

Thus,

$$\mathcal{S}_{s,t} \left(\frac{1}{n} \frac{\partial}{\partial \lambda} r_{\lambda}^{s,t,n}(\omega) \right) = (b_{s,t} - \lambda)^{-(n+1)},$$

which is just the level- $(n + 1)$ case of the corollary. \square

Acknowledgments. We would like to thank Adrian Ioana and Ching Wei Ho for useful conversations.

REFERENCES

- [1] Z. D. Bai, Circular law, *Ann. Probab.* **25** (1997), 494–529.
- [2] H. Bercovici and D. Voiculescu, Lévy–Hinčin type theorems for multiplicative and additive free convolution, *Pacific J. Math.* **153** (1992), 217–248.
- [3] R. Bhatia, Matrix analysis. Graduate Texts in Mathematics, 169. Springer-Verlag, New York, 1997.
- [4] P. Biane, “Free Brownian motion, free stochastic calculus and random matrices.” In Free Probability Theory (Waterloo, ON, 1995), 1–19. Fields Institute Communications 12. Providence, RI: American Mathematical Society, 1997.
- [5] P. Biane, Segal–Bargmann transform, functional calculus on matrix spaces and the theory of semi-circular and circular systems, *J. Funct. Anal.*, **144** (1997), 232–286.
- [6] P. Biane, Processes with free increments, *Math. Z.* **227** (1998), 143–174.
- [7] P. Biane and R. Speicher, Stochastic calculus with respect to free Brownian motion and analysis on Wigner space, *Probab. Theory Related Fields* **112** (1998), 373–409.
- [8] P. Biane and R. Speicher, Free diffusions, free entropy and free Fisher information, *Ann. Inst. H. Poincaré Probab. Statist.* **37** (2001), 581–606.
- [9] Brown, L. G. Lidskiĭ’s theorem in the type II case. In Geometric methods in operator algebras (Kyoto, 1983), 1–35, Pitman Res. Notes Math. Ser., 123, Longman Sci. Tech., Harlow, 1986.
- [10] G. Cébron, Free convolution operators and free Hall transform, *J. Funct. Anal.* **265** (2013), 2645–2708.
- [11] B. Collins and T. Kemp, Liberation of projections, *J. Funct. Anal.* **266** (2014), 1988–2052.
- [12] B. K. Driver and B. C. Hall, Yang–Mills theory and the Segal–Bargmann transform, *Comm. Math. Phys.* **201** (1999), 249–290.
- [13] B. K. Driver, B. C. Hall, and T. Kemp, The large- N limit of the Segal–Bargmann transform on $U(N)$, *J. Funct. Anal.* **265** (2013), 2585–2644.
- [14] B. Fuglede and R. V. Kadison, On determinants and a property of the trace in finite factors, *Proc. Nat. Acad. Sci. U. S. A.* **37** (1951), 425–431.
- [15] B. Fuglede and R. V. Kadison, Determinant theory in finite factors, *Ann. of Math. (2)* **55** (1952), 520–530.
- [16] J. Ginibre, Statistical ensembles of complex, quaternion, and real matrices, *J. Math. Phys.* **6** (1965), 440–449.
- [17] V. L. Girko, The circular law. (Russian) *Teor. Veroyatnost. i Primenen.* **29** (1984), 669–679.
- [18] F. Götze and A. Tikhomirov, The circular law for random matrices, *Ann. Probab.* **38** (2010), 1444–1491.
- [19] L. Gross and P. Malliavin, Hall’s transform and the Segal–Bargmann map. In Itô’s stochastic calculus and probability theory (N. Ikeda, S. Watanabe, M. Fukushima and H. Kunita, Eds.), 73–116, Springer, 1996.
- [20] B. C. Hall, The Segal–Bargmann “coherent state” transform for compact Lie groups. *J. Funct. Anal.* **122** (1994), 103–151.
- [21] B. C. Hall, A new form of the Segal–Bargmann transform for Lie groups of compact type, *Canad. J. Math.* **51** (1999), 816–834.
- [22] B. C. Hall, Harmonic analysis with respect to heat kernel measure, *Bull. Amer. Math. Soc. (N.S.)* **38** (2001), 43–78.
- [23] B. C. Hall, The Segal–Bargmann transform and the Gross ergodicity theorem. In Finite and infinite dimensional analysis in honor of Leonard Gross (New Orleans, LA, 2001), 99–116, Contemp. Math., 317, Amer. Math. Soc., Providence, RI, 2003.
- [24] B. C. Hall and A. N. Sengupta, The Segal–Bargmann transform for path-groups, *J. Funct. Anal.* **152** (1998), 220–254.
- [25] B. C. Hall, The Segal–Bargmann transform for unitary groups in the large- N limit, preprint arXiv:1308.0615 [math.RT].

- [26] C.-W. Ho, The two-parameter free unitary Segal–Bargmann transform and its Biane–Gross–Malliavin identification, *J. Funct. Anal.* **271** (2016), 3765–3817.
- [27] T. Kemp, The large- N limits of Brownian motions on $GL(N)$, *Int. Math. Res. Not.*, (2016), 4012–4057.
- [28] T. Kemp, I. Nourdin, G. Peccati, and R. Speicher, Wigner chaos and the fourth moment. *Ann. Prob.* **40** (2012), 1577–1635.
- [29] H. P. McKean, Stochastic Integrals. Probability and Mathematical Statistics 5. New York: Academic Press, 1969
- [30] J. A. Mingo and R. Speicher, Free probability and random matrices. Fields Institute Monographs, 35. Springer, New York; Fields Institute for Research in Mathematical Sciences, Toronto, ON, 2017.
- [31] A. Nica and R. Speicher, Lectures on the combinatorics of free probability. London Mathematical Society Lecture Note Series, 335. Cambridge University Press, Cambridge, 2006. xvi+417 pp.
- [32] M. Reed and B. Simon, Methods of modern mathematical physics. I. Functional analysis. Second edition. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1980. xv+400 pp.
- [33] I. M. Singer, On the master field in two dimensions. *In* Functional analysis on the eve of the 21st century, Vol. 1 (New Brunswick, NJ, 1993), volume 131 of *Progr. Math.*, 263–281. Birkhäuser Boston, Boston, MA, 1995.
- [34] T. Tao and V. Vu, Random matrices: universality of ESDs and the circular law. With an appendix by Manjunath Krishnapur. *Ann. Probab.* **38** (2010), 2023–2065.
- [35] D. V. Voiculescu, Limit laws for random matrices and free products. *Invent. Math.* **104** (1991), 201–220.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556
E-mail address: bhall@nd.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SAN DIEGO, LA JOLLA, CA 92093-0112
E-mail address: tkemp@math.ucsd.edu