

Mechanisms and constraints on yeast MAPK signaling specificity

Bo Hu, Wouter-Jan Rappel, and Herbert Levine ¹,
Center for Theoretical Biological Physics and Department of Physics,
University of California at San Diego, La Jolla, California 92093-0374

¹Corresponding author. Address: Center for Theoretical Biological Physics and
Department of Physics, University of California at San Diego, La Jolla, California
92093-0374, U.S.A., Tel.: (858)534-4844, Email: hlevine@ucsd.edu

Abstract

The survival of cells relies on their ability to respond specifically to diverse environmental signals. Surprisingly, intracellular signaling pathways often share the same or homologous protein components, yet undesirable crosstalk is in general suppressed. This signaling specificity has been well studied in the yeast model system *Saccharomyces cerevisiae*, where the mitogen activated protein kinase (MAPK) cascades are repeatedly employed in mediating distinct biological processes including pheromone-induced mating and filamentous growth under starvation. Although various mechanisms have been proposed to interpret the yeast MAPK signaling specificity, a consistent theory is still lacking. Here, we present a mathematical model that shows signaling specificity can arise through *asymmetric hierarchical inhibition*. The parameters of our model are, where possible, based on experimental data which allow us to determine the constraints imposed by signaling specificity on these parameters. Our model is in broad agreement with experimental observations to date and generates testable predictions that may stimulate further research.

Key words: signaling specificity; MAPK cascade; asymmetric hierarchical inhibition

Introduction

One of the central questions for biological information processing is how cells sense and distinguish extracellular stimuli when intracellular signaling components are shared among pathways. This sharing of components could potentially lead to erroneous cross-activation where the “wrong” downstream pathway is activated following an external stimulus. To maintain signaling specificity cells have evolved a variety of different mechanisms (1). An obvious mechanism that leads to specificity is to physically separate the pathways. This can be done by either sequestration, where key components of the pathways bind to scaffolding proteins, or compartmentalization where the shared component is restricted to a small portion of the cell.

In some cases, however, signaling specificity is achieved in different ways. An example can be found in the budding yeast *Saccharomyces cerevisiae*, where common protein kinases are activated during the mating pheromone signaling, the switch to filamentous growth, and the response to osmotic stress (1). Fig. 1 A shows the interaction between the mating and filamentous growth pathways, both of which are well characterized through extensive experimental investigations (2–4). To illustrate the potential for cross-talk between these two pathways we have colored shared components in yellow. The mating pathway is activated by the α -pheromone receptor which leads to the dissociation of G protein into $G\alpha$ and $G\beta\gamma$ subunits. The latter recruits the scaffold protein Ste5 to the plasma membrane, where the MAPKKK Ste11, the MAPKK Ste7, and the MAPK Fus3 are sequentially activated with the aid of upstream kinases including Ste20 and Ste50. MAPK Fus3 is Ste5-dependent and plays a central role in mating because it controls not only pheromone-dependent gene expression, but also promotes cell cycle arrest and mating projection (5). Among many downstream targets, Fus3 activates the transcription factor Ste12 in the nucleus by phosphorylating two redundant Ste12 inhibitors, Dig1 and Dig2 (6). Activated Ste12 proteins appear to act as dimers and drive the *pheromone response elements* (PREs) of inducible mating genes (7, 8).

The filamentous growth pathway in yeast cells is responsible for the switch to an invasive growth form under nitrogen starvation and shares many components with the mating pathway (9–11). In particular, it also possesses a MAPK cascade with identical MAPKKK and MAPKK components but with a different MAPK (Kss1). A major difference, however, is that the scaffold protein Ste5 is not needed for the activation of the filamentation-specific MAPK Kss1. Kss1 activates Ste12 as well as the filamentation-specific transcription factor Tec1; active Ste12 and Tec1 bind cooperatively to drive the

filamentation response elements (FREs) (12). Remarkably, Tec1 is degraded rapidly in the presence of active Fus3 after pheromone exposure, providing a unilateral inhibition to avoid cross-activation (13–15). When such negative effect of Fus3 is eliminated by mutants expressing stable Tec1, Fus3 appears to be as potent as Kss1 to activate the filamentation program upon pheromone stimulation, indicating that Fus3 has dual regulatory functions in filamentous growth (14). Finally, both pathways induce the gene *STE12*, and hence Ste12 is upregulated by itself; also, there is a positive feedback loop that upregulates Tec1, as *TEC1* is among filamentation-specific genes (16).

A variety of mechanisms at the MAPK cascade level have been proposed to interpret the specificity in yeast signaling. For example, the scaffold protein Ste5 has long been thought to promote signaling specificity by sequestration (17, 18). Even though some experiments suggest that specificity is promoted by the selective activation of Ste5 (19), there are several arguments why specificity cannot arise at the MAPK cascade level alone but needs to involve mechanisms operating at the gene transcription level. First of all, after pheromone stimulation the MAPKK Ste7 not only phosphorylates Fus3 but also Kss1. Activated Kss1 will phosphorylate Tec1, resulting in filamentation gene expression. How can the downstream gene regulatory circuit filter out the information transmitted by Kss1 after pheromone treatment? Secondly, although only Kss1 is phosphorylated under nutrient deprivation, Kss1 activates both Ste12 and Tec1. The activation of Ste12 is likely to drive undesirable mating-specific genes through its positive feedback loop. Again, how is the downstream specificity achieved when Ste12 could leak to the mating program?

The first question can be answered by the inhibitory function of Fus3 on Tec1. Experiments have demonstrated that activated Fus3 can suppress the activation of Tec1 by degrading it, thus providing a mechanism that prevents the onset of the filamentation pathway as a result of the activation of Kss1 through the mating signaling (13–15). The solution to the second question is less clear and needs to be sought at the downstream level. In this view, the MAPK module acts like an intermediate circuit that integrates and preprocesses environmental cues, and controls the duration and strength of the intermediate products, Fus3 and Kss1. These products can be regarded as the input for the downstream gene regulatory network.

In this paper, we propose a mathematical model to study how the downstream gene regulatory network can achieve signaling specificity. Our model shows how this signaling property can be achieved through *asymmetric hierarchical inhibition* where the activation of the mating program is determined

upstream from that of the filamentation pathway. An essential ingredient of our model is the degradation of the transcription factor dimers (see Fig. 1 B), without which signaling specificity may be destroyed. In contrast to the *mutual inhibition* mechanism, hierarchical inhibition may give rise to monostable and history-independent output, as demonstrated in our simulations. The rate constants used in our model are mostly derived from experimental data and a careful examination of the parameter space reveals interesting constraints on some key parameters. In general, our simulation results are consistent with known experiments and the model generates several testable experimental predictions.

Model

Our model is restricted to the signaling process and gene regulation in the nucleus, with active MAP kinases Fus3* and Kss1* as the input, where * stands for activated components. The key players in our model are thus the transcription factors Ste12 and Tec1. In the absence of stimulation, Ste12 and Tec1 form inactive complexes with Dig1 and Dig2. Experiments indicate that phosphorylation of Dig1/2 by the MAP kinases decreases their affinity for Ste12 and Tec1. In addition, Dig1 and Dig2 bind cooperatively to Ste12, while Tec1 and Dig2 bind competitively to Ste12 (20). Since Dig1/2 is abundantly present in the nucleus, the above observations imply that Ste12 and Tec1 may exist mainly in three macrostates: inactive complexes (Ste12-Dig1-Dig2 and Tec1-Ste12-Dig1, each with a rough stoichiometry of 1:1:1 (20)), active monomers (Ste12* and Tec1*), and active dimers (Ste12*-Ste12* and Tec1*-Ste12*). Since the level of gene expression for the mating and the filamentation program is proportional to the level of the program-specific transcription factors we will take these active dimers as output variables.

In our model development, we started with a detailed representation of the two pathways (see Fig. 1 A). This network resulted in 8 equations for 8 variables which take the activated Fus3*, denoted here by x , and activated Kss1* (y) as input. A full description of this model can be found in the supporting information. To clarify the basic mechanisms underlying signal specificity, we have reduced this full model to a more amenable four variable model. The procedure employed in this reduction is detailed in the supporting information and involves quasi-steady state approximations. Importantly, the qualitative features of the reduced and detailed model are identical. The reduced model has only four variables: the complexes con-

taining inactive Ste12 (S_0), the active monomer Ste12* (S_1), the complexes containing inactive Tec1 (T_0) and the active monomer Tec1* (T_1). The output of this model, the Ste12*-Ste12* dimer (S_2) and the Ste12*-Tec1* dimer (T_2), can be related to the variables of the model through $S_2 = S_1^2/K_S$ and $T_2 = S_1T_1/K_T(x)$ where K_S and $K_T(x)$ are equilibrium disassociation constants for which the specific expressions can be found in the supporting information.

Fig. 1 B shows a schematic representation of the reduced model. The input x and y facilitate the activation of both S_1 and T_1 , which together can form heterodimers T_2 . Besides, S_1 itself dimerizes to the homodimer S_2 . All forms of Ste12 and Tec1, including the dimers, can degrade. Through gene expression, there is a positive feedback from S_2 to S_0 and from T_2 to T_0 . The figure also highlights the three processes that break the symmetry between the two pathways. These are:

1) The branching from Ste12 to Tec1 via the hetero-dimerization process $S_1 + T_1 \rightleftharpoons T_2$. This process is controlled in our model by the parameter q which is the ratio of the on rates for the heterodimer and homodimer. This competitive interaction is based on experimental results which show that abolishing the interaction of Tec1 with Ste12 eliminates Tec1-mediated gene expression, suggesting that the association of Tec1 with Ste12 is responsible for the filamentation transcription activity (20). The relevant rates for these interactions, however, are less known.

2) The Tec1 degradation which is accelerated in the presence of Fus3* and described as a sigmoidal Fus3*-dependent function with a maximum value of F_λ . This process is based on recent experiments which show that Fus3* phosphorylates Tec1, resulting in its degradation by the SCF^{Cdc4} ubiquitin ligase (14). It has been documented that the half-life of Tec1 is extended at least three-fold in *fus3* null mutants (14, 21), providing us with an initial estimate for the degradation parameter F_λ . In contrast, the Ste12 degradation shows much weaker dependence on the input and is assumed to be constant.

3) The stronger promoter activity of the *TEC1* gene relative to the promoter activity of *STE12*. Like the Tec1 degradation, the promoter activities are mathematically described as the sigmoidal Hill functions each with a maximum of F_T and F_S , respectively (see supporting information). The experimental picture suggests that F_S is smaller than F_T and we have used the parameter range shown in Table 2 of the supporting information.

In addition to these parameters, we have also investigated the dependence of our model on a parameter w that controls the degradation of the dimers. This parameter can take on values between 0 and 1, with 0 corre-

sponding to completely stable dimers and with 1 corresponding to a dimer decay rate equal to that of their monomers. Experimentally, Tec1 appears to interact with Ste12 via its C-terminal, while Fus3 binds directly to the N-terminal residues of Tec1 (13). Thus, the heterodimer Tec1*-Ste12* may be easily accessible to Fus3* and experience Fus3-mediated degradation which, as incorporated in our model, selectively degrades its Tec1* component. Little is known about the stability of the dimers, but we will see in the Results section that this parameter can have significant consequences.

The equations describing our reduced model can be written as:

$$\frac{dS_0}{dt} = -g_S(x, y)S_0 + d_S S_1 - \lambda_S S_0 + \lambda_S S_b P_S(S_2) \quad (1)$$

$$\frac{dS_1}{dt} = g_S(x, y)S_0 - d_S S_1 - \lambda_S S_1 - 2w\lambda_S S_2 - w\lambda_T(0)T_2 \quad (2)$$

$$\frac{dT_0}{dt} = -g_T(x, y)T_0 + d_T T_1 - \lambda_T(x)T_0 + \lambda_T(0)T_b P_T(T_2) \quad (3)$$

$$\frac{dT_1}{dt} = g_T(x, y)T_0 - d_T T_1 - \lambda_T(x)T_1 - w\lambda_T(x)T_2. \quad (4)$$

Here, $g_S(x, y)$ and $g_T(x, y)$ are the input-dependent activation rates with maxima g_{Sm} and g_{Tm} , respectively, d_S and d_T are the deactivation rates, λ_S and $\lambda_T(x)$ are the degradation rates with the latter having a maximum of $F_\lambda \lambda_T(0)$, S_b and T_b are the initial concentrations of Ste12 and Tec1, and $P_S(S_2)$ and $P_T(T_2)$ are the promoter activities for *STE12* and *TEC1* with maxima F_S and F_T , respectively. Explicit expressions for the rates can be found in the supporting information where the parameter values listed in Table 2 are derived, where possible, from experimental data.

These equations were integrated using Matlab 7 (The MathWorks). To assess specificity we employed a two-window simulation protocol. During the first time window x was chosen large while y was set to zero, corresponding to high levels of pheromone and no nutritional deprivation. We have verified that relaxing levels of y in the first window resulted in almost identical results. Note that this is not surprising since only the activation rates for S_0 and T_0 depend on y and saturate for the parameter values we have chosen. During the second time window x was set to zero while y was increased to a high value, mimicking nutritional deprivation in the absence of pheromone stimulation. The duration of both windows was chosen long enough to reach a steady state response.

To quantify the specificity of our model we use two previously introduced metrics (22, 23). The first one measures simply the ratio of the S_2

and T_2 response during each window: $S_X = \ln(S_2|_x/T_2|_x)$ in the first time window and $S_Y = \ln(T_2|_y/S_2|_y)$ in the second. Clearly, signaling specificity is achieved when both S_X and S_Y are positive and large. The second metric is called the signaling fidelity. A large fidelity means that the activation of a pathway through its authentic upstream signal is larger than the activation resulting from cross-talk. For the S_2 signaling it is defined as the ratio of S_2 during the first time window and S_2 during the second window: $V_X = \ln(S_2|_x/S_2|_y)$. Similarly, we define the fidelity for T_2 as $V_Y = \ln(T_2|_y/T_2|_x)$. Note that those metrics are mathematically related: $V_X + V_Y = S_X + S_Y$.

Results

A typical example of a numerical simulation, using the default parameter set $(F_S, F_T, F_\lambda, q, w) = (2, 8, 4, 1, 1)$ together with the parameters of Table 2 in the Supporting Information, is plotted in Fig. 2 A which shows the output of our model (Ste12*-Ste12*, red line and Tec1*-Ste12*, green dashed line) as a function of time. Fig. 2 B shows the specificity measure S_X and S_Y for the time trace in A. We have also computed the fidelity for this time trace and found that $V_X \doteq 2.5$ and $V_Y \doteq 2.0$ at steady state. Thus, our model can exhibit both signaling specificity and fidelity through the appropriate choice of parameters.

Next, we vary two of the five essential numerical parameters, while keeping the remaining three constant. Examples of these simulations are shown in Fig. 3. In the first panel, we show S_X and S_Y as a function of F_S and F_T . The surface is color coded with red corresponding to high values of the metric and blue for the low values. Note that at the intersection between the two surfaces both metrics are around 2, indicating good specificity. In the second panel, we show similar curves for V_X and V_Y . The results are combined in the third panel in which we plot the region where $S_{X,Y}$ is above a threshold value as blue points and where $V_{X,Y}$ above this value as green points. Since experimental data on this threshold is not available we have chosen here the value $\ln(5)$ as an illustrative example. The overlap between the two regions, corresponding to a parameter region with both high specificity and fidelity, is plotted as red points.

We have investigated the specificity and fidelity using all possible combinations of two of the five main parameters. Examples of these parameter sweeps are shown in Fig. 4, A–D. Again, a large region in parameter space exists where both the specificity and fidelity can be achieved. Fig. 4 C shows

the same slice in parameter space as Fig. 4 D but for $w = 0.2$. The overlap region becomes significantly larger for increasing w , illustrating the importance of an appreciable degradation rate for the dimers. We will discuss this in more detail below.

Through extensive numerical simulations we found the following conditions necessary for signaling specificity in our model:

1) F_S , characterizing the feedback strength of S_2 on S_0 , should be chosen to be smaller than F_T which controls the feedback of T_2 on T_0 . This can be seen in Fig. 3 C where the overlap in parameter space is well below the line $F_S = F_T$. Fig. 3 C also shows that F_T needs to have a certain minimum value which is about 5 for our chosen parameter values.

2) The dimers need to have a minimal basal degradation rate. This can be seen in Fig. 4 A where the overlap region shrinks for decreasing values of w and even disappears for $w < 0.1$. It is also shown in Fig. 4, E and F, where we plot S_2 and T_2 as well as the signaling specificity metrics during our two-window numerical experiments for $w = 0$ (completely stable dimers). During the first time window, S_2 is significantly higher than T_2 . However, S_2 remains high in the second window, destroying signaling specificity as negative S_Y shown in Fig. 4 F.

3) For a given set of parameters, we found that the values for q that exhibit signaling specificity and fidelity are bounded by a minimum and a maximum value. This can be seen in Fig. 4, C and D where, for fixed values of F_λ , the red region starts at a smaller value of q and ends at a larger q .

4) The Fus3*-mediated degradation, described by the parameter F_λ , demands a certain minimum value, found to be around 3 in simulations. This is illustrated in Fig. 4, B–D.

In summary, we found that signaling specificity imposed the following parameter constraints: $1 < F_S < 5 < F_T$, $F_\lambda > 3$, $w > 0.1$ and $q_{min} < q < q_{max}$, where q_{min} and q_{max} mostly depend on the choice of w and F_λ .

In addition to these parameter constraints, our simulations also exclude the bistable property, i.e. the existence of two stable steady state solutions for the same set of parameters. Bistability was argued in a recent study to be the consequence of mutual inhibition and the resulting history dependent output was observed in experiments probing the high osmolarity response and pheromone-induced mating in yeast (24).

To understand the absence of multiple stable steady solutions, we further reduce our four variable model by solving Eqs. 1 and 3 for S_0 and T_0 . Substituting the results into Eqs. 2 and 4 yields two nonlinear coupled

ordinary differential equations:

$$\frac{dS_1}{dt} = \frac{d_S S_1 + \lambda_S S_b P_S(S_2)}{1 + \lambda_S/g_S(x, y)} - d_S S_1 - \lambda_S S_1 - 2w\lambda_S S_2 - w\lambda_T(0)T_2 \quad (5)$$

$$\frac{dT_1}{dt} = \frac{d_T T_1 + \lambda_T(0)T_b P_T(T_2)}{1 + \lambda_T(x)/g_T(x, y)} - d_T T_1 - \lambda_T(x)T_1 - w\lambda_T(x)T_2, \quad (6)$$

where, as we have discussed before, S_2 and T_2 are functions of S_1 and T_1 . Note that the only term that couples the two equations is $w\lambda_T(0)T_2$ which arises from the heterodimer degradation. The steady state solution can also be found at the intersection of the nullclines of the above equations, defined as the curves in S_1 vs. T_1 space for which $\frac{dS_1}{dt} = 0$ and $\frac{dT_1}{dt} = 0$. Extensive numerical exploration of parameter space reveals that these curves intersect only once. Thus, there exists a unique steady state which rules out a history dependent trajectory in phase space. The uniqueness of our output can also be seen in Fig. 5 A where we plot S_2 and T_2 as functions of x and y . Obviously, the outputs are single-valued for this representative set of parameters. Our results are markedly different than the ones from a recent theoretical model in the supplementary information of (24), which displays history-dependent bistability between mating and filamentous growth. We will comment further on the differences between the two models in the Discussion section.

To gain insight in the importance of nonvanishing decay of dimers $w > 0$, we then solve the steady equations for S_2 and T_2 using Eqs. 5 and 6:

$$S_2 = \frac{S_b P_S(S_2)}{2w(1 + \lambda_S/g_S)} - (1 + \frac{d_S}{g_S + \lambda_S}) \frac{S_1}{2w} - \frac{\lambda_T(0)}{2\lambda_S} T_2, \quad (7)$$

$$T_2 = \frac{\lambda_T(0)}{\lambda_T(x)} \frac{T_b P_T(T_2)}{w(1 + \lambda_T(x)/g_T)} - (1 + \frac{d_T}{g_T + \lambda_T(x)}) \frac{T_1}{w}. \quad (8)$$

Since $S_1 = \sqrt{K_S S_2}$ and $T_1 = K_T(x)T_2/S_1 = K_T(x)T_2/\sqrt{K_S S_2}$, the above two equations can be rewritten in terms of S_2 and T_2 only:

$$S_2 + (1 + \frac{d_S}{g_S + \lambda_S}) \frac{\sqrt{K_S S_2}}{2w} = \frac{S_2 P_S(S_2)}{2w(1 + \lambda_S/g_S)} - \frac{\lambda_T(0)}{2\lambda_S} T_2, \quad (9)$$

$$T_2 = T_b P_T(T_2) \left[\left(\frac{\lambda_T(x)}{\lambda_T(0)} \left(1 + \frac{\lambda_T(x)}{g_T} \right) (w + (1 + \frac{d_T}{g_T + \lambda_T(x)}) \frac{K_T(x)}{\sqrt{K_S S_2}}) \right) \right]^{-1}. \quad (10)$$

In Eq. 9, the last term represents the draining effect of Tec1 on Ste12 reservoir due to the turnover of Tec1*-Ste12*. This effect ensures that when

y is large and x is very low, the level of S_2 remains low and the mating pathway is suppressed. Note that this term becomes negligible when $w \rightarrow 0$, leading to high levels of S_2 in the presence of large y and small x , and thus to the destruction of specificity for the filamentation pathway.

This draining effect is also controlled by the parameter q which appears implicitly in these equations through $K_T(x)$ since $K_T(x) \sim 1/q$. Therefore, T_2 in equilibrium is proportional to q and choosing large values of q will increase the draining effect. However, even though large q promotes specificity in the filamentation pathway, it may ruin the specificity for the mating pathway. This can be seen by realizing that for large x and small w , the output T_2 is approximately proportional to q/F_λ . Thus, there is a trade-off between the hetero-dimerization rate and the degradation rate of T_2 (see Fig. 4 A), and choosing large values of q may potentially produce high T_2 output even at high concentrations of Fus3*.

Our results can be schematically summarized in the phase diagram shown in Fig. 5 B, where the response of our model in the Fus3*(x)-Kss1*(y) space are represented in a color-coded manner. At low concentrations of Fus3* and Kss1*, the model response is low in both S_2 and T_2 (see also Fig. 5 A). In this blue-colored region, these levels are probably too small to activate either pathway. Upon increasing the level of Kss1* while maintaining a low level of Fus3*, it will activate the filamentous growth pathway (yellow region). Once the Fus3* level has crossed a certain threshold, the mating program is initiated (red region) while the filamentation program is suppressed. This threshold in x is also clear in Fig. 5 A and largely independent of y . Therefore, Fus3 seems to be the determining factor for pathway specificity, consistent with experimental results and previous arguments (14).

Discussion

Mechanisms of our model

The goal of any specificity model for the mating and filamentation pathways should be to give high S_2 (Ste12*-Ste12*) output levels and low T_2 (Tec1*-Ste12*) levels in the presence of high x (Fus3*)/low y (Kss1*) and the reverse for high y /low x input. How is this accomplished in our model? High values of Fus3* lead to increased activation rates of Ste12 but also of Tec1. The latter, however, is suppressed because of the Fus3*-mediated degradation (controlled by F_λ). The result is a high level of S_2 and a low level of T_2 . Note, however, that the hetero-dimerization process, controlled by the parameter q , cannot be too large (see Fig. 4 C). The relatively high

rate of hetero-dimerization, i.e. large q , can potentially overcome the enhanced degradation rates and may bring high T_2 output even in pheromone signaling.

High concentrations of Kss1* in response to nutritional starvation also trigger the activation of Ste12 and Tec1 which further form dimers. The heterodimer T_2 is degraded at a rate controlled by the parameter w , balanced by *de novo* synthesis of Tec1 and new formation of heterodimer. As long as w is larger than a certain threshold, this degradation/synthesis cycle functions as a continuing drain on the Ste12 reservoir, finally reducing the S_2 output. Similarly, q also has a minimum bound (Fig. 4, C and D) below which the hetero-dimerization could be too slow to effectively absorb and drain the Ste12 reservoir. Furthermore, the genetic positive feedback of Tec1, controlled by the parameter F_T , enlarges the Tec1 reservoir and amplifies its draining capacity on Ste12 resource. Consequently, choosing F_T too small will weaken this draining effect and cause erroneous cross-activation. In addition, the positive feedback loop containing S_2 , tuned by F_S , boosts the levels of S_2 . For large values of F_S this feedback can be sufficient to overcome the draining effect and impair specificity. Hence, there is a constraint on the values of F_S , as seen in Fig. 3 C.

From the above, we conclude that the *asymmetric hierarchical* inhibitions in our model are crucial for yeast signaling specificity. *Asymmetric* since the three crucial processes described earlier break the symmetry between the mating and filamentation pathway and *hierarchical* since the inhibitory effect of Fus3* on Tec1 works at a higher level of the pathway and depends only weakly upon the genetic feedback regulation. This is in contrast to the draining effect by the dimer degradation which operates at a pathway level that is further downstream. As a consequence, the decision of yeast cell fate is made hierarchically. At the top of the decision tree, the concentration of Fus3* is evaluated: regardless of the level of Kss1*, high Fus3* will drastically reduce the Tec1 pool and direct the cell to the mating program. For low concentrations of Fus3*, the Fus3-dependent inhibitory effect is weak, and the level of Kss1* is evaluated further down the decision tree: at high Kss1*, the draining effect by Tec1*-Ste12* becomes dominant and the filamentous growth pathway is activated. Our proposed hierarchical inhibition mechanism is different from the mutual inhibition mechanism observed where the inhibition usually occurs at the same level or appears symmetric in the signaling pathways. An example can be found in the yeast osmolar and pheromone responses where the MAPKs Fus3 and Hog1 inhibit each other after activation. This mutual inhibition mechanism may easily lead to bistable behavior as observed recently (24). In contrast, the hierar-

chical inhibition mechanism may not favor bistable property in nature, as characterized in Fig. 5 B.

Comparison with previous model

A recent model for the mating and filamentation pathway relies on a mutual inhibitory interaction that takes into account a drop of Tec1 level in the presence of Fus3* and the draining effect by the hetero-dimerization process (without dimer degradation) on the Ste12 resource (24). There are, however, several significant differences between the two models. First, the parameter values in our model are, where possible, based on experimental data while the parameters in the model in Ref. (24) appear to be chosen more freely. Thus, one should be able to compare the results of our model directly to experimental results. Second, the Tec1 genetic feedback was taken to be linearly dependent on the concentration of Tec1*-Ste12* in their model McClean et al. (24) and could reach arbitrary values in principle. Thus, the draining effect by hetero-dimerization alone can be sustained and unlimited. The inclusion of a more realistic sigmoidal function for this feedback, as presented here, leads to a qualitatively different behavior. Specifically, the draining effect through hetero-dimerization becomes transient and limited when the genetic feedback of Tec1 expression is saturated. Hence, additional mechanism needs to be incorporated to establish a continuing drain on the Ste12 resource. We argue here that the heterodimer degradation might provide such a mechanism to couple with the hetero-dimerization process. As long as this degradation is large enough there will be an outflux from Ste12* to Tec1*-Ste12*, preventing the build-up of Ste12*-Ste12* and ensuring the signaling specificity at high concentrations of Kss1*.

Finally, we have verified through simulations that replacing the sigmoidal feedback in our model with a linear version gives rise to bistable behavior, even when $w = 0$, i.e. dimers are completely stable. Furthermore, we have performed simulations of the model in (24) using our two-window protocol. We found that the existence of bistability depends critically on the strength of the feedback and that reducing this strength more than 3-fold from the parameter value in (24) leads to monostable behavior. Thus, the history-dependent bistability found in (24) is likely a parameter-dependent phenomenon and may not be a natural result of the “cross-pathway mutual inhibition”. This difference in qualitative outcomes should be accessible through experiments as the outcome of the bistable system will be history dependent while our model predicts a unique outcome of the experiments, independent of the input history.

Comparison with experiments

Our model is able to capture a number of experimental observations. First, eliminating the Fus3*-mediated Tec1 degradation, accomplished by setting $F_\lambda = 1$, destroys the specificity in our model. This can be seen for example in Fig. 4 B where the region with both high specificity and fidelity requires a minimum value of F_λ . For reasonably large values of F_T , the abolishment of Fus3*-mediated Tec1* degradation results in the activation of the filamentous growth pathway. This agrees with experiments in which *Tec1*^{T273V} blocks Tec1 ubiquitination and degradation and allows the induction of filamentation genes in response to pheromone (14).

Second, the signaling specificity in our model breaks down for $F_S > F_T$ (see Fig. 3 C). In other words, a requirement for specificity is that the Tec1 genetic feedback is stronger than the Ste12 genetic feedback. This agrees with recent measurements of yeast gene expression, which show that Ste12 is weakly up-regulated by its dimers, while the positive feedback on Tec1 is stronger (25).

Third, the activation of Kss1 is not required to trigger the mating program, as demonstrated in our phase diagram Fig. 5 B. In fact, the level of Fus3* is the determining factor in the cell's fate. This is consistent with experimental documents which suggest that Fus3 is capable of all functions that Kss1 is known of (14). Note that Fus3 has additional mating specific functions, such as pheromone-induced G1 phase arrest, polarity formation for shmooing, and maintaining pathway specificity via degrading Tec1. The additional functions of Fus3 may explain why Fus3 activation has to be under tight control *in vivo* and is strictly dependent on the Ste5 scaffold (26).

Experimental predictions

Several predictions can be made on the basis of our model, some of which may be experimentally accessible. First of all, our model predicts that the stability of heterodimer Tec1*-Ste12* critically affects the level of filamentation gene expression. In our model, this is controlled by the parameter w . Decreasing w significantly elevates the levels of both S_2 and T_2 , while choosing w close to zero endangers signaling specificity (see Fig. 4, E and F). Recent experiments have shown the ability to obtain tunable degradation of a tagged protein through a synthetic gene network in *Saccharomyces cerevisiae* (27). Our predictions about dimer stability may be testable using similar experimental techniques in the future.

Also, overactivation of Kss1 in the presence of pheromone stimulation will not result in filamentous differentiation. This can be most easily seen in the phase diagram which show that even for large values of Kss1* the Fus3* determines the outcome. In addition, overexpression of *STE12* by its transcription homodimer, corresponding to choosing $F_S \gg 1$ in the model, may destroy the filamentation signaling specificity. This is shown in Fig. 3 C where keeping F_T fixed and increasing F_S is demonstrated to lose specificity. Conversely, and also shown in Fig. 3 C, overexpression of *TEC1* by its transcription heterodimer, i.e. $F_T \gg F_S$, may kill the mating signaling specificity.

Finally, we predict a single steady state solution which is reached in a history independent fashion. Hence, the final outcome should be the same regardless of the sequence of external cues given to the cell. Perhaps experiments similar to the ones in (24) are possible to determine the trajectory in phase space and should contribute to our understanding of the underlying mechanisms for eukaryotic signaling specificity.

Supplementary Information

An online supplementary material to this article can be found by visiting BJ Online at <http://www.biophysj.org>.

Acknowledgements

This research has been supported by the NSF-sponsored Center for Theoretical Biological Physics (grant nos. PHY-0216576 and PHY-0225630). We thank Dr. Lee Bardwell for a critical reading of the manuscript and valuable comments.

References

1. Schwartz, M. A., and H. D. Madhani, 2004. Principles of MAP kinase signaling specificity in *Saccharomyces cerevisiae*. *Annu. Rev. Genet.* 38:725–748.
2. Liu, H., C. A. Styles, and G. R. Fink, 1993. Elements of the yeast pheromone response pathway required for filamentous growth of diploids. *Science*. 262:1741–1744.

3. Roberts, R. L., and G. R. Fink, 1994. Elements of a single MAP kinase cascade in *Saccharomyces cerevisiae* mediate two developmental programs in the same cell type: mating and invasive growth. *Genes Dev.* 8:2974–2985.
4. Schaeffer, H. J., and M. J. Weber, 1999. Mitogen-activated protein kinases: specific messages from ubiquitous messengers. *Mol. Cell Biol.* 19:2435–2444.
5. Elion, E. A., 2000. Pheromone response, mating and cell biology. *Curr. Opin. Microbiol.* 3:573–581.
6. Cook, J. G., L. Bardwell, S. J. Kron, and J. Thorner, 1996. Two novel targets of the MAP kinase Kss1 are negative regulators of invasive growth in the yeast *Saccharomyces cerevisiae*. *Genes Dev.* 10:2831–2848.
7. Yuan, Y. L., and S. Fields, 1991. Properties of the DNA-binding domain of the *Saccharomyces cerevisiae* STE12 protein. *Mol. Cell Biol.* 11:5910–5918.
8. Olson, K. A., C. Nelson, G. Tai, W. Hung, C. Yong, C. Astell, and I. Sadowski, 2000. Two regulators of Ste12p inhibit pheromone-responsive transcription by separate mechanisms. *Mol. Cell Biol.* 20:4199–4209.
9. Madhani, H. D., and G. R. Fink, 1998. The control of filamentous differentiation and virulence in fungi. *Trends Cell Biol.* 8:348–353.
10. Pan, X., T. Harashima, and J. Heitman, 2000. Signal transduction cascades regulating pseudohyphal differentiation of *Saccharomyces cerevisiae*. *Curr. Opin. Microbiol.* 3:567–572.
11. Gancedo, J. M., 2001. Control of pseudohyphae formation in *Saccharomyces cerevisiae*. *FEMS Microbiol. Rev.* 25:107–123.
12. Madhani, H. D., and G. R. Fink, 1997. Combinatorial control required for the specificity of yeast MAPK signaling. *Science.* 275:1314–1317.
13. Brückner, S., T. Köhler, G. H. Braus, B. Heise, M. Bolte, and H. U. Mösch, 2004. Differential regulation of Tec1 by Fus3 and Kss1 confers signaling specificity in yeast development. *Curr. Genet.* 46:331–342.
14. Chou, S., L. Huang, and H. Liu, 2004. Fus3-regulated Tec1 degradation through SCFCdc4 determines MAPK signaling specificity during mating in yeast. *Cell.* 119:981–990.

15. Bao, M. Z., M. A. Schwartz, G. T. Cantin, J. R. Yates III, and H. D. Madhani, 2004. Pheromone-dependent destruction of the Tec1 transcription factor is required for MAP kinase signaling specificity in yeast. *Cell*. 119:991–1000.
16. Zeitlinger, J., I. Simon, C. T. Harbison, N. M. Hannett, T. L. Volkert, G. R. Fink, and R. A. Young, 2003. Program-specific distribution of a transcription factor dependent on partner transcription factor and MAPK signaling. *Cell*. 113:395–404.
17. Harris, K., R. E. Lamson, B. Nelson, T. R. Hughes, M. J. Marton, C. J. Roberts, C. Boone, and P. M. Pryciak, 2001. Role of scaffolds in MAP kinase pathway specificity revealed by custom design of pathway-dedicated signaling proteins. *Curr. Biol*. 11:1815–1824.
18. Park, S. H., A. Zarrinpar, and W. A. Lim, 2003. Rewiring MAP kinase pathways using alternative scaffold assembly mechanisms. *Science*. 299:1061–1064.
19. Flatauer, L. J., S. F. Zadeh, and L. Bardwell, 2005. Mitogen-activated protein kinases with distinct requirements for Ste5 scaffolding influence signaling specificity in *Saccharomyces cerevisiae*. *Mol. Cell Biol*. 25:1793–1803.
20. Chou, S., S. Lane, and H. Liu, 2006. Regulation of mating and filamentation genes by two distinct Ste12 complexes in *Saccharomyces cerevisiae*. *Mol. Cell Biol*. 26:4794–4805.
21. Wang, Y., and H. G. Dohlman, 2006. Pheromone-regulated sumoylation of transcription factors that mediate the invasive to mating developmental switch in yeast. *J. Biol. Chem*. 281:1964–1969.
22. Komarova, N. L., X. Zou, Q. Nie, and L. Bardwell, 2005. A theoretical framework for specificity in cell signaling. *Mol. Syst. Biol*. 1:2005.0023.
23. Bardwell, L., X. Zou, Q. Nie, and N. L. Komarova, 2007. Mathematical models of specificity in cell signaling. *Biophys. J*. 92:3425–3441.
24. McClean, M. N., A. Mody, J. R. Broach, and S. Ramanathan, 2007. Cross-talk and decision making in MAP kinase pathways. *Nat. Genet*. 39(Suppl.):409–414.

25. Ren, B., F. Robert, J. J. Wyrick, O. Aparicio, E. G. Jennings, I. Simon, J. Zeitlinger, J. Schreiber, N. Hannett, E. Kanin, T. L. Volkert, C. J. Wilson, S. P. Bell, and R. A. Young, 2000. Genome-wide location and function of DNA binding proteins. *Science*. 290:2306–2309.
26. Andersson, J., D. M. Simpson, M. Qi, Y. Wang, and E. A. Elion, 2004. Differential input by Ste5 scaffold and Msg5 phosphatase route a MAPK cascade to multiple outcomes. *EMBO. J.* 23:2564–2576.
27. Grilly, C., J. Stricker, W. L. Pang, M. R. Bennett, and J. Hasty, 2007. A synthetic gene network for tuning protein degradation in *Saccharomyces cerevisiae*. *Mol. Syst. Biol.* 3:127.

Figure Legends

Figure 1.

(A): Illustration of the mating pathway, initiated by pheromone stimulations, and the filamentous growth pathway, activated by nutrient starvation. Shared components are colored in yellow and details of the interactions can be found in the text. (B): Schematic representation of the core of our model with forward processes indicated in red, backward processes in green and degradation processes in blue. As input, the model takes the $Fus3^*$ and $Kss1^*$ levels. These input variables, called x and y here, activate $Ste12$ and $Tec1$, leading to the dimers $Ste12$ - $Ste12$ (S_2) and $Ste12$ - $Tec1$ (T_2), the output of our model. The specificity determining parameters are indicated and include q , controlling the relative hetero-dimerization rate of T_1 and S_1 ; w , the relative degradation rate of the dimers; F_λ , tuning the $Fus3^*$ -dependent degradation rate of $Tec1$; F_S , adjusting the positive feedback of the gene expression on $Ste12$; and F_T , determining the genetic positive feedback of $Tec1$.

Figure 2.

A typical numerical result using our stimulation protocol. In the first time window, the input to the mating pathway (x) is high while the input to the filamentation pathway (y) is low. During the second window, the input is reversed. The duration of both windows are set to 200min. (A): Time traces of our model output (activated $Ste12^*$ - $Ste12^*$ dimer, S_2 , and the activated $Ste12^*$ - $Tec1^*$ dimer, T_2). (B): Corresponding signaling specificity S_X and S_Y computed from (A).

Figure 3.

The response of our model for varying F_S and F_T . (A): the specificity S_X and S_Y as a function of F_S and F_T . The surface is color coded as indicated by the color bar. (B): the fidelity V_X and V_Y as a function of F_S and F_T . (C): The result of a parameter sweep in F_S vs. F_T space. The parameter combination for which the specificity is larger than $\ln(5)$ is shown as blue points and where the fidelity is larger than $\ln(5)$ is shown as green points. The red points correspond to parameter values for which both the specificity and the fidelity are greater than $\ln(5)$.

Figure 4.

(A-D): The regions of high specificity and fidelity for different parameter sweeps with threshold $\ln(5)$, using the same color coding as in the previous figure. (E-F): The two-window simulation results in the absence of dimer degradation ($w = 0$).

Figure 5.

(A): The output of our model for a representative set of parameters. The value of S_2 and T_2 is uniquely defined for each pair of (x, y) . (B): The schematic phase diagram of our model with blue denoting no response, red corresponding to the activation of the mating pathway, and yellow labeling the activation of filamentous growth.




Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5: