

# Topics in Probability (Math 289A): Lecture 13

Lecturer: Prof. Williams

Scribe: Toni Gui

November 21, 2017

**Skorokhod Problem** Given  $x \in \mathbb{D}_+^d$ , want to find  $(y, z) \in \mathbb{D}_+^d \times \mathbb{D}_+^d$  such that  $z(t) = x(t) + Ry(t) \in \mathbb{R}_+^d \quad \forall t \geq 0$  and  $y(0) = 0, y$  is non-decreasing,  $\int_{[0, \infty)} z(t) dy(t) = 0$ .

If  $R = (v_1 | \cdots | v_d)$  is completely-**S**, then there is a solution of the Skorokhod problem for each  $x \in \mathbb{D}_+^d$ .

Solutions may not be unique even if  $R$  is a **P**-matrix. This non-uniqueness is a problem for "strong" construction of reflecting Brownian motion.

## Weak Formulation for Semimartingale Reflecting Brownian Motion (SRBM)

An SRBM associated with  $R$  and starting from  $x_0 \in \mathbb{R}_+^d$  is a  $d$ -diml continuous adapted process  $Z$  defined on some filtered probability space  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}, \mathbb{P})$  such that  $Z(t) = x_0 + B(t) + RY(t) \in \mathbb{R}_+^d, \forall t \geq 0$ . where

- (i)  $B$  is a standard  $d$ -diml Brownian motion that is an  $\{\mathcal{F}_t\}$ -martingale;
- (ii)  $Y(0) = 0, Y$  is continuous, non-decreasing and  $\{\mathcal{F}_t\}$  adapted,  $\int_{[0, t]} Z(s) dY(s) = 0$ .

**Theorem** (Reiman-Williams, 87; Taylor-Williams, 93) A necessary condition to have an SRBM (in weak sense) for each  $x_0 \in \mathbb{R}_+^d$  is that  $R$  is completely-**S**. If  $R$  is completely-**S**, then for each  $x_0 \in \mathbb{R}_+^d$ , there exists an SRBM (in weak sense) and for such an SRBM with associated  $(B, Y)$ ,  $(Z, B, Y)$  is unique in law.

( A cool theorem proved by Reiman-Williams:  $\int_{[0, t]} 1_{F_i \cap F_j}(Z(s)) dY_i(s) = 0$  a.s.,  $i \neq j, \forall t \geq 0$ , here  $F_i = \{x \in \mathbb{R}_+^d : x_i = 0\}$ .)

*Example of an SDE which has a weak solution but no strong solution:  $dX_t = \text{sgn}(X_t) dB_t$*

here  $\text{sgn}(x) = \begin{cases} 1 & \text{if } x > 0, \\ -1 & \text{if } x \leq 0. \end{cases}$

**Oscillation Inequalities** (Section 3.5 in Skorokhod problem notes) Suppose  $R$  is completely-**S**. Suppose that  $\delta \geq 0, 0 \leq t_1 < t_2 < \infty$  and  $z, x, y \in \mathbb{D}([t_1, t_2], \mathbb{R}^d)$  such that

- (i)  $z(t) = x(t) + Ry(t), t \in [t_1, t_2]$ ,
- (ii)  $z(t) \in \mathbb{R}_+^d, t \in [t_1, t_2]$ ,
- (iii)  $y(t) \geq 0, y$  is non-decreasing,  $\int_{(t_1, t_2]} 1_{(\delta, \infty)}(z_i(s)) dy_i(s) = 0$ .

Then there is a constant  $C > 0$  depending only on  $R$  such that

$$\text{Osc}(y, [t_1, t_2]) \leq C(\text{Osc}(x, [t_1, t_2]) + \delta)$$

$$Osc(z, [t_1, t_2]) \leq C(Osc(x, [t_1, t_2]) + \delta)$$

where, as before.

$$Osc(f, [t_1, t_2]) = \sup\{|f(t) - f(s)| : t_1 \leq s < t \leq t_2\}$$

Can see proof in notes.

**Idea for Proof of Existence of SRBM:** Suppose set  $X(t) = x_0 + B(t)$ , and  $(Z^\delta, Y^\delta)$  is obtained by jumping back into the orthant a distance  $\delta$  whenever a boundary face is reached. From construction,  $(Z^\delta, Y^\delta, X)$  are adapted to filtration generated by  $X$ :  $X$  is a martingale wrt filtration generated by  $(Z^\delta, Y^\delta, X)$ . Let  $\delta_n \rightarrow 0$  as  $n \rightarrow \infty$ , we can use oscillation inequality to prove that  $\{(Z^{\delta_n}, X, Y^{\delta_n})\}_{n=1}^\infty$  is  $C$ -tight:

$$Y^{\delta_n}(T) \leq Osc(Y^{\delta_n}, [0, T]) \leq C(Osc(X, [0, T]) + \delta)$$

Suppose  $(\tilde{Z}, \tilde{X}, \tilde{Y})$  is some weak limit point ( $Z^\delta \rightarrow \tilde{Z}, X \rightarrow \tilde{X}, Y^\delta \rightarrow \tilde{Y}, \tilde{Y}(0) = 0$ ), then  $(\tilde{Z}, \tilde{X}, \tilde{Y})$  is an SRBM.

**Idea for Proof of Uniqueness of SRBM** (Taylor-Williams):

$Z(t) = x_0 + B(t) + RY(t)$ . Do proof by induction on dimension. True for  $d = 1$ . Suppose true for all completely- $\mathbf{S}$   $R$  matrices in dimension  $< d$ . Want to show true for dimension  $d$  (reduce to the case where we start from the origin). Use Brownian scaling and ergodicity of process hitting spherical caps of radius  $2^{-n}$  as it comes out of the origin.

What if  $R$  depends on position?

Weak Uniqueness is still very open other than when have strong spectral radius type condition. (Shashashvili, Dupuis-Ishii—case 2)

Existence: via oscillation inequality (Kang-Williams, 2007, The Annals of Applied Probability, Volume 17, 741-779) (piecewise smooth boundaries)

$R$  is completely- $\mathbf{S} \iff R^T$  is completely- $\mathbf{S}$ .