

MATH180C: Introduction to Stochastic Processes II

[Lecture A00: math-old.ucsd.edu/~ynemish/teaching/180cA](http://math-old.ucsd.edu/~ynemish/teaching/180cA)

[Lecture B00: math-old.ucsd.edu/~ynemish/teaching/180cB](http://math-old.ucsd.edu/~ynemish/teaching/180cB)

Today: Birth and death processes.

Next: PK 6.5

Week 2:

- homework 1 (due Friday April 8)
- NO IN-PERSON LECTURE ON WEDNESDAY

The Yule process

Setting: In a certain population each individual during any (small) time interval of length h gives a birth to one new individual with probability $\beta h + o(h)$, independently of other members of the population. All members of the population live forever. At time 0 the population consists of one individual.

Question: What is the distribution on the size of the population at a given time t ?

The Yule process

Let $X_t, t \geq 0$, be the size of the population at time t .

$X_0 = 1$ (start from one common ancestor).

Compute $\tilde{P}_n(t) = P(X_t = n | X_0 = 1)$

If $X_t = n$, then during a time interval of length h

$$(a) P(X_{t+h} = n+1 | X_t = n) = n\beta h + o(h) \quad (1 - \beta h + o(h))^{n-1}$$

$$(b) P(X_{t+h} = n | X_t = n) = 1 - n\beta h + o(h)$$

$$(c) P(X_{t+h} > n+1 | X_t = n) = o(h)$$

$$(b) P(0 \text{ births} | X_t = n) = (1 - \beta h + o(h))^n = 1 - n\beta h + o(h)$$

← all n indiv. give 0 births

(a), (b), (c) $\Rightarrow (X_t)_{t \geq 0}$ is a pure birth process with rates $\lambda_n = n\beta$

$P_n(t)$ satisfies the system of differential equations

The Yule process

$$(*) \begin{cases} \tilde{P}'_{1,0}(t) = -\beta \tilde{P}_{1,0}(t) & \tilde{P}_1(0) = 1 \\ \tilde{P}'_{2,1}(t) = -2\beta \tilde{P}_{2,1}(t) + \beta \tilde{P}_{1,0}(t) & P_2(0) = 0 \\ \vdots & \vdots \\ \tilde{P}'_{n,n-1}(t) = -n\beta \tilde{P}_{n,n-1}(t) + (n-1)\beta \tilde{P}_{n-2,n-2}(t) & P_n(0) = 0 \\ \vdots & \vdots \\ \vdots & \vdots \end{cases}$$

The same system with shifted indices

$$\tilde{P}_1(t) = P_0(t) \quad \tilde{P}_n(t) = P_{n-1}(t) \quad \text{with } \lambda_n = \beta(n+1)$$

$$P_n(t) = \lambda_0 \cdots \lambda_{n-1} \left(B_{0n} e^{-\lambda_0 t} + \cdots + B_{nn} e^{-\lambda_n t} \right) \quad \lambda_0 \cdots \lambda_{n-1} = \beta^n n!$$

$$B_{kn} = \prod_{\substack{l=0 \\ l \neq k}}^n \frac{1}{\lambda_l - \lambda_k}$$

$$B_{kn} = \prod_{l=0}^{k-1} \frac{1}{\lambda_l - \lambda_k} \prod_{l=k+1}^n \frac{1}{\lambda_l - \lambda_k} = \frac{1}{\beta^n (-1)^k k! (n-k)!}$$

The Yule process

$$P_n(t) = \lambda_0 \cdots \lambda_{n-1} (B_{0n} e^{-\lambda_0 t} + \cdots + B_{nn} e^{-\lambda_n t})$$

$$= \sum_{k=0}^n \cancel{\beta^n} n! \frac{(-1)^k}{\cancel{\beta^n} k! (n-k)!} e^{-\beta(n+1)t}$$

$$= e^{-\beta t} \sum_{k=0}^n \binom{n}{k} (-1)^k (e^{-\beta t})^k$$


$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$$

$$= e^{-\beta t} \sum_{k=0}^n \binom{n}{k} (-e^{-\beta t})^k 1^{n-k} = e^{-\beta t} (1 - e^{-\beta t})^n$$

$$\tilde{P}_n(t) = P_{n-1}(t) = e^{-\beta t} (1 - e^{-\beta t})^{n-1} = p(1-p)^{n-1} \quad p = e^{-\beta t}$$


$$\parallel \\ P(X_t = n) \Rightarrow X_t \sim \text{sGeom}(e^{-\beta t})$$

Graphical representation. Exponential sojourn times




Pure death processes

Pure birth process



What if the chain moves in the opposite direction?



Pure death process:

- exponential sojourn times with rates μ_i
- only negative jumps of magnitude 1 allowed

Pure death processes

Infinitesimal description:

Pure death process $(X_t)_{t \geq 0}$ of rates $(\mu_k)_{k=1}^N$ is a continuous time MC taking values in $\{0, 1, 2, \dots, N-1, N\}$ (state 0 is absorbing) with stationary infinitesimal transition probability functions

$$(a) P_{k, k-1}(h) = \mu_k h + o(h) \quad k=1, \dots, N$$

as $h \rightarrow 0$


$$(b) P_{kk}(h) = 1 - \mu_k h + o(h), \quad k=1, \dots, N$$

$$(c) P_{kj}(h) = 0 \quad \text{for } j > k.$$

State 0 is absorbing ($\mu_0 = 0$)

Pure death process

$$S_k \sim \text{Exp}(\mu_k)$$



Sojourn time / jump description:

Pure death process of rates $(\mu_k)_{k=1}^N$ is a nonincreasing right-continuous process taking values in $\{0, 1, \dots, N\}$

- with sojourn times $S_1, S_2, S_3, \dots, S_N$ being independent exponential r.v.s of rates $\mu_1, \mu_2, \dots, \mu_N$ and
- jumps $X_{W_{i+1}} - X_{W_i} = -1$ of magnitude 1

Differential equations for pure death processes

Define $P_n(t) = P(X_t = n | X_0 = N)$ distribution of X_t
↑ starting in state N

(a), (b), (c) implies (check)

$$\begin{cases} P_n'(t) = -\mu_n P_n(t) + \mu_{n+1} P_{n+1}(t) & \text{for } n = 0 \dots N-1 \\ P_N'(t) = -\mu_N P_N(t) & \end{cases} \quad \text{(note that } \mu_0 = 0)$$

Initial conditions: $P_N(0) = 1$, $P_n(0) = 0$ for $n = 0, \dots, N-1$

Solve recursively: $P_N(t) = e^{-\mu_N t} \rightarrow P_{N-1}(t) \rightarrow \dots \rightarrow P_0(t)$

General solution (assume $\mu_i \neq \mu_j$)

$$P_n(t) = \mu_{n+1} \dots \mu_N \left(A_{n,n} e^{-\mu_n t} + \dots + A_{N,n} e^{-\mu_N t} \right), \quad A_{k,n} = \prod_{\substack{\ell=n \\ \ell \neq k}}^N \frac{1}{\mu_\ell - \mu_k}$$

Linear death process

[Discussion section]

Similar to Yule process:

death rate is proportional to the size of the population

$$\mu_k = dk \quad (\text{linear dependence on } k)$$

Compute $P_n(t)$: • $\mu_{n+1} \cdots \mu_N = \alpha^{N-n} \frac{N!}{n!}$

$$\bullet A_{kn} = \prod_{\substack{e=n \\ e+k}}^N \frac{1}{\mu_e - \mu_k} = \frac{1}{\alpha^{N-n} (-1)^{n-k} (k-n)! (N-k)!}$$

$$\left\{ \begin{array}{l} \mu_e - \mu_k = \alpha(e-k) \end{array} \right.$$

$$\bullet P_n(t) = \alpha^{N-n} \frac{N!}{n!} \cdot \frac{1}{\alpha^{N-n}} \sum_{k=n}^N \frac{1}{(-1)^{n-k} (k-n)! (N-k)!} \cdot e^{-k\alpha t} \quad \left\{ \begin{array}{l} j = k-n \\ k = j+n \end{array} \right.$$


$$= \frac{N!}{n!} \sum_{j=0}^{N-n} \frac{(-1)^j e^{-(j+n)\alpha t}}{j! (N-n-j)!}$$

$$= \frac{N!}{n!} e^{-n\alpha t} \sum_{j=0}^{N-n} \frac{1}{j! (N-n-j)!} (-e^{-\alpha t})^j = \frac{N!}{n! (N-n)!} e^{-n\alpha t} (1 - e^{-\alpha t})^{N-n}$$

$X_t \sim \text{Bin}(N, e^{-\alpha t})$

Interpretation of $X_t \sim \text{Bin}(n, e^{-\alpha t})$ [Discussion section]

Consider the following process: Let $\xi_i, i=1 \dots N$, be i.i.d. r.v.s, $\xi_i \sim \text{Exp}(\alpha)$. Denote by X_t the number of ξ_i 's that are bigger than t (ξ_i is the lifetime of an individual, X_t = size of the population at t). $X_0 = N$.



Then: $S_k \sim \text{Exp}(\alpha k)$, independent

$\hookrightarrow (X_t)_{t \geq 0}$ is a pure death process

Probability that an individual

survives to time t is $e^{-\alpha t}$

Probability that exactly n

individuals survive to time t is

$$\binom{N}{n} e^{-\alpha n t} (1 - e^{-\alpha t})^{N-n} = P(X_t = n)$$