

MATH180C: Introduction to Stochastic Processes II

<https://mathweb.ucsd.edu/~ynemish/teaching/180c>

Today: Stationary distribution and long-
run behavior of CTMC

Next: PK 2.4

Week 4:

- HW3 due Saturday, April 29 on Gradescope
- Midterm 1: Friday, April 28

$$i, j \in \{0, 1, 2, 3, \dots, N\}$$

$$P_{ij}(t) = P(X_t = j \mid X_0 = i)$$

$$P(t) = (P_{ij}(t))$$

$$\frac{d}{dt} P(t) = Q P(t), \quad P(0) = I$$

$$\frac{d}{dt} P(t) = P(t) Q$$

Discrete time (Y_n)

$$P = (P_{ij})$$

$$P_{ij} = P(Y_1 = j \mid Y_0 = i)$$

(P^n) - n-step
transition prob
matrix.

Long run behavior of discrete time MC. Summary

Let $(X_n)_{n \geq 0}$ be a discrete time MC on $\{0, \dots, N\}$ with stationary transition probability matrix $P = (P_{ij})_{i,j=0}^N$.

- P is called **regular** if there exists k such that $[P^k]_{ij} > 0$ for all i, j . [P is regular iff (X_n) is irreducible and aperiodic]

Thm. If P is **regular**, then there exist $\pi_0, \dots, \pi_N \in \mathbb{R}$ s.t.

$$1) \pi_i > 0 \quad \forall i$$

$$2) \sum_{i=0}^N \pi_i = 1$$

$$3) \forall j \quad \lim_{n \rightarrow \infty} P_{ij}^{(n)} = \pi_j$$

(π_0, \dots, π_N) is called limiting

(stationary) distribution of (X_n)

(π_0, \dots, π_N) is uniquely defined by the system of equations

$$\begin{cases} \pi_j = \sum_{i=0}^N \pi_i P_{ij}, \\ \sum_{i=0}^N \pi_i = 1 \end{cases}$$

$$(\pi_0, \pi_1, \dots, \pi_N) = (\pi_0, \dots, \pi_N) P$$

$$P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$


Long run behavior of continuous time MC.

Let $(X_t)_{t \geq 0}$ be a continuous time MC, $X_t \in \{0, \dots, N\}$ and let $(Y_n)_{n \geq 0}$ be the embedded jump chain.

Def. $(X_t)_{t \geq 0}$ is called irreducible if its jump chain $(Y_n)_{n \geq 0}$ is irreducible (consisting of one communicating class)

Thm. If $(X_t)_{t \geq 0}$ is irreducible, then

$$P_{ij}(t) > 0 \text{ for all } i, j \text{ and } t > 0$$



Idea of the proof:

- Y_n is irreducible $\Rightarrow \exists i_1, \dots, i_{k-1}$ s.t.

$$P(Y_k = j, Y_{k-1} = i_{k-1}, \dots, Y_1 = i_1 \mid Y_0 = i) > 0$$


- $P(k\text{-th jump} \leq t < (k+1)\text{-th jump}) > 0 \quad \forall t > 0$

Long run behavior of continuous time MC

Remarks: Continuous time MCs are "aperiodic"

All irreducible continuous time MCs are "regular"

Example.



$(Y_n)_{n \geq 0}$ has period 2

$$P_{00}(t) = P(X_t = 0 | X_0 = 0) > P(S_0 > t) = e^{-t}$$

Thm. If $(X_t)_{t \geq 0}$ is irreducible, then there exists π_0, \dots, π_N

1) $\pi_i > 0, \sum_{i=0}^N \pi_i = 1$

2) $\lim_{t \rightarrow \infty} P_{ij}(t) = \pi_j$ for all i

$$\lim_{t \rightarrow \infty} P(t) = \begin{pmatrix} \pi_0 & \pi_1 & \pi_2 & \dots & \pi_N \\ \pi_0 & \pi_1 & \pi_2 & \dots & \pi_N \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \pi_0 & \pi_1 & \pi_2 & \dots & \pi_N \end{pmatrix}$$

3) $\pi = (\pi_0, \dots, \pi_N)$ is uniquely determined by $\pi Q = 0, \sum_{i=0}^N \pi_i = 1$

π is called limiting/stationary/equilibrium distribution of (X_t)

Long run behavior of continuous time MC $\downarrow_0 P(t)$

Remark about 3): $\pi Q = 0$ is equivalent to $\pi P(t) = \pi \quad \forall t$

(\Rightarrow) If $\pi Q = 0$, then using Kolmogorov backward equation

$$(\pi P(t))' = \pi P'(t) \stackrel{K(B)}{=} \pi Q P(t) = 0$$


so $\pi P(t)$ is independent of t . Since $P(0) = I$, we get

$$\forall t \quad \pi P(t) = \pi P(0) = \pi I = \pi$$

(\Leftarrow) If $\pi P(t) = \pi$, then $(\pi P(t))' = 0$. Using Kolmogorov forward equation

$$0 = (\pi P(t))' = \pi P'(t) = \pi P(t)Q = \pi Q$$

Example: Two-state MC



$$Q = \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}$$

$$(\pi_0, \pi_1) \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} = (0, 0)$$

$$-\pi_0 + \pi_1 = 0$$

$$\pi_0 = \pi_1$$

$$\pi_0 - \pi_1 = 0$$

$$\pi_0 + \pi_1 = 1$$

$$\Rightarrow (\pi_0, \pi_1) = \left(\frac{1}{2}, \frac{1}{2}\right)$$

$$\pi_0 + \pi_1 = 1$$

From Lecture 9: if $Q = \begin{pmatrix} -\alpha & \alpha \\ \beta & -\beta \end{pmatrix}$, then

$$P(t) = I + \frac{1}{\alpha + \beta} Q - \frac{1}{\alpha + \beta} e^{-(\alpha + \beta)t} Q \xrightarrow{t \rightarrow \infty} I + \frac{1}{\alpha + \beta} Q. \text{ If } \alpha = \beta = 1$$

$$\lim_{t \rightarrow \infty} P(t) = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}$$

Note, that the jump process (Y_n) does not have limiting distribution!

$$\tilde{P}^{Y_n} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Long run behavior of discrete time MC. Summary (2)

Let $(X_n)_{n \geq 0}$ be a discrete time MC on $\{0, 1, \dots\}$ with stationary transition probability matrix $P = (P_{ij})_{i,j=0}^{\infty}$

Define $R_i = \min\{n: X_n = i\}$, $m_i = E(R_i | X_0 = i)$ mean duration between visits

Thm. If $(X_n)_{n \geq 0}$ is recurrent irreducible aperiodic, then

$$\lim_{n \rightarrow \infty} P_{ij}^{(n)} = \frac{1}{m_j} \quad \forall j$$

If $\lim_{n \rightarrow \infty} P_{ij}^{(n)} > 0$ for some (all) j , then MC is positive recurrent

$\lim_{n \rightarrow \infty} P_{ij}^{(n)} = 0$ for some (all) j , then MC is null recurrent.

If (X_n) is positive recurrent, $(\pi_j)_{j=0}^{\infty}$, $\pi_j = \lim_{n \rightarrow \infty} P_{ij}^{(n)}$ is called stationary distribution, uniquely determined by

$$\pi_j = \sum_{i=0}^{\infty} \pi_i P_{ij} \quad \forall j, \quad \sum_{i=0}^{\infty} \pi_i = 1, \quad \pi_i > 0$$

Long run behavior of continuous time MC (2)

Let $(X_t)_{t \geq 0}$ be a continuous time MC, $X_t \in \{0, 1, \dots\}$
and let $(Y_n)_{n \geq 0}$ be the embedded jump chain.

Define $R_i = \min \{t > S_0 : X_t = i\}$,

$m_i = E(R_i | X_0 = i)$ - mean return time from i to i

If $m_i < \infty$, then i is positive recurrent (class property).

Thm 1) If $(X_t)_{t \geq 0}$ is irreducible, then

$$\lim_{t \rightarrow \infty} P_{ij}(t) = \frac{1}{q_j m_j} =: \pi_j \geq 0$$

2) $(X_t)_{t \geq 0}$ is positive recurrent iff there exists a (unique)

solution $(\pi'_j)_{j=0}^{\infty}$ to $\sum_{i=0}^{\infty} \pi'_i q_{ij} = 0$, $\sum_{i=0}^{\infty} \pi'_i = 1$, $\pi'_i > 0$

in which case $\pi_j = \pi'_j$ and $(\pi_j)_{j=0}^{\infty}$ is called

limiting/stationary distribution.

Remarks

- 1) Until now we discussed only the transition probabilities. But in order to describe completely MC (X_t) we need also the initial / starting distribution

$$\nu = (\nu_0, \nu_1, \dots), \quad \nu_i = P(X_0 = i)$$

$$(X_t) \longleftrightarrow (\nu, Q)$$

- 2) Distribution of X_{t_1} is given by $\nu P(t_1)$
- $$P(X_{t_1} = i) = [\nu P(t_1)]_i$$

More generally

$$P(X_0 = i_0, X_{t_1} = i_1, \dots, X_{t_n} = i_n) =$$

- 3) Stationary distribution remains unchanged in time
- $$\pi P(t) = \pi \Rightarrow X_0 \sim \pi, \text{ then } X_t \sim \pi \quad P(f) = \pi$$

Remarks

4) Similarly as in the discrete case, π_j gives the fraction of time spent in state j in long run

$$\lim_{T \rightarrow \infty} E \left[\frac{1}{T} \int_0^T \mathbb{1}_{\{X_t = j\}} dt \mid X_0 = i \right] = \pi_j$$

(compare with $\lim_{m \rightarrow \infty} E \left[\frac{1}{m} \sum_{n=0}^{m-1} \mathbb{1}_{\{X_n = j\}} \mid X_0 = i \right] = \pi_j$ for discrete time MC)

5) If we can find $(\pi_i)_{i=0}^{\infty}$ such that $\pi_i q_{ij} = \pi_j q_{ji}$ then $(\pi_i)_{i=0}^{\infty}$ satisfies $\pi Q = 0$

$$\text{Indeed, } \sum_{j=0}^{\infty} \pi_i q_{ij} = \pi_i \sum_{j=0}^{\infty} q_{ij} = 0 = \sum_{j=0}^{\infty} \pi_j q_{ji} = (\pi Q)_i$$

What you should know for midterm 1 (minimum):

- definition of continuous time MC, Markov property, transition probabilities, generator
- representations of MC: infinitesimal (generator), jump-and-hold, transition probabilities, rate diagram and relations between them (in particular Q and $P(t)$)
- computing absorption probabilities and mean time to absorption
- computing stationary distributions for finite and infinite state MCs and interpretation of $(\pi_i)_{i=0}^{\infty}$
- basic properties of birth and death processes