

# BLOW UP FOR THE 2D EULER EQUATION ON SOME BOUNDED DOMAINS

ALEXANDER KISELEV AND ANDREJ ZLATOŠ

ABSTRACT. We find a smooth solution of the 2D Euler equation on a bounded domain which exists and is unique in a natural class locally in time, but blows up in finite time in the sense of its vorticity losing continuity. The domain's boundary is smooth except at two points, which are interior cusps.

## 1. INTRODUCTION

Consider the 2D Euler equation on a bounded domain  $D \subseteq \mathbb{R}^2$ , in vorticity formulation:

$$\omega_t + u \cdot \nabla \omega = 0, \quad \omega(0, \cdot) = \omega_0. \quad (1.1)$$

The incompressible velocity  $u$  satisfies the no-flow condition  $u \cdot n = 0$  on  $\partial D$  (with  $n$  the unit outer normal to  $\partial D$ ), and is found from the vorticity  $\omega$  via the Biot-Savart law

$$u(t, x) = \int_D K_D(x, y) \omega(t, y) dy, \quad (1.2)$$

where the Kernel  $K_D$  is given by  $K_D(x, y) := \nabla^\perp G_D(x, y)$ , with  $\nabla^\perp = (-\partial_{x_2}, \partial_{x_1})$  and  $G_D \geq 0$  the Green's function for  $D$ .

If the domain  $D$  and the initial data  $\omega_0$  are smooth, results on existence, uniqueness, and global regularity of solutions of the 2D Euler equation go back to the 1930s works of Wolibner [15] and Hölder [6]. Even though the solutions in this case are globally regular, infinite-time growth of their derivatives and small scale creation appear to be ubiquitous in two dimensional fluids.

The best known upper bound on the growth of the gradient of the vorticity is double-exponential in time (see [18], even though the result is also implicit in earlier works). The question of how fast it can actually grow has received much attention in the literature. First results of this type were due to Yudovich [16, 17], who constructed examples with unbounded growth of the vorticity gradient on the boundary of the domain. Recently, such growth at the boundary was shown to be fairly generic in [13] and [8]. Better quantitative estimates were obtained by Nadirashvili [14], who built examples with at least linear growth, and Denisov [2], whose construction gives a super-linear lower bound for the vorticity gradient. Denisov also constructed examples with a double-exponential growth for an arbitrary but finite time [3]. In these examples, growth happens in the bulk of the fluid. Very recently, Kiselev and Šverák [7] constructed examples with a double-exponential in time growth of the vorticity gradient on the boundary, which is of course the fastest possible rate of growth. Whether such growth can happen in the bulk remains open. The best result here so far is due to Zlatoš [20], who built examples on the torus with at least exponential in time growth.

The 2D Euler equation in less regular domains and with less regular initial data has also been studied extensively. Yudovich theory [19] (see also [11, 12]) guarantees existence and uniqueness of a bounded weak solution  $\omega$  if the domain is  $C^\infty$  and the initial data  $\omega_0$  is  $L^\infty$ . Uniqueness comes from the fact that the fluid velocity corresponding to a bounded vorticity  $\omega$  is log-Lipshitz. This makes bounded vorticities a very natural class to consider (although incremental generalizations are possible). The smoothness assumptions on the boundary of the domain can also be weakened, and Yudovich theory can be extended to domains which are only  $C^{1,1}$  [5]. Moreover, if the initial data and domain have a higher regularity, the solution inherits it as well [12]. The 2D Euler equation has also been studied on certain types of domains with singular points, including domains with corners. We refer to the recent works [1, 4, 5, 9, 10] for some results on existence, uniqueness, and regularity of solutions, as well as for further references.

Our main purpose in this note is to provide an example showing that if the boundary of the domain is not everywhere sufficiently regular, then much more dramatic effects than just a fast growth of the derivatives of solutions can take place. We show that there are solutions of the 2D Euler equation on domains with two cusps (and smooth elsewhere) which are smooth locally in time but blow up in finite time, in the sense of the vorticity losing continuity. We note that our construction will rely only on the above existence and uniqueness results for smooth domains and initial data, due to a special structure of our example.

The example involves a “double stadium” domain. Let

$$\tilde{D}_r^\pm := [(-r, r) \times (\pm 1 - r, \pm 1 + r)] \cup B_r(-r, \pm 1) \cup B_r(r, \pm 1)$$

be the “stadium” domain in  $\mathbb{R}^2$  with width  $2r > 0$  and centered at  $(0, \pm 1)$ . While  $\tilde{D}_r^\pm$  are defined explicitly, they are only  $C^{1,1}$ . Let us therefore define domains  $D_r^\pm$  by making the connections between the circular arcs and the horizontal intervals (which form  $\partial\tilde{D}_r^\pm$ ) infinitely smooth. This can be done by adjusting the circular arcs near the points of contact with the intervals, and such an adjustment can be made as small as we want in  $C^1$ . Let us do this so that again all the  $D_r^\pm$  are rescaled copies of each other, centered at  $(0, \pm 1)$ . Finally, let

$$D' := D_1^+ \cup D_1^- \cup [(-1, 1) \times \{0\}]$$

and denote  $D := D_1^+$  the upper half of  $D'$ .

If  $\omega_0 \in L^\infty(D')$  is odd in  $x_2$ , then the Yudovich theory shows that there is a unique odd-in- $x_2$  weak solution  $\omega$  to (1.1) on  $D' \times \mathbb{R}$ , namely the one whose restriction to  $D \times \mathbb{R}$  solves (1.1) there. Moreover, if  $\omega_0 \in C(D')$ , then it follows from Yudovich theory that  $\omega(t, \cdot) \in C(D)$  for all  $t > 0$ . The following theorem provides a necessary and sufficient condition for  $\omega$  to remain continuous on the whole domain  $D'$ .

**Theorem 1.1.** *Let  $\omega_0 \in C(\bar{D}')$  be non-negative on  $D$  and odd in  $x_2$ . Then the unique odd-in- $x_2$  solution  $\omega$  to (1.1) on  $D' \times \mathbb{R}$  is continuous for all  $t > 0$  if and only if  $\omega_0(\partial D) = 0$ .*

*Remark.* If in addition we have  $\omega_0 \in C^\infty(\bar{D})$ , then classical regularity theory gives us  $\omega(t, \cdot) \in C^\infty(D)$  for all  $t > 0$ . If furthermore  $\omega_0$  is supported away from  $[-1, 1] \times \{0\}$ , then finite speed of propagation (i.e., bounded  $u$ ) implies  $\omega(t, \cdot) \in C^\infty(D')$  for all small enough  $t > 0$ . By Theorem 1.1, such a solution will become discontinuous in finite time unless  $\omega_0$  vanishes on all of  $\partial D$ .

We note that our primary goal here is to demonstrate existence of the phenomenon of blow up for the Euler equation on some two-dimensional domains. We therefore choose to do this in the shortest way possible, using the double stadium domain  $D'$  with two cusps, instead of writing a more general and more technical paper. It is however not difficult to see that our result extends to many other domains with the same symmetry as  $D'$ , and one cusp would in fact be sufficient. Indeed, for the latter, one can use an extension of the Yudovich theory to domains  $D$  with  $\partial D$  smooth except at one point, which is a convex corner with a right angle (see, e.g., [10] and the references therein). This  $D$  can then replace our stadium domain, and the corresponding  $D'$  would be a symmetric cardioid-shaped domain with its boundary smooth except at a single cusp.

## 2. PROOF OF THEOREM 1.1

If  $\omega_0(\partial D) = 0$ , then the result follows from Yudovich theory because both  $D$  and  $\partial D$  are invariant under the flow map generated by  $u$ .

Let us now assume that  $\omega_0(x_0) > 0$  for some  $x_0 \in \partial D$ . We will show that if  $X'(t; x) = u(t, X(t; x))$  and  $X(0; x) = x$  for  $x \in \bar{D}$ , then there is  $T > 0$  such that  $X(T; x_0) = 0$ . This, the continuity of  $\omega_0$  and of  $X(t; \cdot)$  on  $\bar{D}$ , and oddness of  $\omega$  in  $x_2$  then show that  $\omega(T, \cdot)$  is discontinuous at the origin.

Let  $\delta > 0$  be such that  $|\omega_0^{-1}([\delta, \infty))| \geq 30\delta (> (4 + 2\pi)2\delta + \delta)$ . Because  $u$  is incompressible and  $\omega(t, \cdot) \geq 0$  on  $D$ , it follows that for each  $t \geq 0$ ,

$$|\{y \in D_{1-2\delta}^+ \mid \omega(t, y) \geq \delta\}| \geq \delta. \tag{2.1}$$

Next let

$$\kappa := \inf_{x \in \partial D_{1-\delta}^+ \text{ \& } y \in \bar{D}_{1-2\delta}^+} G_D(x, y).$$

Continuity of  $G_D$  away from the diagonal  $x = y$ , positivity of  $G_D$  on  $D$ , and compactness of  $\bar{D}_r^+$  show that  $\kappa > 0$ . Let  $v > 0$  solve  $\Delta v = 0$  on  $D \setminus \bar{D}_{1-\delta}^+$  with  $v = 0$  on  $\partial D$  and  $v = \kappa$  on  $\partial D_{1-\delta}^+$ . By the Hopf Lemma it follows that  $\varepsilon := \inf_{x \in \partial D} |\nabla v(x)| > 0$ .

We then have  $G_D(x, y) \geq v(x)$  for  $x \in D \setminus \bar{D}_{1-\delta}^+$  and  $y \in \bar{D}_{1-2\delta}^+$ , because the inequality holds for  $x \in \partial(D \setminus \bar{D}_{1-\delta}^+)$  (and both functions are harmonic in  $x$ ). So we have  $|K_D(x, y)| \geq \varepsilon$  for  $x \in \partial D$  and  $y \in \bar{D}_{1-2\delta}^+$ . This and (2.1) show that for  $x \in \partial D$ , the right-hand side of (1.2) equals  $a(t, x)(n_2(x), -n_1(x))$ , with  $a(t, x) \geq \varepsilon\delta^2$ . Thus  $X(t; x_0)$  travels clockwise along  $\partial D$  with velocity  $\geq \varepsilon\delta^2$ , which means that  $X(T; x_0) = 0$  for some  $T > 0$ . The proof is finished.

**Acknowledgements.** AK acknowledges partial support by NSF grants DMS-1104415 and DMS-1159133, and by a Guggenheim Fellowship. AZ acknowledges partial support by NSF grants DMS-1056327 and DMS-1159133.

## REFERENCES

- [1] C. Bardos, F. Di Plinio, and R. Temam, *The Euler equations in planar nonsmooth domains*, J. Math. Anal. Appl. **407** (2013), 69–89.
- [2] S. Denisov, *Infinite superlinear growth of the gradient for the two-dimensional Euler equation*, Discrete Contin. Dyn. Syst. A **23** (2009), 755–764.
- [3] S. Denisov, *Double-exponential growth of the vorticity gradient for the two-dimensional Euler equation*, Proc. Amer. Math. Soc. **143** (2015), 1199–1210.

- [4] F. Di Plinio and R. Temam, *Grisvard's shift theorem near  $L^\infty$  and Yudovich theory on polygonal domains*, SIAM J. Math. Anal. **47** (2015), 159–178.
- [5] D. Gérard-Varet and C. Lacave, *The two-dimensional Euler equations on singular domains*, Arch. Ration. Mech. Anal. **209** (2013), 131–170.
- [6] E. Hölder, *Über unbeschränkte Fortsetzbarkeit einer stetigen ebenen Bewegung in einer unbegrenzten inkompressiblen Flüssigkeit (German)*, Math. Z. **37** (1933), 727–738.
- [7] A. Kiselev and V. Šverák, *Small scale creation for solutions of the incompressible two dimensional Euler equation*, Ann. of Math (2) **180** (2014), 1205–1220.
- [8] H. Koch, *Transport and instability for perfect fluids*, Math. Ann. **323** (2002), 491–523.
- [9] C. Lacave, *Uniqueness for two dimensional incompressible ideal flow on singular domains*, SIAM J. Math. Anal., to appear. (arXiv:1109.1153)
- [10] C. Lacave, E. Miot and C. Wang, *Uniqueness for the 2D Euler equations on domains with corners*, Indiana Univ. Math. J. **63** (2014), 1725–1756.
- [11] A. Majda and A. Bertozzi, *Vorticity and Incompressible Flow*, Cambridge University Press, 2002.
- [12] C. Marchioro and M. Pulvirenti, *Mathematical Theory of Incompressible Nonviscous Fluids*, Applied Mathematical Sciences, **96**, Springer-Verlag, New York, 1994.
- [13] A. Morgulis, A. Shnirelman and V. Yudovich, *Loss of smoothness and inherent instability of 2D inviscid fluid flows*, Comm. Partial Differential Equations **33** (2008), no. 4-6, 943–968.
- [14] N. S. Nadirashvili, *Wandering solutions of the two-dimensional Euler equation (Russian)*, Funktsional. Anal. i Prilozhen. **25** (1991), 70–71; translation in Funct. Anal. Appl. **25** (1991), 220–221.
- [15] W. Wolibner, *Un théorème sur l'existence du mouvement plan d'un fluide parfait, homogène, incompressible, pendant un temps infiniment long (French)*, Mat. Z. **37** (1933), 698–726.
- [16] V. I. Judovič, *The loss of smoothness of the solutions of Euler equations with time (Russian)*, Dinamika Splošn. Sredy **16** (1974), 71–78.
- [17] V. I. Yudovich, *On the loss of smoothness of the solutions of the Euler equations and the inherent instability of flows of an ideal fluid*, Chaos **10** (2000), 705–719.
- [18] V. I. Yudovich, *The flow of a perfect, incompressible liquid through a given region*, Sov. Phys. Dokl. **7** (1963), 789–791.
- [19] V. I. Yudovich, *Non-stationary flows of an ideal incompressible fluid*, Zh. Vych. Mat. **3** (1963), 1032–1066.
- [20] A. Zlatoš, *Exponential growth of the vorticity gradient for the Euler equation on the torus*, Adv. Math **268** (2015), 396–403.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TX 77005, USA  
 EMAIL: KISELEV@RICE.EDU

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 53706, USA  
 EMAIL: ZLATOS@MATH.WISC.EDU